

Introducing

AFEXON[®] Tab.

Characteristics


- Aju Pharm's self-manufactured IMD
- The first Aceclofenac+Eperisone HCl combination in Korea
- Superior pain relief compared to Aceclofenac 100mg and comparable safety profile.



9.8 x 9.8 x 5.4mm

1. Upper: Eperisone HCl 75mg SR layer
2. Lower: Aceclofenac 100mg IR layer

AFEXON Tab showed superior pain reduction compared to Aceclofenac monotherapy in patients with acute low back pain.




*p-value < 0.0001
 † LS Mean change difference = [change in test group] - [change in reference group]
 ‡ p-value(ANCOVA)

Study design

Subject patient: 421 patients with acute low back pain accompanied by musculoskeletal muscle spasms.
 Test drug: AFEXON Tab, | Reference drug: Aceclofenac 100mg | Test method: One tablet per dose, twice daily for 7 days, administered orally after meals.
 Endpoint: 100mm Pain VAS change | Superiority criterion: Upper limit of the 95% CI is less than 0.
 † LS Mean change difference = [change in test group] - [change in reference group]
 ‡ p-value(ANCOVA): P-value for the difference in change between the two groups adjusted for baseline values.

AFEXON Tab significantly reduced the change in total ODI score compared to Aceclofenac monotherapy in patients with acute low back pain.




*p-value < 0.0001

Study design

Subject patient: 421 patients with acute low back pain accompanied by musculoskeletal muscle spasms.
 Test drug: AFEXON Tab, | Reference drug: Aceclofenac 100mg | Test method: One tablet per dose, twice daily for 7 days, administered orally after meals.
 Endpoint: ODI (Oswestry Disability Index)
 † LS Mean change difference = [change in test group] - [change in reference group]
 ‡ p-value(ANCOVA): P-value for the difference in change between the two groups adjusted for baseline values.

AFEXON Tab showed no significant difference in safety outcomes compared to Aceclofenac monotherapy.



Study design

Subject patient: 421 patients with acute low back pain accompanied by musculoskeletal muscle spasms.
 Test drug: AFEXON Tab, | Reference drug: Aceclofenac 100mg | Test method: One tablet per dose, twice daily for 7 days, administered orally after meals.