

HANLIM PHARM PRODUCT LIST

For Human Health and Future



For Human Health and Future
HANLIM PHARM. CO., LTD.

CONTENTS

Introduction	04
History	06
Location	07
Research & Development	08
Export Status & Certificates	09
Products	10

INTRODUCTION



Protecting the precious human life from various forms of disease

Founded in 1974, HANLIM PHARM. CO., LTD. has established itself as one of the foremost pharmaceutical companies in the Republic of Korea. In alignment with its vision of *"Protecting precious human life from various forms of diseases,"* HANLIM has remained grounded in its commitment to advancing public health through substantial investment in research and development, as well as adopting the latest technology and equipment for the production of innovative, high-quality medicines.



HISTORY

About Hanlim

Since its establishment, we have consistently put our entire effort into developing and providing an extensive range of high-quality pharmaceutical products, particularly in IMD and Ophthalmology. We focus on improving patients' lives by developing and supplying impactful products. Equipped with high-tech manufacturing facilities, our plant is designed to produce various dosage forms. With our unwavering commitment to improving every patient's life, we strive to create significant and sustainable value for the future by expanding our research and development pipeline across various therapeutic areas.

1974	Established
1993	Designated as KGMP company (Ministry of Food and Drug Safety)
2000	Established subsidiary company for API, H.L Genomics Co., Ltd.
2003	Achieved one million dollar export medal
2005	Organized ophthalmic division
2007	Achieved three million dollar export medal
2009	Achieved five million dollar export medal
2014	Achieved ten million dollar export medal
2016	Designated as PIC/s GMP company (Ministry of Food and Drug Safety)
2018	Approved for ISO 13485
2020	Approved for ISO 37001
2022	Established subsidiary company SANGMYUNG innovation Co., Ltd.
2025	New manufacturing plant under construction

New Drug Award



LODIEN Tab. KFDA Technology Award 2007	LODIEN Tab. IR52 Jang Young-Shil Award 2008	RISENEX-PLUS Tab. KFDA Technology Award 2011	NASAFLEX Nasal Spray KFDA Technology Award 2014
BRONPASS Tab. KFDA Technology Award 2022	LUMINOMARK Inj. KFDA Technology Award 2024	LUMINOMARK Inj., LUMINOIMAGER-C IR52 Jang Young-Shil Award 2025	

Korea Innovative Pharmaceutical Company



2012 Selected as K-Inno Pharm	2015 Selected as K-Inno Pharm	2018 Selected as K-Inno Pharm	2021 Selected as K-Inno Pharm
2024 Selected as K-Inno Pharm			

LOCATION

| Head Office



- **Address :**
42, Seocho-daero 52-gil, Seocho-gu,
Seoul, Republic of Korea

| Manufacturing Plant



- **Address :**
2-27, Yeongmun-ro, Cheoin-gu,
Yongin-si, Gyeonggi-do, Republic of Korea
- **KGMP Approval**
- **PIC/s GMP Approval / ISO13485**
- **Approved Dosage Forms :**
Tablets, Capsules, Injection, Ophthalmic solution,
Ointment, Nasal spray, Internal solution

| H.L. Genomics



- **Address :**
1552, Bogaewonsam-ro, Wonsam-myeon,
Cheoin-gu, Yongin-si, Gyeonggi-do, Republic of Korea
- **BGMP Approval**
- **PMDA Approval**
- **Products :** 21 kinds of APIs

| SANGMYUNG innovation Co., Ltd.,



- **Address :**
2-27, Yeongmun-ro, Cheoin-gu,
Yongin-si, Gyeonggi-do, Republic of Korea
- **Products :** Ocular Implants, Microarray Patches,
Ophthalmic Drops

RESEARCH & DEVELOPMENT

Thinking of human health and the future, HANLIM PHARM. CO., LTD. R&D Centers are equipped with advanced technologies and staffed by dedicated researchers who focus on developing innovative products, including not only modified new drugs but also new chemical entities (NCEs). By continuing to invest in research and development, Hanlim has been recognized by the government's Korea Health Industry Development Institute (KHDI) as one of Korea's Best Innovative Pharmaceutical Companies since 2012, acknowledged for its high R&D investment-to-revenue ratio and outstanding performance.



PERFORMANCE

PERFORMANCE		API	Indication	Clinical study	Remark (Approval date)
	Lodien Tab.	S-amlodipine nicotinate	Hypertension	Phase 1, 3, 4	Chiral, New Salt (JUL, 2006)
	Risenex Plus Tab.	Risedronate +Vit D3	Osteoporosis	Phase 3, 4	<u>Once a week</u> New Comb. (JUN, 2010)
	Risenex M Tab.	Risedronate +Vit D3	Osteoporosis	Phase 1, 3, 4	<u>Once a month</u> New Comb. (APR, 2012)
	Nasaflex Nasal Spray	Mometasone furoate +Azelastine HCl	Allergic rhinitis	Phase 3, 4	New Comb. (NOV, 2012)
	Belion Tab.	Bepotastine salicylate	Allergic rhinitis	Phase 1	New salts (MAR, 2013)
	T-sporin Eye Drops	Cyclosporin	Dry eye	Phase 3	New Formulation (MAY, 2014)
	Lodivikar Tab.	S-amlodipine +Olmesartan	Hypertension	Phase 1	New Chiral Combination (SEP, 2014)
	Belion CR Tab.	Bepotastine salicylate	Allergic rhinitis	Phase 1, 3	Controlled Release (JUL, 2018)
	Stafen Cap.	Pitavastatin Ca +Fenofibrate	Dyslipidemia	Phase 1, 3	New Combination (APR, 2019)
	Bronpass Tab.	7 Herbal ingredients	Acute bronchitis	Phase 2a, 2b, 3	New Herb Drug (APR, 2021)
	Lodien-T Tab.	S-amlodipine + Telmisartan	Hyertension	Phase 1	New Chiral Combination (MAY, 2021)
	Luminomark inj.	Indocyanine Green	Surgery marker	Phase 2, 3	New Indication (MAY, 2023)

EXPORT STATUS


HANLIM PHARM. CO., LTD. has been making countless efforts to supply quality products to the world for decades. Supported by our qualified global capabilities, we dedicate significant time and energy to align our GMP standards with advanced and global requirements. Now, our products are being exported to over 35 countries around the world.



Region	Countries
Asia	Brunei, Cambodia, Hong Kong, Indonesia, Macau, Malaysia, Mongolia, Myanmar, Philippines, Singapore, Sri Lanka, Taiwan, Thailand, Vietnam
Africa	Ethiopia, Nigeria
MENA	Iran, Iraq, Saudi Arabia, Syria, United Arab Emirates (UAE)
North America	United States
CIS	Azerbaijan, Georgia, Uzbekistan
LATAM	Bolivia, Costa Rica, Dominican Republic, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama, Peru +

CERTIFICATES

PIC/s GMP Certificate



ISO 13485 Certificate



PRODUCTS

Gastrointestinal Drug	11
Cardiovascular Related Agents	12
Hemostatics, Anticoagulants and Related Agents	17
Allergy & ENT Disease Related Agents	18
Anti-inflammatory & Analgesics, Anti-rheumatics	20
Narcotic Analgesics	25
Anaesthetics & Adjuvants	28
Hypnotics	29
Skeletal Muscle Relaxants	30
Endocrine & Metabolism Related Agents	30
Ophthalmologic Drugs	33
Anti-epileptics	49
Psychoneural Drugs	50
Corticosteroids	50
Urological Agent	52
Diagnosis Agent & Diagnosis NIR Imaging Device	52

※ The dosage and administration may vary depending on the symptoms/age of patient.
For more detailed information, please refer to the insert paper.

Gastrointestinal Drug

ZOPYRIN Enteric-coated Tab.

Packing : 30Tabs/BTL, 100Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years



Composition

Each tablet contains
Sulfasalazine 500mg

Indications

- Rheumatoid arthritis resistant to NSAIDs
- Ulcerative colitis


Dosage & Administration

- Rheumatoid arthritis:
Initial: 1 tablet tablet 1 time a day
Maintain: 1 tablet 2~4 times a day

PEPTAZOLE Tab. (20mg)

Packing : 28Tabs/BTL

Shelf life : 3 years



Composition

Each tablet contains
Pantoprazole sodium sesquihydrate 22.55mg (Pantoprazole 20mg)

Indications

- Mild gastro-esophageal reflux disease and associated symptoms
- Long-term management and prevention of relapse in reflux esophagitis
- Prevention of gastro-esophageal ulcers caused by NSAIDs in patients who should consume NSAIDs consistently


Dosage & Administration

- 1 tablet 1 time a day before breakfast

PEPTAZOLE Tab. (40mg)

Packing : 28Tabs/BTL

Shelf life : 3 years



Composition

Each tablet contains
Pantoprazole sodium sesquihydrate 45.1mg (Pantoprazole 40mg)

Indications

- Combination with antibiotic therapy for the prevention of recurrent H.pylori associated gastric and duodenal ulcers
- Gastric and duodenal ulcer
- Moderate to severe esophageal reflux
- Zollinger-Ellison syndrome and other pathological gastric acid excess

Dosage & Administration

- 1 tablet 1~2 times a day
* Patient with severe liver impairment: Maximum 20mg 1 time a day

Gastrointestinal Drug

DAGES Cap.

Packing : 10Caps/PTP, 20Caps/PTP, 500Caps/BTL

Shelf life : 2 years



Composition	Each capsule contains Pepsin 25mg, Papain 50mg, Diastase 15mg, Cellulase 15mg, Pancreatin 50mg, Pancrelipase 13mg, UDCA 25mg
Indications	- Dyspepsia, anorexia (loss of appetite), overeating, upset stomach, abdomen bloating due to dyspepsia - Supplements for the following diseases: Liver diseases due to lack of bile secretion, disease in biliary tract and gall
Dosage & Administration	1~2 capsules 3 times a day after a meal

Cardiovascular Related Agents

LODIEN Tab. (1.25mg)

Packing : 30Tabs/PTP, 300Tabs/BTL

Shelf life : 3 years



Composition	Each tablet contains S-Amlodipine nicotinate 1.627mg (S-Amlodipine 1.25mg)
Indications	- Cardiac ischemia caused by hypertension, chronic stable angina of the coronary arteries or vasospasm (variant angina) of coronary vascular system - Reduce risks of angina and coronary artery revascularization in CAD patients without cardiac failure or with not less than 40% of cardiac output
Dosage & Administration	2.5mg as S-amlodipine 1 time a day (Maximum: 5mg a day)

LODIEN Tab. (2.5mg)

Packing : 30Tabs/PTP, 300Tabs/BTL

Shelf life : 3 years




Composition	Each tablet contains S-Amlodipine nicotinate 3.253mg (S-Amlodipine 2.5mg)
Indications	- Cardiac ischemia caused by hypertension, chronic stable angina of the coronary arteries or vasospasm (variant angina) of coronary vascular system - Reduce risks of angina and coronary artery revascularization in CAD patients without cardiac failure or with not less than 40% of cardiac output
Dosage & Administration	2.5mg as S-amlodipine 1 time a day (Maximum: 5mg a day)

LODIEN Tab. (5mg)

Packing : 30Tabs/PTP, 300Tabs/BTL

Shelf life : 3 years




Composition	Each tablet contains S-Amlodipine nicotinate 6.506mg (S-Amlodipine 5mg)
Indications	- Cardiac ischemia caused by hypertension, chronic stable angina of the coronary arteries or vasospasm (variant angina) of coronary vascular system - Reduce risks of angina and coronary artery revascularization in CAD patients without cardiac failure or with not less than 40% of cardiac output
Dosage & Administration	2.5mg as S-amlodipine 1 time a day (Maximum: 5mg a day)

Cardiovascular Related Agents

LODIEN-T Tab. (40/2.5mg)

Packing : 30Tabs/BTL

Shelf life : 3 years



Composition Each tablet contains
Telmisartan 40mg, S-Amlodipine nicotinate 3.253mg (S-Amlodipine 2.5mg)


Indications • Essential hypertension that is not adequately controlled by S-amlodipine or telmisartan monotherapy

Dosage & Administration • 1 tablet 1 time a day with water; with or without a meal
* Recommend to take at the same time every day

LODIEN-T Tab. (40/5mg)

Packing : 30Tabs/BTL

Shelf life : 3 years



Composition Each tablet contains
Telmisartan 40mg, S-Amlodipine nicotinate 6.506mg (S-Amlodipine 5.0mg)


Indications • Essential hypertension that is not adequately controlled by S-amlodipine or telmisartan monotherapy

Dosage & Administration • 1 tablet 1 time a day with water; with or without a meal
* Recommend to take at the same time every day

LODIEN-T Tab. (80/2.5mg)

Packing : 30Tabs/BTL

Shelf life : 3 years



Composition Each tablet contains
Telmisartan 80mg, S-Amlodipine nicotinate 3.253mg (S-Amlodipine 2.5mg)


Indications • Essential hypertension that is not adequately controlled by S-amlodipine or telmisartan monotherapy

Dosage & Administration • 1 tablet 1 time a day with water; with or without a meal
* Recommend to take at the same time every day

LODIEN-T Tab. (80/5mg)

Packing : 30Tabs/BTL

Shelf life : 3 years



Composition Each tablet contains
Telmisartan 80mg, S-Amlodipine nicotinate 6.506mg (S-Amlodipine 5.0mg)

Indications • Essential hypertension that is not adequately controlled by S-amlodipine or telmisartan monotherapy

Dosage & Administration • 1 tablet 1 time a day with water; with or without a meal
* Recommend to take at the same time every day

Cardiovascular Related Agents

LODIVIKAR Tab. (2.5/20mg)

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains S-Amlodipine 2.5mg, Olmesartan medoxomil 20mg
	Indications	• Essential hypertension in patients whose blood pressure is not adequately controlled by amlodipine (S-amlodipine) or olmesartan medoxomil monotherapy
	Dosage & Administration	• 1 tablet 1 time a day with water; with or without a meal * Recommend to take at the same time every day

LODIVIKAR Tab. (2.5/40mg)

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains S-Amlodipine 2.5mg, Olmesartan medoxomil 40mg
	Indications	• Essential hypertension in patients whose blood pressure is not adequately controlled by amlodipine (S-amlodipine) or olmesartan medoxomil monotherapy
	Dosage & Administration	• 1 tablet 1 time a day with water; with or without a meal * Recommend to take at the same time every day

LODIVIKAR Tab. (5/20mg)

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains S-Amlodipine 5mg, Olmesartan medoxomil 20mg
	Indications	• Essential hypertension in patients whose blood pressure is not adequately controlled by amlodipine (S-amlodipine) or olmesartan medoxomil monotherapy
	Dosage & Administration	• 1 tablet 1 time a day with water; with or without a meal * Recommend to take at the same time every day

LODIVIKAR Tab. (5/40mg)

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains S-Amlodipine 5mg, Olmesartan medoxomil 40mg
	Indications	• Essential hypertension in patients whose blood pressure is not adequately controlled by amlodipine (S-amlodipine) or olmesartan medoxomil monotherapy
	Dosage & Administration	• 1 tablet 1 time a day with water; with or without a meal * Recommend to take at the same time every day

Cardiovascular Related Agents

CRECHECK Tab. (2.5mg)

Packing : 30Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Rosuvastatin calcium 2.6mg (Rosuvastatin 2.5mg)
	Indications	- Dietary supplements for primary hypercholesterolemia (type IIa), complex hyperlipidemia (type IIb), and familial hypercholesterolemia homozygote - Delay the progress of arteriosclerosis by reducing total cholesterol (TC) and LDL-c to the target level in hypercholesterinemia patients - Dietary supplements for primary dysbetalipoproteinemia - Reduce the risk for cardiovascular diseases
	Dosage & Administration	- Initial: 5mg 1 time a day - Maintain: 10mg 1 time a day * Adjust maintenance dose at an interval of 4 weeks or longer (Maximum: 20mg a day)

CRECHECK Tab. (5mg)

Packing : 30Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Rosuvastatin calcium 5.2mg (Rosuvastatin 5mg)
	Indications	- Dietary supplements for primary hypercholesterolemia (type IIa), complex hyperlipidemia (type IIb), and familial hypercholesterolemia homozygote - Delay the progress of arteriosclerosis by reducing total cholesterol (TC) and LDL-c to the target level in hypercholesterinemia patients - Dietary supplements for primary dysbetalipoproteinemia - Reduce the risk for cardiovascular diseases
	Dosage & Administration	- Initial: 5mg 1 time a day - Maintain: 10mg 1 time a day * Adjust maintenance dose at an interval of 4 weeks or longer (Maximum: 20mg a day)

CRECHECK Tab. (10mg)

Packing : 30Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Rosuvastatin calcium 10.4mg (Rosuvastatin 10mg)
	Indications	- Dietary supplements for primary hypercholesterolemia (type IIa), complex hyperlipidemia (type IIb), and familial hypercholesterolemia homozygote - Delay the progress of arteriosclerosis by reducing total cholesterol (TC) and LDL-c to the target level in hypercholesterinemia patients - Dietary supplements for primary dysbetalipoproteinemia - Reduce the risk for cardiovascular diseases
	Dosage & Administration	- Initial: 5mg 1 time a day - Maintain: 10mg 1 time a day * Adjust maintenance dose at an interval of 4 weeks or longer (Maximum: 20mg a day)

CRECHECK Tab. (20mg)

Packing : 30Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Rosuvastatin calcium 20.8mg (Rosuvastatin 20mg)
	Indications	- Dietary supplements for primary hypercholesterolemia (type IIa), complex hyperlipidemia (type IIb), and familial hypercholesterolemia homozygote - Delay the progress of arteriosclerosis by reducing total cholesterol (TC) and LDL-c to the target level in hypercholesterinemia patients - Dietary supplements for primary dysbetalipoproteinemia - Reduce the risk for cardiovascular diseases
	Dosage & Administration	- Initial: 5mg 1 time a day - Maintain: 10mg 1 time a day * Adjust maintenance dose at an interval of 4 weeks or longer (Maximum: 20mg a day)

Cardiovascular Related Agents

CREDOUBLE Tab. (10/5mg)

Packing : 30Tabs/PTP, 100Tabs/BTL

Shelf life : 3 years



Composition Each tablet contains
Ezetimibe 10mg, Rosuvastatin calcium 5.2mg (Rosuvastatin 5mg)

Indications • Primary hypercholesterolaemia

Dosage & Administration • Initial: 10/5mg 1 time a day
Range: 10/5mg~10/20mg a day
* Adjust maintenance dose after checking lipid level in blood at an interval of 4 weeks or longer

CREDOUBLE Tab. (10/10mg)

Packing : 30Tabs/PTP, 100Tabs/BTL

Shelf life : 3 years



Composition Each tablet contains
Ezetimibe 10mg, Rosuvastatin calcium 10.4mg (Rosuvastatin 10mg)

Indications • Primary hypercholesterolaemia

Dosage & Administration • Initial: 10/5mg 1 time a day
Range: 10/5mg~10/20mg a day
* Adjust maintenance dose after checking lipid level in blood at an interval of 4 weeks or longer

CREDOUBLE Tab. (10/20mg)

Packing : 30Tabs/PTP, 100Tabs/BTL

Shelf life : 3 years



Composition Each tablet contains
Ezetimibe 10mg, Rosuvastatin calcium 20.8mg (Rosuvastatin 20mg)

Indications • Primary hypercholesterolaemia

Dosage & Administration • Initial: 10/5mg 1 time a day
Range: 10/5mg~10/20mg a day
* Adjust maintenance dose after checking lipid level in blood at an interval of 4 weeks or longer

Cardiovascular Related Agents

STAFEN Cap.

Packing : 30Caps/PTP, 300Caps/BTL

Shelf life : 3 years



- | | |
|------------------------------------|--|
| Composition | Each tablet contains
Pitavastatin calcium 2mg, Fenofibrate granule 241.51mg (Fenofibrate 160mg) |
| Indications | <ul style="list-style-type: none"> Treatment for complex-dyslipidemia (or combined hyperlipidemia) patients with high risk for Coronary Heart Disease (CHD) whose LDL-Cholesterol level is adequately controlled by pitavastatin 2mg monotherapy, while their triglyceride level remains high and HDL-Cholesterol level remains low |
| Dosage & Administration | <ul style="list-style-type: none"> 1 capsule 1 time a day after a meal |

PITADUCE Tab. (2mg)

Packing : 30Tabs/PTP, 100Tabs/BTL

Shelf life : 3 years



- | | |
|------------------------------------|---|
| Composition | Each tablet contains
Pitavastatin calcium 2mg |
| Indications | <ul style="list-style-type: none"> Primary hypercholesterolemia and mixed dyslipidemia |
| Dosage & Administration | <ul style="list-style-type: none"> 1~2mg 1 time a day (Maximum: 4mg a day) |

PITADUCE Tab. (4mg)

Packing : 30Tabs/PTP, 100Tabs/BTL

Shelf life : 2 years



- | | |
|------------------------------------|---|
| Composition | Each tablet contains
Pitavastatin calcium 4mg |
| Indications | <ul style="list-style-type: none"> Primary hypercholesterolemia and mixed dyslipidemia |
| Dosage & Administration | <ul style="list-style-type: none"> 1~2mg 1 time a day (Maximum: 4mg a day) |

Hemostatics, Anticoagulants and Related Agents

HANLIM HEPARIN SODIUM Inj. (25,000 IU)

Packing : 5mL*10Vials

Shelf life : 3 years



- | | |
|------------------------------------|--|
| Composition | Each vial(5mL) contains
Heparin sodium 25,000 IU |
| Indications | <ul style="list-style-type: none"> Prevention of cerebral thrombosis due to thrombosis/cerebral hemorrhage Prevention and treatment of thrombosis and pulmonary embolism after surgery Prevention and treatment of thromboembolic atrial fibrillation Diagnosis and treatment of acute and chronic blood clotting disorders Prevention and treatment of blood clotting Coronary occlusion/peripheral vascular embolism |
| Dosage & Administration | <ul style="list-style-type: none"> IV injection or mix with perfusing blood at extracorporeal circulation |

Hemostatics, Anticoagulants and Related Agents

HANLIM PROTAMINE SULFATE Inj.

Packing : 5mL*5Amps

Shelf life : 3 years



Composition	Each ampoule(5mL) contains Protamine sulfate 50mg
Indications	- Neutralization of heparin overdose - Neutralization of heparin activity after extracorporeal circulation like hemodialysis, heart-lung machine, selective brain perfusion cooling method, and etc.
Dosage & Administration	1~1.5mL (10~15mg) with respect to 1,000 unit of heparin

Allergy & ENT Disease Related Agents

NASAFLEX Nasal Spray

Packing : 18mL

Shelf life : 3 years



Composition	Each mL contains Mometasone furoate 0.5mg, Azelastine HCl 1.4mg
Indications	Perennial allergic rhinitis
Dosage & Administration	1 actuation in each nostril 2 times a day * Each actuation includes Mometasone furoate 0.05mg and Azelastine HCl 0.14mg

NASATIVE Nasal Spray

Packing : 18g

Shelf life : 2 years



Composition	Each mL contains Mometasone furoate 0.5mg
Indications	- Seasonal or perennial allergic rhinitis - Nasal polyps - Acute rhinosinusitis without severe bacterial infection
Dosage & Administration	Allergic rhinitis: 2 actuations in each nostril 1 time a day (Minimum 1 actuation, maximum 4 actuations in each nostril a day)

DESONA Nasal Solution

Packing : 10mL

Shelf life : 2 years



Composition	Each mL contains Budesonide 1mg
Indications	- Seasonal and perennial allergic rhinitis: Allergic or vasomotor rhinitis - Prevention of recurrence of nasal polyps after treatment and incision
Dosage & Administration	Rhinitis: 4 actuations in each nostril 1 time a day or 2 actuations in each nostril 2 times a day

RHINOVENT Nasal Solution

Packing : 15mL

Shelf life : 2 years



Composition	Each mL contains Ipratropium bromide 0.3mg
Indications	Symptomatic relief of rhinorrhea in allergic and non-allergic rhinitis : Runny nose
Dosage & Administration	1 actuation (42µg) in each nostril 2~3 times a day

Allergy & ENT Disease Related Agents

BELION Tab.

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 2 years



Composition Each tablet contains
Bepotastine salicylate 9.64mg (Bepotastine 7.11mg)

Indications

- Perennial allergic rhinitis
- Chronic urticarial
- Pruritus accompanied by skin diseases (eczema, dermatitis, skin itchiness, prurigo)

Dosage & Administration

- 1 tablet 2 times a day

BELION CR Tab.

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years



Composition Each tablet contains
Bepotastine salicylate 19.28mg (Bepotastine 14.22mg)

Indications

- Perennial allergic rhinitis
- Chronic urticarial
- Pruritus accompanied by skin diseases (eczema, dermatitis, skin itchiness, prurigo)

Dosage & Administration

- 1 tablet 1 time a day after a meal

QUINOVID Otic Solution

Packing : 5mL

Shelf life : 3 years



Composition Each mL contains
Ofloxacin 3mg

Indications

- Infection by this drug-resistant bacteria such as staphylococcus, streptococcus, haemoproteus, pseudomonas aeruginosa, and influenza: Tympanitis and external otitis

Dosage & Administration

- 6~10 drops 2 times a day
- * Lay the head by side and stay for 10 mins

BRONPASS Tab.

Packing : 30Tabs/PTP, 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years



Composition Each tablet contains
Mixture of dried rehmannia glutinosa · Paeonia suffruticosa andrews · Five-flavor-fruit · Asparagus cochinchinensis MERR · Scutellaria baicalensis root · Apricot stone · Stomona radix extract (1.4~1.7→1) · Corn starch (4.8:1) 300mg

Indications

- Acute bronchitis

Dosage & Administration

- 1 tablet 2 times a day

Anti-inflammatory & Analgesics, Anti-rheumatics

CELE-V Cap. (100mg)

Packing : 30Caps/BTL, 300Caps/BTL

Shelf life : 3 years

	Composition	Each capsule contains Celecoxib 100mg
	Indications	• Symptomatic relief of osteoarthritis (degenerative arthritis), rheumatoid arthritis and ankylosing spondylitis • For the relief of acute pain (after surgery or tooth extraction) • Primary dysmenorrhea
	Dosage & Administration	• Osteoarthritis (Degenerative arthritis): 200mg 1 time a day or 100mg 2 times a day

CELE-V Cap. (200mg)

Packing : 30Caps/BTL, 300Caps/BTL


Shelf life : 3 years

	Composition	Each capsule contains Celecoxib 200mg
	Indications	• Symptomatic relief of osteoarthritis (degenerative arthritis), rheumatoid arthritis and ankylosing spondylitis • For the relief of acute pain (after surgery or tooth extraction) • Primary dysmenorrhea
	Dosage & Administration	• Osteoarthritis (Degenerative arthritis): 200mg 1 time a day or 100mg 2 times a day

MEBUTON Tab.

Packing : 30Tabs/BTL, 500Tabs/BTL


Shelf life : 3 years

	Composition	Each tablet contains Nabumetone 500mg
	Indications	• Inflammation and pain of osteoarthritis (degenerative arthritis), rheumatoid arthritis
	Dosage & Administration	• 2 tablets 1 time a day before bed

GABANEURO Cap. (25mg)

Packing : 30Caps/BTL, 100Caps/BTL

Shelf life : 3 years

	Composition	Each capsule contains Pregabalin 25mg
	Indications	• Peripheral and central neuropathies • Epilepsy: Adjunctive therapy with partial seizures with or without secondary systemic symptoms • Fibromyalgia syndrome
	Dosage & Administration	• Neuropathy: 1) Peripheral - Initial: 150mg in 2 divided doses a day - After 3~7 days: Increase up to 300mg - After: If needed, increase up to 600mg at an interval of 7 days 2) Central - Initial: 150mg in 2 divided doses a day - After 7 days: Increase up to 300mg - After additional 7 days: Increase up to 600mg

Anti-inflammatory & Analgesics, Anti-rheumatics

GABANEURO Cap. (50mg)

Packing : 30Caps/BTL, 100Caps/BTL

Shelf life : 3 years



Composition	Each capsule contains Pregabalin 50mg
Indications	- Peripheral and central neuropathies - Epilepsy: Adjunctive therapy with partial seizures with or without secondary systemic symptoms - Fibromyalgia syndrome
Dosage & Administration	- Neuropathy: - Peripheral - Initial: 150mg in 2 divided doses a day - After 3~7 days: Increase up to 300mg - After: If needed, increase up to 600mg at an interval of 7 days - Central - Initial: 150mg in 2 divided doses a day - After 7 days: Increase up to 300mg - After additional 7 days: Increase up to 600mg

GABANEURO Cap. (75mg)

Packing : 30Caps/BTL, 100Caps/BTL

Shelf life : 3 years




Composition	Each capsule contains Pregabalin 75mg
Indications	- Peripheral and central neuropathies - Epilepsy: Adjunctive therapy with partial seizures with or without secondary systemic symptoms - Fibromyalgia syndrome
Dosage & Administration	- Neuropathy: - Peripheral - Initial: 150mg in 2 divided doses a day - After 3~7 days: Increase up to 300mg - After: If needed, increase up to 600mg at an interval of 7 days - Central - Initial: 150mg in 2 divided doses a day - After 7 days: Increase up to 300mg - After additional 7 days: Increase up to 600mg

GABANEURO Cap. (150mg)

Packing : 30Caps/BTL, 100Caps/BTL

Shelf life : 3 years




Composition	Each capsule contains Pregabalin 150mg
Indications	- Peripheral and central neuropathies - Epilepsy: Adjunctive therapy with partial seizures with or without secondary systemic symptoms - Fibromyalgia syndrome
Dosage & Administration	- Neuropathy: - Peripheral - Initial: 150mg in 2 divided doses a day - After 3~7 days: Increase up to 300mg - After: If needed, increase up to 600mg at an interval of 7 days - Central - Initial: 150mg in 2 divided doses a day - After 7 days: Increase up to 300mg - After additional 7 days: Increase up to 600mg

Anti-inflammatory & Analgesics, Anti-rheumatics

GABANEURO Cab. (300mg)

Packing : 30Caps/BTL, 100Caps/BTL

Shelf life : 3 years

	Composition	Each capsule contains Pregabalin 300mg
	Indications	- Peripheral and central neuropathies - Epilepsy: Adjunctive therapy with partial seizures with or without secondary systemic symptoms - Fibromyalgia syndrome
	Dosage & Administration	- Neuropathy: - Peripheral - Initial: 150mg in 2 divided doses a day - After 3~7 days: Increase up to 300mg - After: If needed, increase up to 600mg at an interval of 7 days - Central - Initial: 150mg in 2 divided doses a day - After 7 days: Increase up to 300mg - After additional 7 days: Increase up to 600mg

GABANEURO ER Tab. (75mg)

Packing : 30Tabs/PTP

Shelf life : 3 years

	Composition	Each tablet contains Pregabalin 75mg
	Indications	Treatment of peripheral neuropathic pain in adults
	Dosage & Administration	- Initial : 150 mg once daily, administered after the evening meal - Increased up to 600 mg per day at intervals of 1 week - Taper gradually with at least 1-week intervals between dose reductions

GABANEURO ER Tab. (150mg)

Packing : 30Tabs/PTP

Shelf life : 3 years

	Composition	Each tablet contains Pregabalin 150mg
	Indications	Treatment of peripheral neuropathic pain in adults
	Dosage & Administration	- Initial: 150 mg once daily, administered after the evening meal - Increased up to 600 mg per day at intervals of 1 week - Taper gradually with at least 1-week intervals between dose reductions

GABANEURO ER Tab. (300mg)

Packing : 30Tabs/PTP

Shelf life : 3 years

	Composition	Each tablet contains Pregabalin 300mg
	Indications	Treatment of peripheral neuropathic pain in adults
	Dosage & Administration	- Initial: 150 mg once daily, administered after the evening meal - Increased up to 600 mg per day at intervals of 1 week - Taper gradually with at least 1-week intervals between dose reductions

Anti-inflammatory & Analgesics, Anti-rheumatics

JAKFAS Tab. (5mg)

Packing : 30Tabs/BTL

Shelf life : 4 years

	Composition	Each tablet contains Tofacitinib 5.0mg
	Indications	• Rheumatoid arthritis • Psoriatic arthritis • Ankylosing spondylitis • Ulcerative colitis
	Dosage & Administration	• Please refer to the insert

JAKFAS Tab. (10mg)

Packing : 30Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Tofacitinib 10.0mg
	Indications	• Ulcerative colitis
	Dosage & Administration	• Please refer to the insert

ENSITRA Tab.

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Tramadol HCl 37.5mg, Acetaminophen 325mg
	Indications	• Acute and chronic pain of moderate to severe level
	Dosage & Administration	• Initial: 2 tablets every 6 hours (Maximum: 8 tablets a day)

ENSITRA Semi Tab.

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Tramadol HCl 18.75mg, Acetaminophen 162.5mg
	Indications	• Acute and chronic pain of moderate to severe level
	Dosage & Administration	• Initial: 4 tablets every 6 hours (Maximum: 16 tablets a day)

ENSITRA ER Tab.

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Tramadol HCl 75mg, Acetaminophen 650mg
	Indications	• Acute and chronic pain of moderate to severe level
	Dosage & Administration	• Initial: 1 tablet every 12 hours (Maximum: 4 tablets a day)

Anti-inflammatory & Analgesics, Anti-rheumatics

ENSITRA Semi ER Tab.

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years



- | | |
|------------------------------------|--|
| Composition | Each tablet contains
Tramadol HCl 37.5mg, Acetaminophen 325mg |
| Indications | <ul style="list-style-type: none"> Acute and chronic pain of moderate to severe level |
| Dosage & Administration | <ul style="list-style-type: none"> Initial: 2 tablets every 12 hours (Maximum: 8 tablets a day) |

R-A Tab. (10mg)

Packing : 30Tabs/BTL, 100Tabs/BTL

Shelf life : 3 years




- | | |
|------------------------------------|---|
| Composition | Each tablet contains
Leflunomide 10mg |
| Indications | <ul style="list-style-type: none"> Treatment of rheumatoid arthritis Symptomatic relief of active psoriatic arthritis |
| Dosage & Administration | <ul style="list-style-type: none"> Loading: 100mg 1 time a day for 3 days Maintain: 20mg a day |

R-A Tab. (20mg)

Packing : 30Tabs/BTL, 30Tabs/AluAlu, 100Tabs/BTL

Shelf life : 3 years



- | | |
|------------------------------------|---|
| Composition | Each tablet contains
Leflunomide 20mg |
| Indications | <ul style="list-style-type: none"> Treatment of rheumatoid arthritis Symptomatic relief of active psoriatic arthritis |
| Dosage & Administration | <ul style="list-style-type: none"> Loading: 100mg 1 time a day for 3 days Maintain: 20mg a day |

ZOPYRIN Enteric-coated Tab.

Packing : 30Tabs/BTL, 100Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years



- | | |
|------------------------------------|---|
| Composition | Each tablet contains
Sulfasalazine 500mg |
| Indications | <ul style="list-style-type: none"> Rheumatoid arthritis resistant to NSAIDs Ulcerative colitis |
| Dosage & Administration | <ul style="list-style-type: none"> Rheumatoid arthritis:
Initial: 1 tablet 1 time a day
Maintain: 1 tablet 2~4 times a day |

Anti-inflammatory & Analgesics, Anti-rheumatics

HALOXIN Tab. (100mg)

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Hydroxychloroquine sulfate 100mg
	Indications	- Rheumatoid arthritis, juvenile rheumatoid arthritis, discoid and systemic lupus erythematosus, and photosensitive dermatological diseases - Prevention and treatment of malaria (P. vivax, P. malariae, P. ovale and P. falciparum)
	Dosage & Administration	Rheumatoid arthritis, juvenile rheumatoid arthritis, discoid and systemic lupus erythematosus, and dermatological conditions caused or aggravated by sunlight: Initial: 200~400mg a day (Divided doses in case of 400mg)

HALOXIN Tab. (200mg)

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Hydroxychloroquine sulfate 200mg
	Indications	- Rheumatoid arthritis, juvenile rheumatoid arthritis, discoid and systemic lupus erythematosus, and photosensitive dermatological diseases - Prevention and treatment of malaria (P. vivax, P. malariae, P. ovale and P. falciparum)
	Dosage & Administration	Rheumatoid arthritis, juvenile rheumatoid arthritis, discoid and systemic lupus erythematosus, and dermatological conditions caused or aggravated by sunlight: Initial: 200~400mg a day (Divided doses in case of 400mg)

Narcotic Analgesics

ULTIAN Inj. (1mg)

Packing : 5Vials

Shelf life : 18 months

	Composition	Each vial contains Remifentanyl HCl 1.10mg
	Indications	- Analgesic agent for use during induction and/or maintenance of general anesthesia - Provision of analgesia in mechanically ventilated intensive care patients
	Dosage & Administration	IV injection: After diluted to 20~250µg/mL

ULTIAN Inj. (2mg)

Packing : 5Vials

Shelf life : 2 years

	Composition	Each vial contains Remifentanyl HCl 2.21mg
	Indications	- Analgesic agent for use during induction and/or maintenance of general anesthesia - Provision of analgesia in mechanically ventilated intensive care patients
	Dosage & Administration	IV injection: After diluted to 20~250µg/mL

ULTIAN Inj. (5mg)

Packing : 5Vials

Shelf life : 3 years

	Composition	Each vial contains Remifentanyl HCl 5.53mg
	Indications	- Analgesic agent for use during induction and/or maintenance of general anesthesia - Provision of analgesia in mechanically ventilated intensive care patients
	Dosage & Administration	IV injection: After diluted to 20~250µg/mL

Narcotic Analgesics

HANLIM FENTANYL Inj. (1mL(A))

Packing : 1mL*10Amps

Shelf life : 3 years



Composition Each mL contains
Fentanyl citrate 78.5µg (Fentanyl 50µg)

Indications

- Short-term analgesic: At time of/before/induction/maintenance of anesthesia, and right after surgery
- Narcotic analgesic adjuvants for systemic or topical anesthesia
- Preanesthetic medication for induction and maintenance of anesthesia, coadministering with neuroleptics, like droperidol
- Anesthetic adjuvants with oxygen in selective high-risk patients: Open-heart surgery, complex neurologic and orthopedic surgery

Dosage & Administration

- Short term analgesic:
IV injection: 50~100µg (1~2mL) 30~60 mins before surgery

HANLIM FENTANYL Inj. (2mL(A))

Packing : 2mL*10Amps

Shelf life : 3 years



Composition Each mL contains
Fentanyl citrate 78.5µg (Fentanyl 50µg)

Indications

- Short-term analgesic: At time of/before/induction/maintenance of anesthesia, and right after surgery
- Narcotic analgesic adjuvants for systemic or topical anesthesia
- Preanesthetic medication for induction and maintenance of anesthesia, coadministering with neuroleptics, like droperidol
- Anesthetic adjuvants with oxygen in selective high-risk patients: Open-heart surgery, complex neurologic and orthopedic surgery

Dosage & Administration

- Short term analgesic:
IV injection: 50~100µg (1~2mL) 30~60 mins before surgery

HANLIM FENTANYL Inj. (10mL(A))

Packing : 10mL*10Amps

Shelf life : 3 years



Composition Each mL contains
Fentanyl citrate 78.5µg (Fentanyl 50µg)

Indications

- Short-term analgesic: At time of/before/induction/maintenance of anesthesia, and right after surgery
- Narcotic analgesic adjuvants for systemic or topical anesthesia
- Preanesthetic medication for induction and maintenance of anesthesia, coadministering with neuroleptics, like droperidol
- Anesthetic adjuvants with oxygen in selective high-risk patients: Open-heart surgery, complex neurologic and orthopedic surgery

Dosage & Administration

- Short term analgesic:
IV injection: 50~100µg (1~2mL) 30~60 mins before surgery

Narcotic Analgesics

HANLIM FENTANYL Inj. (10mL(V))

Packing : 10mL*10Vials

Shelf life : 3 years



Composition Each mL contains
Fentanyl citrate 78.5µg (Fentanyl 50µg)

Indications

- Short-term analgesic: At time of/before/induction/maintenance of anesthesia, and right after surgery
- Narcotic analgesic adjuvants for systemic or topical anesthesia
- Preanesthetic medication for induction and maintenance of anesthesia, coadministering with neuroleptics, like droperidol
- Anesthetic adjuvants with oxygen in selective high-risk patients: Open-heart surgery, complex neurologic and orthopedic surgery

Dosage & Administration

- Short term analgesic:
IV injection: 50~100µg (1~2mL) 30~60 mins before surgery

HANLIM FENTANYL Inj. (20mL(A))

Packing : 20mL*5Amps

Shelf life : 3 years



Composition Each mL contains
Fentanyl citrate 78.5µg (Fentanyl 50µg)

Indications

- Short-term analgesic: At time of/before/induction/maintenance of anesthesia, and right after surgery
- Narcotic analgesic adjuvants for systemic or topical anesthesia
- Preanesthetic medication for induction and maintenance of anesthesia, coadministering with neuroleptics, like droperidol
- Anesthetic adjuvants with oxygen in selective high-risk patients: Open-heart surgery, complex neurologic and orthopedic surgery

Dosage & Administration

- Short term analgesic:
IV injection: 50~100µg (1~2mL) 30~60 mins before surgery

HANLIM FENTANYL Inj. (20mL(V))

Packing : 20mL*5Vials

Shelf life : 3 years



Composition Each mL contains
Fentanyl citrate 78.5µg (Fentanyl 50µg)

Indications

- Short-term analgesic: At time of/before/induction/maintenance of anesthesia, and right after surgery
- Narcotic analgesic adjuvants for systemic or topical anesthesia
- Preanesthetic medication for induction and maintenance of anesthesia, coadministering with neuroleptics, like droperidol
- Anesthetic adjuvants with oxygen in selective high-risk patients: Open-heart surgery, complex neurologic and orthopedic surgery

Dosage & Administration

- Short term analgesic:
IV injection: 50~100µg (1~2mL) 30~60 mins before surgery

Anaesthetics & Adjuvants

ROCNIUM Inj. (2.5mL)

Packing : 2.5mL*10Vials

Shelf life : 3 years



Composition	Each vial(2.5mL) contains Rocuronium bromide 25mg
Indications	• Adjuvants for systemic anaesthesia in order to help tracheal intubation during Rapid/normal Sequence Induction (RSI) and to induce muscular skeleton during surgery • Adjuvants to facilitate tracheal intubation and mechanical ventilation at ICU
Dosage & Administration	• IV injection : 0.6mg/kg~1.0mg/kg

ROCNIUM Inj. (5mL)

Packing : 5mL*10Vials

Shelf life : 3 years



Composition	Each vial(5mL) contains Rocuronium bromide 50mg
Indications	• Adjuvants for systemic anaesthesia in order to help tracheal intubation during Rapid/normal Sequence Induction (RSI) and to induce muscular skeleton during surgery • Adjuvants to facilitate tracheal intubation and mechanical ventilation at ICU
Dosage & Administration	• IV injection : 0.6mg/kg~1.0mg/kg

BRITURN Inj. (1mL)

Packing : 1mL*10Vials

Shelf life : 3 years



Composition	Each mL contains Sugammadex sodium 108.8mg
Indications	• Reversal of neuromuscular blockade induced by rocuronium or vecuronium
Dosage & Administration	• General reversal: IV injection: 4mg/kg 1 time

BRITURN Inj. (2mL)

Packing : 2mL*10Vials

Shelf life : 3 years



Composition	Each mL contains Sugammadex sodium 108.8mg
Indications	• Reversal of neuromuscular blockade induced by rocuronium or vecuronium
Dosage & Administration	• General reversal: IV injection: 4mg/kg 1 time

BRITURN Inj. (4mL)

Packing : 4mL*10Vials

Shelf life : 3 years



Composition	Each mL contains Sugammadex sodium 108.8mg
Indications	• Reversal of neuromuscular blockade induced by rocuronium or vecuronium
Dosage & Administration	• General reversal: IV injection: 4mg/kg 1 time

Anaesthetics & Adjuvants

BRITURN Inj. (5mL)

Packing : 5mL*10Vials

Shelf life : 3 years



Composition	Each mL contains Sugammadex sodium 108.8mg
Indications	• Reversal of neuromuscular blockade induced by rocuronium or vecuronium
Dosage & Administration	• General reversal: IV injection: 4mg/kg 1 time

BUNOL Inj.

Packing : 1mL*10Amps

Shelf life : 2 years



Composition	Each mL contains Butorphanol tartrate 2mg
Indications	• Treatment of pain required opioid analgesics • Preoperative or pre-anesthetic medication • For the relief of pain during labor
Dosage & Administration	• Pain required opioid analgesics IV injection : 1mg every 3~4 hours as needed IM injection: 2mg every 3~4 hours as needed

Hypnotics

HANLIM DEXTOMINE Inj. (2mL)

Packing : 2mL*5Vials

Shelf life : 2 years



Composition	Each vial contains Dexmedetomidine HCl 236µg (Dexmedetomidine 200µg)
Indications	• For sedation of initially intubated and mechanically ventilated patients during treatment in an intensive care setting • For sedation of non-intubated patients prior to and/or during surgical and other procedures - Monitored Anesthesia Care, MAC - Awake Fiberoptic Intubation, AFI
Dosage & Administration	• Sedation of initially intubated and mechanically ventilated patients during treatment in an intensive care setting: IV injection: loading 1µg/kg for 10~20 mins

ENTOBAR Inj. (100mg)

Packing : 2mL*10Amps

Shelf life : 3 years



Composition	Each ampoule(2mL) contains Pentobarbital sodium 100mg
Indications	• Before anaesthesia or at an introductory phase of systemic anaesthesia • Sedation of anxiety and nervousness, insomnia • Suppression of convulsions
Dosage & Administration	• IV injection: Initial 100mg (Maximum: 500mg) IM injection: 150~200mg 1 time (Maximum: 250mg a day)

Skeletal Muscle Relaxants

MYALON Tab.

Packing : 30Tabs/BTL, 500Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Afloqualone 20mg
	Indications	- Improvement of myotonia due to the following diseases: Shoulder-arm-neck syndrome, lowback pain - Spastic paralysis due to the following diseases: Cerebrovascular disorders, cerebral palsy, spastic spinal paralysis, multiple sclerosis post-traumatic aftereffect and other cerebrospinal diseases
	Dosage & Administration	1 tablet 3 times a day

Endocrine & Metabolism Related Agents

RISENEX-M Tab.

Packing : 1Tab/PTP


Shelf life : 3 years

	Composition	Each tablet contains Sodium risedronate 150mg, Cholecalciferol 0.75mg(30,000 IU)
	Indications	Treatment and prevention of postmenopausal osteoporosis in women
	Dosage & Administration	1 tablet 1 time a month

RISENEX-PLUS Tab.

Packing : 4Tabs/PTP

Shelf life : 2 years

	Composition	Each tablet contains Sodium risedronate 35mg Cholecalciferol conc. 56mg(5,600 IU)
	Indications	- Treatment and prevention of postmenopausal osteoporosis in women - Treatment of osteoporosis in men
	Dosage & Administration	1 tablet 1 time a week

PANORIN Inj.

Packing : 1mL*10Amps

Shelf life : 2 years

	Composition	Each ampoule(1mL) contains Disodium pamidronate 18.75mg (15mg on dried basis)
	Indications	- Tumour-induced hypercalcaemia - Paget's disease of bone, osteolytic lesions and bone pain in patients with bone metastases associated with breast cancer or multiple myeloma
	Dosage & Administration	Paget's disease of bone: 30mg a day for 3 days

URINON Tab.

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Benzbromarone 50mg
	Indications	Improvement of hyperuricemia: Gout, hypertension with hyperuricemia
	Dosage & Administration	Gout: Initial: 1/2~1 tablet 1 time a day Maintain: 1 tablet 1~3 times a day

Endocrine & Metabolism Related Agents

USORIC Tab. (40mg)

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years




Composition	Each tablet contains Febuxostat 40mg
Indications	Treatment of chronic hyperuricaemia in patients with gout
Dosage & Administration	40mg or 80mg 1 time a day with or without a meal

USORIC Tab. (80mg)

Packing : 30Tabs/BTL, 300Tabs/BTL

Shelf life : 3 years



Composition	Each tablet contains Febuxostat 80mg
Indications	Treatment of chronic hyperuricaemia in patients with gout
Dosage & Administration	40mg or 80mg 1 time a day with or without a meal

DEXINEURO Tab.

Packing : 30Tabs/BTL, 100Tabs/BTL

Shelf life : 3 years



Composition	Each tablet contains R-thioctic acid tromethamine 480mg (R-Thioctic acid 300mg)
Indications	To relieve diabetic polyneuropathy
Dosage & Administration	1 tablet 1 time a day with sufficient amount of water about 30 minutes before breakfast.

HANLIM VASOPRESSIN Inj.

Packing : 1mL*10Amps

Shelf life : 3 years



Composition	Each ampoule(1mL) contains Vasopressin (USP) 20 Vasopressin Unit.
Indications	Differential diagnosis of pituitary diabetes insipidus and pituitary or renal diabetes insipidus, removal of enteric gas (abdominal distention, pretreatment in cholecystography and pyelography), emergency care of acute oesophageal variceal bleeding
Dosage & Administration	Pituitary gland diabetes insipidus: SC/IM injection: 2~10 unit as vasopressin 2~3 times a day as needed

TERIPIN Inj.

Packing : 1Vial+1Amp(Diluent)

Shelf life : 3 years



Composition	Each vial contains Terlipressin acetate 1mg	Each ampoule(5mL) contains NaCl 45mg
Indications	- Treatment of esophageal variceal bleeding - Type 1 hepatorenal syndrome	
Dosage & Administration	Esophageal variceal bleeding: Initial: 1~2mg as terlipressine acetate Maintain: 1mg every 4~6 hours for 2~3 days	

Endocrine & Metabolism Related Agents

TERIPIN-S Inj.

Packing : 8.5mL*5Amps

Shelf life : 2 years



- | | |
|------------------------------------|--|
| Composition | Each ampoule(8.5mL) contains Terlipressin acetate 1mg |
| Indications | <ul style="list-style-type: none"> • Treatment of esophageal variceal bleeding • Type 1 hepatorenal syndrome |
| Dosage & Administration | <ul style="list-style-type: none"> • Esophageal variceal bleeding:
Initial: 1~2mg as terlipressine acetate
Maintain: 1mg every 4~6 hours for 2~3 days |

HANLIM CARBETOCIN Inj.

Packing : 1mL*5Amps

Shelf life : 3 years



- | | |
|------------------------------------|--|
| Composition | Each ampoule(1mL) contains Carbetocin 100µg |
| Indications | <ul style="list-style-type: none"> • Prevention of postpartum hemorrhage due to uterine atony following delivery of an infant by Caesarean section under epidural or spinal anaesthesia |
| Dosage & Administration | <ul style="list-style-type: none"> • IV injection: 100µg(1mL) for 1 min 1 time |

TRACSIBAN Inj. (6.75mg)

Packing : 0.9mL*1Vial

Shelf life : 3 years



- | | |
|------------------------------------|--|
| Composition | Each vial(0.9mL) contains Atosiban acetate 7.1577mg (Atosiban 6.75mg) |
| Indications | <ul style="list-style-type: none"> • To delay imminent pre-term birth in pregnant women with: <ul style="list-style-type: none"> - Regular uterine contractions of at least 30 seconds duration at a rate of ≥ 4 per 30 minutes - A cervical dilation of 1 to 3 cm (0-3 cm for nulliparas) and effacement of $\geq 50\%$ - A gestational age from 24 until 33 completed weeks - A normal fetal heart rate |
| Dosage & Administration | <ul style="list-style-type: none"> • 3 consecutive steps: <ol style="list-style-type: none"> 1) IV Bolus of 0.9mL: over 1 min, 6.75mg 2) IV Loading for 3 hours: 24mL/hr, 18mg/hr 3) Continuous IV infusion: 8mL/hr, 6mg/hr |

TRACSIBAN Inj. (37.5mg)

Packing : 5mL*1Vial

Shelf life : 3 years



- | | |
|------------------------------------|--|
| Composition | Each vial(5mL)contains Atosiban acetate 39.765mg (Atosiban 37.5mg) |
| Indications | <ul style="list-style-type: none"> • To delay imminent pre-term birth in pregnant women with: <ul style="list-style-type: none"> - Regular uterine contractions of at least 30 seconds duration at a rate of ≥ 4 per 30 minutes - A cervical dilation of 1 to 3 cm (0-3 cm for nulliparas) and effacement of $\geq 50\%$ - A gestational age from 24 until 33 completed weeks - A normal fetal heart rate |
| Dosage & Administration | <ul style="list-style-type: none"> • 3 consecutive steps: <ol style="list-style-type: none"> 1) IV Bolus of 0.9mL: over 1 min, 6.75mg 2) IV Loading for 3 hours: 24mL/hr, 18mg/hr 3) Continuous IV infusion: 8mL/hr, 6mg/hr |

Ophthalmologic Drugs

NUMAREN Eye Drops

Packing : 15mL

Shelf life : 3 years



Composition	Each mL contains Hypromellose 3mg, Dextran 70 1mg
Indications	• Dry eyes due to deficient tear or mucous secretion
Dosage & Administration	• 1~2 drops several times a day

NUMAREN Eye Drops (Single-dose)

Packing : 0.5mL*30ea

Shelf life : 3 years



Composition	Each mL contains Hypromellose 3mg, Dextran 70 1mg
Indications	• Dry eyes due to deficient tear or mucous secretion
Dosage & Administration	• 1~2 drops several times a day

TEARCLEAN Eye Drops

Packing : 15mL

Shelf life : 2 years



Composition	Each mL contains Sodium chloride 5.5mg, Potassium chloride 1.5mg, Glucose 0.05mg
Indications	• For the relief of ocular discomfort, dryness, fatigue and blurry vision due to contact lenses
Dosage & Administration	• 2~3 drops 5~6 times a day

TEARCLEAN Eye Drops (Single-dose)

Packing : 0.5mL*30ea

Shelf life : 2 years



Composition	Each mL contains Sodium chloride 5.5mg, Potassium chloride 1.5mg, Glucose 0.05mg
Indications	• For the relief of ocular discomfort, dryness, fatigue and blurry vision due to contact lenses
Dosage & Administration	• 2~3 drops 5~6 times a day

CUTIS Eye Drops

Packing : 15mL

Shelf life : 2 years



Composition	Each mL contains Sodium chloride 5.5mg, Potassium chloride 1.5mg
Indications	• For the Relief of ocular fatigue, dryness, discomfort and blurry vision due to contact lenses
Dosage & Administration	• 2~3 drops 5~6 times a day

CUTIS Eye Drops (Single-dose)

Packing : 0.5mL*30ea

Shelf life : 2 years



Composition	Each mL contains Sodium chloride 5.5mg, Potassium chloride 1.5mg
Indications	• For the Relief of ocular fatigue, dryness, discomfort and blurry vision due to contact lenses
Dosage & Administration	• 2~3 drops 5~6 times a day

Ophthalmologic Drugs

HANLIM POVIDONE Eye Drops

Packing : 10mL

Shelf life : 3 years



- | | |
|------------------------------------|---|
| Composition | Each mL contains
Povidone 20mg |
| Indications | • For dry eyes and when wearing hard contact lenses |
| Dosage & Administration | • 2 drops 4~5 times a day |

MAXCLEAR Eye Drops (Single-dose)

Packing : 0.5mL*60ea

Shelf life : 2 years



- | | |
|------------------------------------|---|
| Composition | Each mL contains
Carboxymethylcellulose sodium 5mg |
| Indications | • For the temporary relief of ocular burning, irritation, and discomfort due to dry eyes or exposure to wind or sun |
| Dosage & Administration | • 1~2 drops as needed |

EYEPURE Eye Drops (Single-dose)

Packing : 0.5mL*30ea

Shelf life : 2 years



- | | |
|------------------------------------|---|
| Composition | Each mL contains
Trehalose dihydrate 33.16mg (Trehalose 30.0mg) |
| Indications | • Ocular dryness or fatigue caused by the followings: Wind, smoke, air pollution, dust, dry heat, air-conditioning, flight, and use of computer for long time |
| Dosage & Administration | • 1 drop several times a day |

HYALURON Eye Drops

Packing : 5mL, 10mL

Shelf life : 3 years



- | | |
|------------------------------------|---|
| Composition | Each mL contains
Sodium hyaluronate 1mg |
| Indications | • Treatment for disorder of cornea-conjunctive epithelial cell
- Endogenic disease such as Sjogren's syndrome, mucocutanealocular syndrome (Stevens-Johns syndrome) and dry eye syndrome
- Exogenous disease due to surgical procedures, medicine, trauma and wearing of contact lens |
| Dosage & Administration | • 1 drop 5~6 times a day |

HYALURON Eye Drops (Single-dose)

Packing : 0.5mL*60ea, 0.88mL*30ea

Shelf life : 3 years



- | | |
|------------------------------------|---|
| Composition | Each mL contains
Sodium hyaluronate 1mg |
| Indications | • Treatment for disorder of cornea-conjunctive epithelial cell
- Endogenic disease such as Sjogren's syndrome, mucocutanealocular syndrome (Stevens-Johns syndrome) and dry eye syndrome
- Exogenous disease due to surgical procedures, medicine, trauma and wearing of contact lens |
| Dosage & Administration | • 1 drop 5~6 times a day |

Ophthalmologic Drugs

HYALURON MAX Eye Drops 0.15%

Packing : 5mL

Shelf life : 2 years



Composition Each mL contains
Sodium hyaluronate 1.5mg

Indications

- Treatment for disorder of cornea-conjunctive epithelial cell
- Endogenic disease such as Sjogren's syndrome, mucocutanealocular syndrome (Stevens-Johns syndrome) and dry eye syndrome
- Exogenous disease due to surgical procedures, medicine, trauma and wearing of contact lens

Dosage & Administration

- 1 drop 5~6 times a day

HYALURON MAX Eye Drops 0.15% (Single-dose) **Packing :** 0.5mL*60ea

Shelf life : 2 years



Composition Each mL contains
Sodium hyaluronate 1.5mg

Indications

- Treatment for disorder of cornea-conjunctive epithelial cell
- Endogenic disease such as Sjogren's syndrome, mucocutanealocular syndrome (Stevens-Johns syndrome) and dry eye syndrome
- Exogenous disease due to surgical procedures, medicine, trauma and wearing of contact lens

Dosage & Administration

- 1 drop 5~6 times a day

HYALURON MAX Eye Drops 0.3% (Single-dose) **Packing :** 0.5mL*60ea

Shelf life : 3 years



Composition Each mL contains
Sodium hyaluronate 3mg

Indications

- Treatment for disorder of cornea-conjunctive epithelial cell
- Endogenic disease such as Sjogren's syndrome, mucocutanealocular syndrome (Stevens-Johns syndrome) and dry eye syndrome
- Exogenous disease due to surgical procedures, medicine, trauma and wearing of contact lens

Dosage & Administration

- 1 drop 5~6 times a day

T-SPORIN Eye Drops 0.05% (Single-dose)

Packing : 0.4mL*30ea

Shelf life : 3 years



Composition Each mL contains
Cyclosporin 0.5mg

Indications

- Increased tear production in patients whose tear production is presumed to be suppressed due to ocular inflammation associated with keratoconjunctivitis sicca

Dosage & Administration

- 1 drop 2 times a day every 12 hours

Ophthalmologic Drugs

DIQUASOL Eye Drops

Packing : 5mL

Shelf life : 3 years



Composition Each mL contains
Diquafosol sodium 30mg

Indications • Improvement of symptoms related with dry eyes (keratoconjunctiva epidermal lesion)

Dosage & Administration • 1 drop 6 times a day

DIQUASOL Eye Drops (Single-dose)

Packing : 0.4mL*60ea

Shelf life : 3 years



Composition Each mL contains
Diquafosol sodium 30mg

Indications • Improvement of symptoms related with dry eyes (keratoconjunctiva epidermal lesion)

Dosage & Administration • 1 drop 6 times a day

DIGENTA Eye Drops

Packing : 5mL

Shelf life : 3 years



Composition Each mL contains
Gentamicin sulfate(potency) 3mg, Fluorometholone 1mg

Indications • Anterior infection by gentamicin susceptible bacteria (e.g. bacterial conjunctivitis)
• Anterior inflammation with possibility of bacterial infection (e.g. post-surgery treatment)

Dosage & Administration • Bacterial infection:
1 drop 5 times a day
* Severe cases: increase 1 drop every 1 hour for 1~2 days

DIGENTA Eye Ointment

Packing : 4g

Shelf life : 3 years



Composition Each gram contains
Gentamicin sulfate(potency) 3mg, Fluorometholone 1mg

Indications • Anterior infection by gentamicin susceptible bacteria (e.g. bacterial conjunctivitis)
• Anterior inflammation with possibility of bacterial infection (e.g. post-surgery treatment)

Dosage & Administration • Bacterial infection:
5mm length 3~4 times a day

Ophthalmologic Drugs

FUMELON Eye Drops 0.02%

Packing : 5mL

Shelf life : 3 years



Composition Each mL contains Fluorometholone 0.2mg

Indications • Extraocular inflammatory diseases (blepharitis, conjunctivitis, keratitis, episcleritis, etc.)

Dosage & Administration • 1~2 drops 2~4 times a day

FUMELON Eye Drops 0.1%

Packing : 5mL

Shelf life : 3 years



Composition Each mL contains Fluorometholone 1mg

Indications • Inflammatory diseases of extraocular and anterior segment of eyes (blepharitis, conjunctivitis, keratitis, episcleritis, iritis, iridocyclitis, postoperative inflammation, etc.)

Dosage & Administration • 1~2 drops 2~4 times a day

FUMELON Eye Drops 0.1% (Single-dose)

Packing : 0.4mL*30ea

Shelf life : 3 years



Composition Each mL contains Fluorometholone 1mg

Indications • Inflammatory diseases of extraocular and anterior segment of eyes (blepharitis, conjunctivitis, keratitis, episcleritis, iritis, iridocyclitis, postoperative inflammation, etc.)

Dosage & Administration • 1~2 drops 2~4 times a day

FUMELON-PLUS Ophthalmic Susp.

Packing : 5mL

Shelf life : 3 years



Composition Each mL contains Fluorometholone 1mg, Tetrahydrozoline HCl 0.25mg

Indications • Acute non-infectious conjunctivitis or allergic conjunctivitis with severe edema and bleeding
• Non-infectious inflammation of anterior segment of eyes (panuveitis, episcleritis, scleritis)
• Postoperative inflammation after strabotomy, cataract or glaucoma surgery

Dosage & Administration • Initial 24~48 hours: 1 drop every 1 hour
After: 1 drop 2~3 times a day

Ophthalmologic Drugs

FUMELON-PLUS Ophthalmic Susp.(Single-dose)

Packing : 0.4mL*30ea

Shelf life : 3 years



Composition	Each mL contains Fluorometholone 1mg, Tetrahydrozoline HCl 0.25mg
Indications	• Acute non-infectious conjunctivitis or allergic conjunctivitis with severe edema and bleeding • Non-infectious inflammation of anterior segment of eyes (panuveitis, episcleritis, scleritis) • Postoperative inflammation after strabotomy, cataract or glaucoma surgery
Dosage & Administration	• Initial 24~48 hours: 1 drop every 1 hour • After: 1 drop 2~3 times a day

LOTEPRO Ophthalmic Susp. 0.5%

Packing : 5mL

Shelf life : 2 years



Composition	Each mL contains Loteprednol etabonate 5mg
Indications	• Treatment of steroid responsive inflammatory conditions of seasonal allergic conjunctivitis and giant papillary conjunctivitis • Treatment of postoperative inflammation
Dosage & Administration	• 1~2 drops 4 times a day

LOTEPRO Ophthalmic Susp. 0.5% (Single-dose)

Packing : 0.4mL*30ea

Shelf life : 2 years



Composition	Each mL contains Loteprednol etabonate 5mg
Indications	• Treatment of steroid responsive inflammatory conditions of seasonal allergic conjunctivitis and giant papillary conjunctivitis • Treatment of postoperative inflammation
Dosage & Administration	• 1~2 drops 4 times a day

LOTEPLUS Ophthalmic Susp.

Packing : 5mL

Shelf life : 3 years



Composition	Each mL contains Loteprednol etabonate 5mg, Tobramycin 3mg
Indications	• Treatment of superficial bacterial infection or steroid responsive inflammatory conditions of seasonal allergic conjunctivitis
Dosage & Administration	• 1~2 drops 4 times a day

BROFENAC Eye Drops

Packing : 5mL

Shelf life : 2 years



Composition	Each mL contains Bromfenac sodium hydrate 1mg
Indications	• Symptomatic treatment of inflammatory diseases of anterior- and extra-ocular segments
Dosage & Administration	• 1 drop 2 times a day

BROFENAC Eye Drops (Single-dose)

Packing : 0.4mL*30ea

Shelf life : 27 month



Composition	Each mL contains Bromfenac sodium hydrate 1mg
Indications	• Symptomatic treatment of inflammatory diseases of anterior- and extra-ocular segments
Dosage & Administration	• 1 drop 2 times a day

Ophthalmologic Drugs

OLOPANOL Eye Drops

Packing : 5mL

Shelf life : 3 years



Composition Each mL contains
Olopatadine HCl 1.11mg (Olopatadine 1mg)

Indications • Treatment of symptoms of allergic conjunctivitis

Dosage & Administration • 1 drop 2 times a day every 6~8 hours (Maximum: 4 months if needed)

OLOPANOL Eye Drops (Single-dose)

Packing : 0.4mL*10ea

Shelf life : 2 years



Composition Each mL contains
Olopatadine HCl 1.11mg (Olopatadine 1mg)

Indications • Treatment of symptoms of allergic conjunctivitis

Dosage & Administration • 1 drop 2 times a day every 6~8 hours (Maximum: 4 months if needed)

OLO-ONCE Eye Drops

Packing : 3mL

Shelf life : 2 years



Composition Each mL contains
Olopatadine HCl 2.22mg (Olopatadine 2mg)

Indications • Treatment of symptoms of allergic conjunctivitis

Dosage & Administration • 1 drop 1 time a day

OLO-ONCE Eye Drops (Single-dose)

Packing : 0.4mL*10ea

Shelf life : 2 years



Composition Each mL contains
Olopatadine HCl 2.22mg (Olopatadine 2mg)

Indications • Treatment of symptoms of allergic conjunctivitis

Dosage & Administration • 1 drop 1 time a day

OLO-POWER Eye Drops 0.7%

Packing : 2.5mL

Shelf life : 2 years



Composition Each mL contains
Olopatadine HCl 7.76mg (Olopatadine 7mg)

Indications • Treatment of ocular itching related with allergic conjunctivitis

Dosage & Administration • 1 drop 1 time a day

Ophthalmologic Drugs

OLO-POWER Eye Drops 0.7% (Single-dose)

Packing : 0.4mL*10ea

Shelf life : 2 years



Composition	Each mL contains Olopatadine HCl 7.76mg (Olopatadine 7mg)
Indications	• Treatment of ocular itching related with allergic conjunctivitis
Dosage & Administration	• 1 drop 1 time a day

CLESIN Eye Drops

Packing : 5mL

Shelf life : 3 years



Composition	Each mL contains Sodium cromoglycate 20mg
Indications	• Treatment of allergic conjunctivitis and keratitis
Dosage & Administration	• 1~2 drops 4 times a day

NAZORIN Eye Drops

Packing : 15mL

Shelf life : 2 years



Composition	Each mL contains Pheniramine maleate 3mg, Naphazoline HCl 0.25mg
Indications	• For the temporary relief of ocular redness and irritation
Dosage & Administration	• 1~2 drops 4 times a day

NAZORIN Eye Drops (Single-dose)

Packing : 0.5mL*10ea

Shelf life : 2 years



Composition	Each mL contains Pheniramine maleate 3mg, Naphazoline HCl 0.25mg
Indications	• For the temporary relief of ocular redness and irritation
Dosage & Administration	• 1~2 drops 4 times a day

EYEFEE Eye Drops

Packing : 15mL

Shelf life : 3 years



Composition	Each mL contains Aminocaproic acid 10mg, Neostigmine methylsulfate 0.02mg, Naphazoline HCl 0.02mg, Chlorpheniramine maleate 0.1mg
Indications	• Fatigue of eye, conjunctival congestion, discomfort feelings of eye after swimming or by dust and sweat, ophthalmia by ultraviolet rays and other rays, blepharitis, discomfort feelings in wearing contact lenses, eye itching, dim-sighted
Dosage & Administration	• 2~3 drops 5~6 times a day

Ophthalmologic Drugs

EASY-DROP Eye Drops

Packing : 15mL

Shelf life : 3 years



Composition Each mL contains
Hypromellose 2mg, Glycerin 2mg, Polyethylene glycol 400 10mg

Indications

- For the temporary relief of burning and irritation due to dryness of the eye
- For protection against further irritation

Dosage & Administration

- 1~2 drops as needed

CLEAR-DROP Eye Drops

Packing : 15mL

Shelf life : 3 years



Composition Each mL contains
Tetrahydrozoline HCl 0.5mg, Dextran 70 1mg, Povidone K25 10mg, Polyethyleneglycol 400 10mg

Indications

- For the relief of redness of the eye due to minor eye irritations
- For use as a protectant against further irritation or to relieve dryness of the eye

Dosage & Administration

- 1~2 drops 4 times a day at the maximum

HANLIM ORIGINAL Eye Drops

Packing : 15mL

Shelf life : 3 years



Composition Each mL contains
Tetrahydrozoline HCl 0.5mg

Indications

- For the relief of redness of the eye due to minor eye irritations

Dosage & Administration

- 1~2 drops 2~4 times a day

DAILY-Q Eye Drops

Packing : 15mL

Shelf life : 3 years



Composition Each mL contains
Carboxymethylcellulose 5mg, Glycerin 9mg

Indications

- For the temporary relief of burning, irritation, and discomfort due to dryness of the eye or exposure to wind or sun
- May be used as a protectant against further irritation

Dosage & Administration

- 1~2 drops as needed

Ophthalmologic Drugs

DAILY-Q Eye Drops (Single-dose)

Packing : 0.5mL*30ea

Shelf life : 3 years



Composition

Each mL contains
Carboxymethylcellulose 5mg, Glycerin 9mg

Indications

- For the temporary relief of burning, irritation, and discomfort due to dryness of the eye or exposure to wind or sun
- May be used as a protectant against further irritation

Dosage & Administration

- 1~2 drops as needed

HANLIM RELIEF A.C Eye Drops

Packing : 15mL

Shelf life : 3 years



Composition

Each mL contains
Tetrahydrozoline HCl 0.5mg, Zinc sulfate 2.5mg

Indications

- For the temporary relief of discomfort and redness of the eye due to minor eye irritations

Dosage & Administration

- 1~2 drops (Maximum: 4 times a day)

HANMURINE Eye Drops

Packing : 15mL

Shelf life : 3 years



Composition

Each mL contains
Polyvinyl alcohol 5mg, Povidone 6mg

Indications

- For the temporary relief of burning and irritation due to dryness of the eye
- For use as a protection against further irritation or to relieve dryness of the eye

Dosage & Administration

- 1~2 drops in case of eye irritation

HANLIM MAXIMUM REDNESS RELIEF Eye Drops

Packing : 15mL

Shelf life : 3 years



Composition

Each mL contains
Glycerin 5mg, Naphazoline HCl 0.3mg

Indications

- For the relief of redness of the eye due to minor eye irritations
- For the temporary relief of burning and irritation due to dryness of the eye
- For use as a protectant against further irritation or dryness of the eye

Dosage & Administration

- 1~2 drops 1~4 times a day

Ophthalmologic Drugs

HANLIM CMC Eye Drops

Packing : 15mL

Shelf life : 3 years



Composition Each mL contains
Carboxymethylcellulose sodium 5mg

Indications

- For the temporary relief of burning, irritation, and discomfort due to dryness of the eye or exposure to wind or sun
- May be used as a protectant against further irritation

Dosage & Administration

- 1~2 drops as needed

HANLIM MAX RELIEF Eye Drops

Packing : 15mL

Shelf life : 3 years



Composition Each mL contains
Carboxymethylcellulose sodium 2.5mg

Indications

- As a lubricant to relieve dryness of the eye
- As a protectant against further irritation of the eye
- For the temporary relief of burning, irritation, and discomfort including exposure to wind or sun

Dosage & Administration

- 1~2 drops as needed

HANLIM PVA Eye Drops

Packing : 15mL

Shelf life : 3 years



Composition Each mL contains
Polyvinyl alcohol 14mg

Indications

- Relieves dryness of the eye
- Prevents further irritation

Dosage & Administration

- 1~2 drops as needed

HANLIM LUBRICANT Eye Ointment

Packing : 3.5g

Shelf life : 3 years



Composition Each gram contains
White petrolatum 573mg, Light mineral oil 425mg

Indications

- For the temporary relief of burning, irritation, and discomfort due to dryness of the eye or exposure to wind or sun
- May be used as a protectant against further irritation

Dosage & Administration

- A small amount to the eyelids

Ophthalmologic Drugs

HANLIM LUBRICANT FRESH Eye Ointment

Packing : 3.5g

Shelf life : 3 years



Composition Each gram contains
White petrolatum 830mg, Light mineral oil 150mg

Indications

- For the temporary relief of burning, irritation, and discomfort due to dryness of the eye or exposure to wind or sun
- May be used as a protectant against further irritation

Dosage & Administration

- A small amount to the eyelids several times a day

ENCLE Eye Ointment

Packing : 3.5g

Shelf life : 3 years



Composition Each gram contains
Sodium chloride 50mg

Indications

- For the temporary relief of corneal edema

Dosage & Administration

- A small amount to the eyelids every 3~4 hours

BRIDIN-T Eye Drops 0.15%

Packing : 5mL

Shelf life : 2 years



Composition Each mL contains
Brimonidine tartrate 1.5mg

Indications

- For the reduction of elevated intraocular pressure in patients with open-angle glaucoma or ocular hypertension

Dosage & Administration

- 1 drop 3 times a day every 8 hours

BRIDIN-T Eye Drops 0.15% (Single-dose)

Packing : 0.3mL*60ea

Shelf life : 2 years



Composition Each mL contains
Brimonidine tartrate 1.5mg

Indications

- For the reduction of elevated intraocular pressure in patients with open-angle glaucoma or ocular hypertension

Dosage & Administration

- 1 drop 3 times a day every 8 hours

Ophthalmologic Drugs

BRIDIN-PLUS Eye Drops

Packing : 5mL

Shelf life : 18 months



Composition

Each mL contains
Brimonidine tartrate 2mg, Timolol maleate 6.8mg

Indications

- For the reduction of intraocular pressure in patients with chronic open-angle glaucoma or ocular hypertension who are insufficiently responsive to topical beta-blockers

Dosage & Administration

- 1 drop 2 times a day every 12 hours

BRIDIN-PLUS Eye Drops (Single-dose)

Packing : 0.3mL*60ea

Shelf life : 18 months



Composition

Each mL contains
Brimonidine tartrate 2mg, Timolol maleate 6.8mg

Indications

- For the reduction of intraocular pressure in patients with chronic open-angle glaucoma or ocular hypertension who are insufficiently responsive to topical beta-blockers

Dosage & Administration

- 1 drop 2 times a day every 12 hours

LOWIOP Eye Drops

Packing : 5mL

Shelf life : 2 years



Composition

Each mL contains
Dorzolamide HCl 22.26mg, Timolol maleate 6.83mg

Indications

- For the reduction of elevated intraocular pressure in patients with open-angle glaucoma or ocular hypertension who are insufficiently responsive to beta-blockers

Dosage & Administration

- 1 drop 2 times a day

PROTAN Eye Drops

Packing : 2.5mL

Shelf life : 3 years



Composition

Each mL contains
Latanoprost 0.05mg

Indications

- For the reduction of intraocular pressure in the following cases: Open-angle glaucoma, chronic angle-closure glaucoma, ocular hypertension

Dosage & Administration

- 1 drop 1 time a day

Ophthalmologic Drugs

PROTAN-PLUS Eye Drops

Packing : 2.5mL

Shelf life : 2 years



Composition Each mL contains
Latanoprost 0.05mg, Timolol maleate 6.83mg

Indications

- For the reduction of elevated intraocular pressure in patients with open-angle glaucoma or ocular hypertension

Dosage & Administration

- 1 drop 1 time a day

DACLE Eye Drops

Packing : 10mL

Shelf life : 3 years



Composition Each mL contains
Chloramphenicol (potency) 5mg, Dexamethasone disodium phosphate 1mg,
Tetrahydrozoline HCl 0.25mg

Indications

- Acute & chronic keratitis and conjunctivitis with heavy swelling and superficial congestion by allergy and inflammation, iritis, iridocyclitis, scleritis episcleritis bullosa

Dosage & Administration

- 1 drop 4 times a day

CALTE Eye Drops 2%

Packing : 5mL

Shelf life : 3 years



Composition Each mL contains
Carteolol HCl 20mg

Indications

- Glaucoma and ocular hypertension


Dosage & Administration

- 1 drop 2 times a day

ACETAZOL Tab.

Packing : 100Tabs/BTL

Shelf life : 5 years



Composition Each tablet contains
Acetazolamide 250mg

Indications

- For the relief of edema and glaucoma by congestive heart failure, improvement of respiratory acidosis in epilepsy and emphysema of the lung

Dosage & Administration

- Relief of glaucoma:
Oral administration in divided dose for amounts of 250~1,000mg a day

Ophthalmologic Drugs

POSOD Eye Drops

Packing : 10mL

Shelf life : 2 years



Composition

Each mL contains
Potassium iodide 3mg, Sodium iodide 3mg

Indications

- For use in patient with hemorrhage of vitreous body caused by aging, myopia, hypertension, diabetes, periphlebitis and lens opacity as an early symptom of senile cataract

Dosage & Administration

- 1 drop 1~3 times a day

HANLIM HOMAPINE Eye Drops

Packing : 5mL

Shelf life : 3 years



Composition

Each mL contains
Homatropine HBr 20mg

Indications

- Mydriasis and accommodation paralysis for diagnosis or treatment

Dosage & Administration

- 1~2 drops 1 time a day

OCLE Eye Drops

Packing : 5mL

Shelf life : 2 years



Composition

Each mL contains
Tobramycin (potency) 3mg

Indications

- Blepharitis, hordeolum, dacryocystitis, conjunctivitis, keratitis, corneal ulcer

Dosage & Administration

- 1~2 drops every 4 hours
* Severe cases: 2 drops every 1 hour

OCLEX Eye Drops

Packing : 5mL

Shelf life : 2 years



Composition

Each mL contains
Tobramycin (potency) 3mg, Dexamethasone 1mg

Indications

- For steroid-responsive inflammatory ocular conditions for which a corticosteroid is indicated and where superficial bacterial ocular infection or a risk of bacterial ocular infection exists

Dosage & Administration

- Initial 24~48 hours: 1~2 drops every 2 hours
After: 1~2 drops every 4~6 hours

MOXIFORCE Eye Drops

Packing : 5mL

Shelf life : 2 years



Composition

Each mL contains
Moxifloxacin HCl 5mg

Indications

- Treatment of bacterial conjunctivitis, tarsadenitis, keratitis (including corneal ulcer), caused by susceptible strains
- Aseptic therapy before and after ophthalmic surgery

Dosage & Administration

- Bacterial conjunctivitis, tarsadenitis, keratitis (including corneal ulcer):
1 drop 3 times a day

Ophthalmologic Drugs

LEVOCLE Eye Drops

Packing : 5mL

Shelf life : 3 years



Composition	Each mL contains Levofloxacin 5mg
Indications	• Blepharitis, stye, dacryocystitis, conjunctivitis, tarsadenitis, keratitis, corneal ulcer, and aseptic therapy for ocular surgery
Dosage & Administration	• 1 drop 3 times a day

LEVOCLE Eye Drops 1.5%

Packing : 5mL

Shelf life : 3 years



Composition	Each mL contains Levofloxacin 15mg
Indications	• Blepharitis, stye, dacryocystitis, conjunctivitis, tarsadenitis, keratitis, corneal ulcer, and aseptic therapy for ocular surgery
Dosage & Administration	• 1 drop 3 times a day

QUINOVID Eye Drops

Packing : 5mL

Shelf life : 3 years



Composition	Each mL contains Ofloxacin 3mg
Indications	• Blepharitis, stye, dacryocystitis, conjunctivitis, tarsadenitis, keratitis, corneal ulcer, and aseptic therapy for ocular surgery
Dosage & Administration	• 1 drop 3 times a day

QUINOVID Eye Ointment

Packing : 5g

Shelf life : 3 years



Composition	Each gram contains Ofloxacin 3mg
Indications	• Blepharitis, stye, dacryocystitis, conjunctivitis, tarsadenitis, keratitis, corneal ulcer, and aseptic therapy for ocular surgery
Dosage & Administration	• A small amount 3 times a day

NEODEX Eye Ointment

Packing : 5g

Shelf life : 2 years




Composition	Each gram contains Dexamethasone 0.001g, Neomycin sulfate 0.0035g, Polymyxin B sulfate 6,000 IU
Indications	• Treatment of conjunctivitis, blepharitis, keratitis by bacteria sensitive to polymyxin B sulfate and neomycin sulfate
Dosage & Administration	• A small amount 3 times a day

Anti-epileptics

TOPIRON Tab. (25mg)

Packing : 30Tabs/BTL, 100Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Topiramate 25mg
	Indications	• Treatment for epilepsy 1) Monotherapy: Treatment of partial seizure with or without secondary generalized seizures 2) Additional therapy: If the administration of the primary anticonvulsants is not properly controlled; - Partial seizure with or without secondary generalized seizures - Seizures associated with Lennox-Gastaut syndrome - Primary tonic/clonic systemic seizures • Prevention for migraine
	Dosage & Administration	• Epilepsy: Initial week 1: 25mg 1 time after dinner After: Increased at an interval of 1~2 weeks by increments of 25 or 50mg/day, administered in 2 divided doses a day * Recommended dose in monotherapy: 100~200mg a day (Maximum: 500mg a day)

TOPIRON Tab. (50mg)

Packing : 30Tabs/BTL, 100Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Topiramate 50mg
	Indications	• Treatment for epilepsy 1) Monotherapy: Treatment of partial seizure with or without secondary generalized seizures 2) Additional therapy: If the administration of the primary anticonvulsants is not properly controlled; - Partial seizure with or without secondary generalized seizures - Seizures associated with Lennox-Gastaut syndrome - Primary tonic/clonic systemic seizures • Prevention for migraine
	Dosage & Administration	• Epilepsy: Initial week 1: 25mg 1 time after dinner After: Increased at an interval of 1~2 weeks by increments of 25 or 50mg/day, administered in 2 divided doses a day * Recommended dose in monotherapy: 100~200mg a day (Maximum: 500mg a day)

TOPIRON Tab. (100mg)

Packing : 30Tabs/BTL, 100Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Topiramate 100mg
	Indications	• Treatment for epilepsy 1) Monotherapy: Treatment of partial seizure with or without secondary generalized seizures 2) Additional therapy: If the administration of the primary anticonvulsants is not properly controlled; - Partial seizure with or without secondary generalized seizures - Seizures associated with Lennox-Gastaut syndrome - Primary tonic/clonic systemic seizures • Prevention for migraine
	Dosage & Administration	• Epilepsy: Initial week 1: 25mg 1 time after dinner After: Increased at an interval of 1~2 weeks by increments of 25 or 50mg/day, administered in 2 divided doses a day * Recommended dose in monotherapy: 100~200mg a day (Maximum: 500mg a day)

Psychoneural Drugs

EDPA Tab. (10mg)

Packing : 30Tabs/BTL, 100Tabs/BTL

Shelf life : 3 years



Composition Each tablet contains
Escitalopram oxalate 12.77mg (Escitalopram 10mg)

Indications

Treatment of:

- Major depressive episodes
- Panic disorder with or without agoraphobia
- Social anxiety disorder (social phobia)
- Generalized anxiety disorder
- Obsessive-compulsive disorder

Dosage & Administration

- Major depressive episodes:
10mg 1 time a day

EDPA Tab. (20mg)

Packing : 30Tabs/BTL, 100Tabs/BTL

Shelf life : 3 years



Composition Each tablet contains
Escitalopram oxalate 25.54mg (Escitalopram 20mg)

Indications

Treatment of:

- Major depressive episodes
- Panic disorder with or without agoraphobia
- Social anxiety disorder (social phobia)
- Generalized anxiety disorder
- Obsessive-compulsive disorder

Dosage & Administration

- Major depressive episodes:
10mg 1 time a day

Corticosteroids

CACEPIN Tab. (25mg)

Packing : 30Tabs/BTL, 100Tabs/BTL

Shelf life : 3 years



Composition Each tablet contains
Quetiapine fumarate 28.78mg (Quetiapine 25mg)

Indications

- Schizophrenia
- Bipolar disorder
- Acute treatment of manic episodes associated with bipolar disorder type 1
- Acute treatment of depressive episodes of bipolar disorder
- Prevention of a relapse of bipolar disorder in patients who react to mania, mixed or depressive episodes caused by administration of quetiapine

Dosage & Administration

- Schizophrenia:
2 times a day
Initial 4 days: increase to day 1 50mg, day 2 100mg, day 3 200mg, day 4 300mg
Common effective dose: 300~400mg a day
Adjust within 150~750mg a day

Corticosteroids

CACEPIN Tab. (100mg)

Packing : 30Tabs/BTL, 100Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Quetiapine fumarate 115.13mg (Quetiapine 100mg)
	Indications	• Schizophrenia • Bipolar disorder - Acute treatment of manic episodes associated with bipolar disorder type 1 - Acute treatment of depressive episodes of bipolar disorder - Prevention of a relapse of bipolar disorder in patients who react to mania, mixed or depressive episodes caused by administration of quetiapine
	Dosage & Administration	• Schizophrenia: 2 times a day Initial 4 days: increase to day 1 50mg, day 2 100mg, day 3 200mg, day 4 300mg Common effective dose: 300~400mg a day Adjust within 150~750mg a day

CACEPIN Tab. (200mg)

Packing : 30Tabs/BTL, 100Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Quetiapine fumarate 230.26mg (Quetiapine 200mg)
	Indications	• Schizophrenia • Bipolar disorder - Acute treatment of manic episodes associated with bipolar disorder type 1 - Acute treatment of depressive episodes of bipolar disorder - Prevention of a relapse of bipolar disorder in patients who react to mania, mixed or depressive episodes caused by administration of quetiapine
	Dosage & Administration	• Schizophrenia: 2 times a day Initial 4 days: increase to day 1 50mg, day 2 100mg, day 3 200mg, day 4 300mg Common effective dose: 300~400mg a day Adjust within 150~750mg a day

CACEPIN Tab. (300mg)

Packing : 30Tabs/BTL, 100Tabs/BTL

Shelf life : 3 years

	Composition	Each tablet contains Quetiapine fumarate 345.39mg (Quetiapine 300mg)
	Indications	• Schizophrenia • Bipolar disorder - Acute treatment of manic episodes associated with bipolar disorder type 1 - Acute treatment of depressive episodes of bipolar disorder - Prevention of a relapse of bipolar disorder in patients who react to mania, mixed or depressive episodes caused by administration of quetiapine
	Dosage & Administration	• Schizophrenia: Initial: 300mg 400~800mg a day

ZYAPINE Tab. (2.5mg)

Packing : 30Tabs/PTP

Shelf life : 3 years

	Composition	Each tablet contains Olanzapine 2.5mg
	Indications	• Schizophrenia • Bipolar disorder - Treatment of manic and mixed episodes associated with Bipolar I Disorder - Prevention of recurrence in patients with Bipolar Disorder who have responded to olanzapine treatment for a manic episode - Acute treatment of depressive episodes associated with Bipolar I Disorder
	Dosage & Administration	• Please refer to the insert.

Corticosteroids

ISOTAN Tab.

Packing : 60Tabs/PTP, 500Tabs/BTL

Shelf life : 3 years



Composition	Each tablet contains Tofisopam 50mg
Indications	Autonomic imbalance, head injuries, a menopausal disorder, ovarian failure
Dosage & Administration	50mg 3 times a day

Urological Agent

OBUTINE Tab.

Packing : 30Tabs/BTL, 100Tabs/BTL

Shelf life : 3 years



Composition	Each tablet contains Oxybutynin hydrochloride 5mg
Indications	Urinary frequency, urgency, and incontinence associated with unstable bladder conditions due to uninhibited neurogenic bladder (reflex neurogenic bladder).
Dosage & Administration	For adult : divided doses of 6 to 15 mg a day, in 2 to 3 doses. The total daily dose should not exceed 20 mg.

Diagnosis Agent & Diagnosis NIR Imaging Device

LUMINOMARK Inj.

Packing : 1Vial + 1A (Diluent)

Shelf life : 3 years




Composition	Each vial contains Indocyanine Green 0.01mg
Indications	Adjunctive surgical agent to label the non-palpable small area during breast lesion removal using stereotactic guidance or ultrasound.

LUMINOIMAGER-C

Size : 98 x 319 x 85mm

Weight : 860g



Composition	Camera, battery, auxiliary battery, and charger
Key Features	An NIR imaging device designed for the detection and visualization of Luminomark and ICG markers on an integrated LCD screen via near-infrared illumination.
Instructions for Use	Administration of 'Luminomark' and 'ICG' to the area where medical surgery is performed is to be luminiscent and visible to the naked eye.

For Human Health and Future
HANLIM PHARM. CO., LTD.



HEAD OFFICE

42, Seocho-daero 52-gil, Seocho-gu, Seoul, Republic of Korea

MANUFACTURING PLANT

2-27, Yeongmun-ro, Cheoin-gu, Yongin-si, Gyeonggi-do, Republic of Korea

CONTACT PERSON

Jeonghwa Jenny Yang (Ms.) | Director/Pharmacist | Global Business Dept.
Tel.: 822-3489-6214 | E-mail: yjh0717@hanlim.com