

**INFUSOL® PCM**

**Paracetamol 10mg/mL  
Intravenous Infusion  
Sterile and Non-Pyrogenic**



MAL23036018AZ

Scan to view online



- ◆ **First** locally manufactured product by using a superior aseptic blow-fill-seal (BFS) technology.
- ◆ Infusol PCM IV infusion is indicated for the short-term treatment of moderate pain, especially following surgery, and for the short-term treatment of fever, when administration by IV route is clinically justified by an urgent need to treat pain or hyperthermia and/or when other routes of administration are not possible.

# INFUSOL<sup>®</sup> PCM

## Paracetamol 10mg/mL Intravenous Infusion

Composition	Each mL solution for infusion contains 10mg Paracetamol BP. Each 100mL bottle contains 1000mg Paracetamol BP.												
Product Description	<b>INFUSOL<sup>®</sup> PCM</b> is a slightly pinkish-orange and practically free from particles.												
Pharmacodynamic	<b>INFUSOL<sup>®</sup> PCM</b> provides onset of pain relief within 5 to 10 minutes after start of administration. The peak analgesic effect is obtained in 1 hour and the duration of this effect is usually 4 to 6 hours. <b>INFUSOL<sup>®</sup> PCM</b> reduces fever within 30 minutes after the start of administration with a duration of the antipyretic effect of at least 6 hours.												
Recommended Dosage	The 100ml bottle is restricted to adults, adolescents and children weighing more than 33 kg.												
Posology	<table><thead><tr><th>Patient weight (kg)</th><th>Dose/administration</th></tr></thead><tbody><tr><td>≤10kg</td><td>7.5mg/kg</td></tr><tr><td>&gt;10kg to ≤ 33kg</td><td>15mg/kg</td></tr><tr><td>&gt;33kg to ≤ 50kg</td><td>15mg/kg</td></tr><tr><td>&gt;50kg with additional risk factors for hepatotoxicity</td><td>1g</td></tr><tr><td>&gt;50kg and no additional risk factors for hepatotoxicity</td><td>1g</td></tr></tbody></table> <p><i>* For full prescription, i.e. patient body weight, dosage and dilution information, please refer to user information.</i></p>	Patient weight (kg)	Dose/administration	≤10kg	7.5mg/kg	>10kg to ≤ 33kg	15mg/kg	>33kg to ≤ 50kg	15mg/kg	>50kg with additional risk factors for hepatotoxicity	1g	>50kg and no additional risk factors for hepatotoxicity	1g
Patient weight (kg)	Dose/administration												
≤10kg	7.5mg/kg												
>10kg to ≤ 33kg	15mg/kg												
>33kg to ≤ 50kg	15mg/kg												
>50kg with additional risk factors for hepatotoxicity	1g												
>50kg and no additional risk factors for hepatotoxicity	1g												
Method of administration	The paracetamol solution is administered as a 15-minute intravenous infusion.												
Route of administration	Intravenous (I.V)												
Storage condition before and after dilution	Do not store above 30°C.												
Dosage forms and packaging available	100ml x 20 LDPE plastic bottle per carton.												