

Glocare Labs



NUTRACEUTICALS, FOOD SUPPLEMENTS & OTC LIST

NUTRACEUTICALS & FOOD SUPPLEMENTS LIST

Sl. No.	BRANDNAME	ACTIVE INGREDIENTS	DOSAGE FORM	UNIT PACKAGING
HAIR / SKIN / NAILS				
1	Flaxseed oil Capsule	Flaxseed oil 500mg Capsule	Capsule	Blister
2	Almond oil 500mg Capsules	Almond oil 500mg Capsules	Capsule	Blister
3	Conjugated linolenic acid 500mg Capsules	Conjugated linolenic acid 500mg Capsules	Capsule	Blister
4	Vitamin E 400mg + Wheat germ oil 10mg Capsule	Vitamin E 400mg + Wheat germ oil 10mg Capsule	Capsule	Blister
5	Wheat germ oil 500mg Capsules	Wheat germ oil 500mg Capsules	Capsule	Blister
6	Evening primrose oil 500mg Capsule	Evening primrose oil 500mg Capsule	Capsule	Blister
7	Multi dosing Vitamin Capsule	Vitamin A 2500 IU + Vitamin D3 1000 IU + Vitamin E 200mg as oil Lycopene 10% 100mg as suspension + Biotin 5mg + Selenium 70 mcg + Methylcobalamin 1500mcg as pellets (Multidosing technology).	Capsule	Blister
8	Vitamin D3 Capsules	Vitamin D3 1000 IU Capsules (Stabilized in virgin coconut oil)	Capsule	Blister
9	Aloe Vera Capsule 500 mg	Aloe Vera Capsule 500 mg	Capsule	Blister
10	Olive Oil Capsule 500 mg	Olive Oil Capsule 500 mg	Capsule	Blister
11	Betavit AFR	Each Hard Gelatin Capsule contains: Betacarotene -5.6mg + Vitamin C - 100mg + Vitamin E - 29.0 mg + Vitamin B1 - 10mg + Vitamin B2 -10.0mg + Nicotinamide - 20.0mg + Vitamin B6 -24.0mg + Vitamin B12 -10.0mcg + Folic Acid -500mcg + Calcium Pantothenate -5mg + Biotin -100mcg + Iron -10.4mg + Manganese Sulphate -4.0mg + Potassium -5.0mg + Copper - 0.5mg + Zinc -15.0mg + Calcium -75mg + Magnesium - 22.0mg + Selenium -30.0mcg + Chromium -27.0mcg	Capsule	Blister
12	RICHVIT HAIR TABLETS	Each Film coated tablet contains: Vitamin A 800 mcg + Vitamin B1 (Thiamine) 5mg + Vitamin B2 (Riboflavin) 5mg + Vitamin B3 (Niacin) 5mg + Vitamin B5 (Pantothenic Acid) 100mg + Vitamin B6 (Pyridoxine HCl) 5mg + Folic Acid 200mcg + Biotin 250 mcg + Vitamin B12 (Cyanocobalamin) 0.6mg + Vitamin C (Ascorbic Acid) 100mg + Vitamin D3 (Cholecalciferol) 400IU + Calcium 200mg + Copper 1mg + Selenium 55mcg + Zinc 10mg + Methylsulfonylmethane (MSM) 250mg + Marine Collagen 200mg + Silica 20mg + Hyaluronic Acid 50mg + L-Cysteine 15mg + L-Methionine 15mg + L-Lysine 15mg + BCAA Providing 40mg (Branched chain Amino acid (L-Leucine, L-Isoleucine and L-Valine) L-Leucine 20mg + L-Isoleucine 10mg + L-Valine 10mg)	Tablet	Blister
13	RICHVIT RADIANCE CAPS	Each hard gelatin Capsule contains: Vitamin C 40mg + Zinc 5mg + Vitamin E 12mg + Methyl sulfonyl methane 10mg + Niacinamide 8mg + Vitamin A 400mcg + Pantothenic Acid 3mg + Copper 150mcg + Vitamin D3 5mcg + Vitamin B6 1.4mg + AloeVera 200mg + Iodine 150mcg + Folic Acid 200mcg + Biotin 50mcg + Vitamin B12 2.5mcg + Vitamin C 40mg + Zinc 5mg + Vitamin E 12mg + Methyl sulfonyl methane 10mg + Niacinamide 8mg + Vitamin A 400mcg + Pantothenic Acid 3mg + Copper 150mcg + Vitamin D3 5mcg + Vitamin B6 1.4mg + AloeVera 200mg + Iodine 150mcg + Folic Acid 200mcg + Biotin 50mcg + Vitamin B12 2.5mcg	Capsule	Blister
EYE				
14	Lutein and Zeaxanthin Capsule	Lutein and Zeaxanthin Capsule	Capsule	Blister
HEART				
15	Co-Enzyme Q10 -100mg + Piperine 5mg Capsule	Co-Enzyme Q10 -100mg + Piperine 5mg Capsule	Capsule	Blister
16	Garlic oil Capsule	Garlic oil 0.25% 500mg Capsule	Capsule	Blister
17	Fish oil Capsule	Fish oil 500mg	Capsule	Blister

18	Grape Seed Extract Capsule 100 mg	Grape Seed Extract Capsule 100 mg	Capsule	Blister
19	Ginkgo Biloba Capsules 500mg	Ginkgo Biloba Capsules 500mg	Capsule	Blister
20	Vitamin E & EPA Capsules(40mg+100mg)	Vitamin E & EPA Capsules (40mg+100mg)	Capsule	Blister
21	Coconut Oil Capsule	Coconut Oil Capsule	Capsule	Blister
22	Nigella oil	Black Seed Oil 500 mg Capsules	Capsule	Blister
MULTIVITAMIN + AMINOACID				
23	Miltiz Tablet / Syrup,	Each 5ml contains: Thiamine hydrochloride BP 2mg + Riboflavin sodium phosphate BP 2.5mg + Pyridoxine hydrochloride BP 0.75mg + Nicotinamide BP 15mg + D-Panthenol USP 3mg + Cyanocobalamin BP 5mcg Colour: Approved colour used	Capsule	Blister / Bottle
24	Provita + Capsules	L-Threonin E -2.10mg + L-Isoleucine -2.95mg + L-Leucine -9.15mg + L-Lysine Hydrochloride -12.5mg + L-Methionine 9.20mg + L-Phenylalanine- 2.50mg + L- Arginine Hcl -6.60mg + L-Histidine Hcl 1.85mg + L-Tryptophan -1mg + L-Valine 3.40mg + Vitamin B1 Mononitrate -1.5mg + Vitamin B2 - 1.7mg + Vitamin B6 -1.5mg +Nicotinamide -20mg + Calcium Pantothenate -5mg + Folic Acid - 0.75mg + Cyancobalamin - 2.5mcg + Ascorbic Acid - 40mg + Vitamin E -7.50 IU + Potassium Iodide -1.8mg + Ferrous Sulphate -20.5mg + Copper Sulphate 2.00mg + Magnesium Sulphate 35.5mg + Manganese Sulphate 1.40mg + Zinc Sulphate - 3.30mg + Sodium Selenite -10mcg	Capsules	Blister
25	Provita + liquid	L-Isoleucine -5.90mg + L-Leucine -18.3mg + L-Lysine Hydrochloride -25mg + L-Methionine 9.2mg + L-Phenylalanine- 5mg + L-Threonine -4.2mg + L-Tryptophan -5mg + L-Valine 6.7mg + Cholecalciferol -200IU + Vitamin B1 Hcl -1.5mg + Riboflavin Sodium Phosphate -1.5mg + Vitamin B6 - 1.5mg + Nicotinamide -20mg + Calcium Pantothenate -5mg + Folic Acid - 0.75mg Cyanocobalamin - 2.5mcg + Ascorbic Acid -40mg + Vitamin A Palmitate -500IU + Vitamin E -7.50IU	Syrup	Bottle 200ml
26	Richvit Kid Tablets	Vitamin B1 Mononitrate -1mg + Vitamin B2 - 1.2mg + Vitamin B6 - 1.5mg + Vitamin B12- 3mcg + Vitamin D3 -200IU + Biotin 0.25mg + Nicotinamide - 15mg + Pantothenic Acid -4mg + Ascorbic Acid -75mg + Vitamin E - 5mg + Vitamin A(As Acetate) 2500 IU + Magnesium Oxide 25mg + Zinc Sulphate Monohydrate 2mg + Ferrous Sulphate -1.5mg	Tablets	Blister
27	Richvit Syrup	Cholecalciferol -400IU + Vitamin B1 Hcl -1mg + Riboflavin Sodium Phosphate -1.2mg + Vitamin B6 -1mg Nicotinamide -10mg + D-Panthenol -2.5mg + L-Lysine Hcl - 8mg + Vitamin A Palmitate 3000 IU + Ascorbic Acid -40mg + Cyanocobalamin -3mcg	Syrup	Bottle
28	G-Vit Syrup	Vitamin A As Palmitate - 2500 IU + Cholecalciferol – 250 IU + Cyanocobalamine - 10mcg + Riboflavin (As Riboflavin Sodium Phosphate) - 0.75mg + Thiamine Hydrochloride - 0.75mg + Nicotinamide - 7.5mg + D-Panthenol -1.25mg + Vitamin E – 2.5IU + L- Lysine Hydrochloride - 5mg + Zinc Sulphate Heptahydrate Eq. To Elemental Zinc - 22.5mg + Sodium Selenite Pentahydrate Eq. To Selenium -50mcg	Syrup	Bottle
29	G-Vit Drops	Vitamin A As Palmitate - 1000 IU + Cholecalciferol - 100 IU + Cyanocobalamin - 1.5mcg + Riboflavin (As Riboflavin Sodium Phosphate) - 0.6mg + Thiamine Hydrochloride -0.8mg + Pyridoxine Hydrochloride - 0.5mg + Nicotinamide - 8mg + Ascorbic Acid -40mg D - Panthenol - 1mg + Vitamin E-0.5 IU + L - Lysine Hydrochloride - 10mg + Zinc Sulphate Heptahydrate + Eq. To Elemental Zinc – 10mg Colour : Caramel	Drops	Bottle
30	Richvit Drops	Cholecalciferol -100 IU + Vitamin B1 Hcl -1mg + Riboflavin Sodium Phosphate -1.2mg + Vitamin B6 -1mg + Nicotinamide -10mg + D-Panthenol -2.5mg + L-Lysine Hcl - 8mg + Vitamin A Palmitate 1000 IU + Cyanocobalamin -3mcg + Ascorbic Acid- 40mg	Drops	Bottle
DM				
31	Fenugreek oil 500mg Capsule	Fenugreek oil 500mg Capsule	Capsule	Blister

BONE & JOINT

32	Arthoflex Tablets	Glucosamine Sulphate Potassium 750mg + Methyl Sulphonyl Methane -250mg + Copper Sulphate 2mg + Magnesium Sulphate 2mg + Vitamin D3 -125 IU	Tablets	Blister
----	--------------------------	--	---------	---------

MUSCLE DEVELOPMENT

33	Promass Capsules	L-Leucine 200mg + L-Isoleucine 100mg + L-Valine 100mg + L-Glutamine -100mg	Capsules	Blister
----	-------------------------	--	----------	---------

B COMPLEX

34	Becoz Syrup	Vitamin B12 -0.4mcg + L-Lysine Hcl - 50mg + Zinc Sulphate 15mg	Syrup	Bottle
----	--------------------	--	-------	--------

IRON

35	Ferrol Capsules	Sodium Feredate Eq. To Iron - 15mg + Folic Acid -5mg + Cyancobalamin-1.5mcg	Capsules	Blister
----	------------------------	---	----------	---------

36	Feramin Syrup	Ferric Ammonium Citrate -50mg + Vitamin B1 Hcl -5mg + Vitamin B2 Sodium Phosphate -5mg + Vitamin B6 -5mg + D Panthenol -20mg + Nicotinamide - 50mg + Vitamin B 12 -7.5mcg + Vitamin C- 7.5mg + Folic Acid -1mg + Vitamin E - 15 IU + Zinc Sulphatemonohydrate Eq. To Elemental Zinc 11mg + Selenium Dioxide Eq. to Selenium - 60mcg	Syrup	Bottle
----	----------------------	---	-------	--------

37	Feramin Tablets	Methycobalamine -500mcg + Folic Acid -0.9mg + Carbonyl Iron -15mg + Zinc Sulphate monohydrate Eq. to zinc -15mg	Tablets	Blister
----	------------------------	---	---------	---------

APPETITE STIMULANT

38	Cypro Vit Syrup	Cyproheptadine Hcl - 2mg + L-Lysine Hcl -150mg + Flavoured Syrup Base	Syrup	Bottle
----	------------------------	---	-------	--------

ANTIOXIDANTS / MULTIVITAMINS

39	Nataplex P	Each Hard Gelatin Capsule Contains: Vitamin A (As Acetate) 6000 IU + Vitamin D 400IU + Thiamine Mononitrate 3mg + Riboflavin 3mg + Pyridoxine Hydrochloride 3mg + Nicotinamide 20mg + Calcium Pantothenate 2mg + Vitamin C -100mg + Folic Acid 0.9mg + Vitamin B12 - 2mcg + Iron As Ferrous Fumarate 65mg + Calcium As Calcium Carbonate 80mg + Iodine As Potassium Iodide 75mcg + Manganese Sulphate 0.5mg + Potassium Chloride 3.8mg + Molybdenum As Sodium Molybdate Dihydrate 0.1mg + Zinc Sulphate Monohydrate 22.5mg + Magnesium Chloride Eq. to Magnesium -25mg + Copper As Copper Gluconate 0.5mg + Biotin USP 30mcg	Capsules	Blister
----	-------------------	---	----------	---------

40	SUPERVIT-G CAPSULES	Each Hard Gelatin capsule contains: Ascorbic acid 15mg + Cholecalciferol 300 IU + Nicotinamide 7.5mg + Riboflavin 1mg + Thiamine mono nitrate 1mg + Vitamin A (as acetate) 2500 IU	Capsule	Blister
----	----------------------------	---	---------	---------

41	SUPERVIT-G SYRUP	Each 10ml contains: VITAMIN A (As Palmitate) 5000IU + CHOLECALCIFEROL 400IU + THIAMINE HYDROCHLORIDE 1.0mg + RIBOFLAVIN (As Sodium Phosphate) 1.2mg + PYRIDOXINE HYDROCHLORIDE 0.5mg + NICOTINAMIDE 10.0mg.	Syrup	Bottle
----	-------------------------	--	-------	--------

42	ZINCBIOVIT CAPSULES / TABLETS / SYRUP	Each hard gelatin capsule contains: Vitamin A (as acetate) 5000 IU + Cholecalciferol (Vitamin D3) 200 IU + Vitamin E (as dry powder) 25 IU + Thiamine mononitrate 10mg + Riboflavin 10mg + Pyridoxine Hydrochloride 3mg + Cyanocobalamin 5mcg + Ascorbic acid 100mg + Biotin 30mcg + Niacinamide 50mg + Folic acid 0.9mg + Calcium pantothenate 12.5mg + Manganese chloride (Eq.to Manganese) 1.4mg + Copper Sulphate pentahydrate 2.5mg + Calcium Di basic phosphate 162mg + Chromium chloride (Eq.to Elemental Chromium) 65mcg + Zinc sulfate monohydrate (Eq.to.Elemental zinc) 15mg + Dried ferrous sulphate 33.3mg + Sodium selenite (Eq. to elemental selenium) 60mcg.	Capsule	Blister / Bottle
----	--	---	---------	------------------

43	AMINOVIT CAPS	Each hard gelatin Capsule contains L-Threonine 2.10mg + L-Valine 3.40mg + L-Methionine 9.20mg + L-Isoleucine 2.95mg + L-Leucine 9.15 mg + L-Phenylalanine 2.50mg + L-Tryptophan 1.00 mg + L-Lysine hydrochloride 12.50mg + L-Arginine hydrochloride 6.60mg + L-Histidine hydrochloride 1.85mg + Thiamine mononitrate 1.50mg + Riboflavin 1.70mg + Nicotinamide 20.00mg + Pyridoxine hydrochloride 1.50mg + Folic acid 0.75 mg + Calcium pantothenate 5.00 mg + Cyanocobalamin 2.50 mcg + Ascorbic acid 40 mg +	Capsule	Blister
----	----------------------	---	---------	---------

		Alpha tocopheryl acetate 7.5IU + Potassium Iodide 0.10mg + Ferrous sulphate 20.50 mg + Copper sulphate 2.00mg + Magnesium sulphate 7.42mg + Manganese sulphate 1.40mg + Zinc sulphate 3.30 mg + Sodium selenite 10.00mcg		
EYE				
44	Ocugard	Each Film Coated Tablet Contains: Vitamin C 250mg + Ginkgo Biloba Extract 60mg + Alpha Lipoic Acid 37.5mg + Rutin 25mg + Zinc (As Zinc Oxide) 12.5mg + Bilberry Extract 5mg + Lutein 3mg + Beta Carotene -2.5mg + Zeaxanthin -0.5mg + Selenium 13mcg + Vitamin E (As Alpha Tocopheryl Acetate Concentrate) (Powder Form) - 100IU	Tablet	Blister
PROBIOTIC				
45	GASTRO LB PLUS CAPSULES	Each Hard Gelatin capsule contains: Lactobacillus Sporogenes 10 billion + Simethicone 10mg + Vitamin B 12 1 mcg + L-Lysine mono HCL 5mg + Niacinamide 15mg	Capsule	Blister
DIABETES				
46	RICH DIABETES PLUS	Each filmcoated tablets contains: Niacin (as Nicotinamide) 20mg + Zinc Sulphate 3.3mg + Ferrous Fumarate 15mg + Pantothenic Acid 4 mg + Vitamin B12 (Cyanocobalamin) 0.6mg + Vitamin A 2500IU + Riboflavin 10mg + Manganese Sulphate 2mg + Copper Sulphate 2 mg + Vitamin D3 400IU + Chromium chloride 65 mcg + Folic Acid (as Pteroylmonoglutamic Acid) 0.9mg + Biotin 30 mcg + Sodium Selenate 60mcg + Potassium Iodide 150mcg	Tablet	Blister
MEN SPECIAL				
47	RICHVIT ALPHA MAN	Each film coated contains Vitamin D 0.01mg + Potassium 50mg + Vitamin E 20mg + Vitamin C 100mg + Thiamin (Vitamin B1) 5mg + Riboflavin (Vitamin B2) 5mg + Niacin 20mg + Pantothenic Acid 5mg + Vitamin B6 5mg + Vitamin B12 20mcg + Folic Acid 400mcg + Biotin 100 mcg + Vitamin K 30 mcg + Calcium 210mg + Chromium 200mcg + Copper 2mg + Iodine 150mcg + Magnesium 125mg + Manganese 2mg + Molybdenum 75mcg + Selenium 200mcg + Zinc 15mg + Beta Carotene 2.1mg + Coenzyme Q-10 4.9mcg	Tablets	Blister
GROWTH & DEVELOPMENT				
48	RICHVIT GROW TABLETS	Each film coated tablet contains: Vitamin C (as Ascorbic Acid) 350 mg + Vitamin D (as Cholecalciferol D-3)400 IU + Vitamin E (as Natural d-alpha Tocopheryl Succinate) 30 IU + Thiamin (Vitamin B-1) (as Thiamin Mononitrate) 4.5mg + Riboflavin (Vitamin B-2) 5.1 mg + Niacin (as Niacinamide) 20 mg + Vitamin B-6 (as Pyridoxine Hydrochloride) 6 mg + Folic Acid 400 mcg + Vitamin B-12 (as Cyanocobalamin) 18 mcg + Biotin 45 mcg + Pantothenic Acid (as Calcium d-Pantothenate)10mg + Calcium (as Calcium Carbonate, Dicalcium Phosphate and Calcium Citrate)100 mg + Iron (as Carbonyl Iron)15 mg + Phosphorus (as Dicalcium Phosphate) 20 mg + Iodine (as Potassium Iodide)150 mcg + Magnesium (as Magnesium Oxide) 50 mg + Zinc (as Zinc Oxide)15 mg + Selenium (as Selenium Yeast) 5mcg + Copper (as Cupric Sulfate) 2 mg + Manganese (as Manganese Sulphate) 2 mg + Chromium (as GTF Chromium Yeast)20 mcg + Molybdenum (as Molybdenum Yeast) 20 mcg + Potassium (as Potassium Chloride) 25 mg + Carrot Powder 3 mg + Choline (as Choline Bitartrate) 50 mcg + Inositol 50mcg	Tablet	Blister
49	RICHVIT TEEN TABLETS	Each Film coated tablet contains: Vitamin C (as Ascorbic Acid) 350 mg + Vitamin D (as Cholecalciferol D-3) 400 IU + Vitamin E (as Natural d-alpha Tocopheryl Succinate) 30 IU + Thiamin (Vitamin B-1) (as Thiamin Mononitrate) 4.5mg + Riboflavin (Vitamin B-2) 5.1 mg + Niacin (as Niacinamide) 20 mg + Vitamin B-6 (as Pyridoxine Hydrochloride) 6 mg + Folic Acid 400 mcg + Vitamin B-12 (as Cyanocobalamin) 18 mcg + Biotin 45 mcg + Pantothenic Acid (as Calcium d-Pantothenate)10mg + Calcium (as Calcium Carbonate, Dicalcium Phosphate and Calcium Citrate)100 mg + Iron (as Carbonyl Iron)15 mg + Phosphorus (as Dicalcium Phosphate) 20 mg + Iodine (as Potassium Iodide)150 mcg + Magnesium (as Magnesium Oxide) 50 mg + Zinc (as Zinc Oxide)15 mg + Selenium (as Selenium Yeast) 5mcg +	Tablet	Blister

		Copper (as Cupric Sulfate) 2 mg+ Manganese (as Manganese Sulphate) 2 mg + Chromium (as GTF Chromium Yeast)20 mcg + Molybdenum (as Molybdenum Yeast) 20 mcg + Potassium (as Potassium Chloride) 25 mg + Carrot Powder 3 mg + Choline (as Choline Bitartrate) 50 mcg + Inositol 50mcg		
WOMENS				
50	RICHVIT PREGNANCY PLUS	Each hard gelatin Capsule contains: Iron 15mg + Folic acid 0.9mg+ Vitamin B1 2.2mg Vitamin B2 2.8mg+ Vitamin B3 8mg + Vitamin B5 12mg Vitamin B6 2.8mg+ Vitamin B12 (Cyanocobalamin) 25mcg Vitamin C 80mg+ Vitamin D 0.01mg+ Vitamin E 12mg Vitamin K 84mcg+ Calcium 120mg+ Magnesium 105mg	Capsule	Blister
51	RICHVIT WOMEN	Each hard gelatin Capsule contains: L-Carnitine (Also includes Co-Q10, Guarana and Green Tea extracts) 30mg + Alpha Lipoic Acid 20mg + L-Methionine 40mg + N-Acetyl Cysteine 30mg + Green tea extracts 40mg + Guarana extract 100mg + Co-enzyme Q10 2mg + Citrus Bioflavonoids 10mg + Natural Mixed Carotenoids 2mg + Vitamin D 0.01mg + Vitamin E 20 mg + Vitamin K 90 mcg + Vitamin C 80 mg + Thiamin (Vitamin B1) 10mg + Riboflavin (Vitamin B2) 4 mg + Niacin (Vitamin B3) 36mg + Vitamin B6 8mg + Folic acid 400 mcg + Vitamin B12 0.6mg Biotin 50 mcg + Pantothenic acid 6mg + Magnesium 100mg + Iron 14mg + Zinc 15mg + Copper 1.0mg + Manganese 2mg + Selenium 100 mcg + Chromium 50 mcg + Iodine 150 mcg + Vitamin E 6.6 mg	Capsule	Blister
52	Sea buckthorn seed oil	Sea buckthorn seed oil 500mg (HPMC Capsule)	Capsule	Blister
53	Vitamin A & D Capsules (250IV&1000IV)	Vitamin A & D Capsules (250IV & 1000IV)	Capsule	Blister
54	Natural Vit E Capsule 4 mg	Natural Vit E Capsule 4 mg	Capsule	Blister
55	Vitamin B Complex Capsules	Vitamin B Complex Capsules	Capsule	Blister
56	Ener - G Capsule	Each hard gelatin capsule contains: Ginseng Extract 50.0mg + Natural Betacarotene 6.0mg + Vitamin D3 200.0 IU + Vitamin B1 1.5mg + Vitamin B2 1.5mg + Vitamin B6 2.0mg + Vitamin B12 2.0mcg + Vitamin C 60.0mg + Natural Vitamin E 15.0 IU + Nicotinamide 20.0mg + Calcium Pantothenate 6.0mg + Dibasic Calcium Phosphate 255.0mg + Folic Acid 100.0mcg + Iron 12.0mg + Copper 1.0mg + Zinc 5.0mg + Magnesium 2.0mg + Manganese 1.0mg + Potassium 3.5 mg + Selenium 15.0mcg	Capsule	Blister
57	Cypro Vit Tablet	Cyproheptadine Hcl - 2mg	Tablet	Blister
58	RICHVIT SENIOR	Each film coated contains Siberian Ginseng Extract 20mg + Co-Q10 10mg + Pumpkin Seed Extract 100mg + Phosphatidylcholine 7mg + L-Glutathione 10mg + Lutein Esters 2mg + Alpha Lipoic Acid (ALA) 20mg + L-Carnitine 20mg + L-Carnitine 20mg + Citrus Bioflavonoids 10mg + Natural Mixed Carotenoids 2mg + Vitamin A (2666 IU) 800mcg + Vitamin D 0.01mg Vitamin E 12mg + Vitamin C 80mg + Thiamin (Vitamin B1) 14mg + Riboflavin (Vitamin B2) 4mg + Niacin (Vitamin B3) 36mg + Vitamin B6 10mg + Biotin 50mcg + Panthothenic Acid 10mg + Iron 6mg + Zinc 15mg + Copper 1000mcg + Manganese 3.5mg + Selenium 150mcg + Chromium 75mcg + Iodine 150mcg + Silcon 10mg	Tablet	Blister
59	VIT C -500	Each uncoated chewable tablet contains: Ascorbic acid BP 500 mg	Tablet	Strip

CREAM 15 gms



GLORIDERM

Composition:

Beclomethasone Dipropionate USP 0.025% w/w
 Clotrimazole BP 1% w/w
 Neomycin sulphate BP 0.5% w/w
 Preservative : Chlorocresol BP 0.1% w/w

Store below 30°C, Protect from light.

DO NOT FREEZE

To be used only under Medical Supervision.

FOR EXTERNAL USE ONLY

Puncture tube nozzle seal with piercing point of cap

OINTMENT 5 gms



GLOPIROCIN

Composition:

Mupirocin USP 2.0% w/w
 Ointment base q.s.

FOR EXTERNAL USE ONLY

Store below 30°C,
 Protect from light.

DO NOT FREEZE

WARNING: To be sold by retail on the prescription of a Registered Medical Practitioner only.

Puncture nozzle seal with piercing point of cap.

GEL 30 gms



DOLMAX GEL

Composition :

Diclofenac Diethylamine BP
 Equivalent to Diclofenac Sodium 1% w/w in a gel base.

Store below 30°C, Protected from heat & light.

Do not allow to freeze

Replace cap after use

Puncture nozzle seal with spike on cap

FOR EXTERNAL USE ONLY

GEL 30 gms



GLOPHOB GEL

Composition :

Heparin Sodium BP 200IU
 Preservative : Benzyl alcohol BP 0.1% w/w
 Aqueous Base.

Use as directed by the physician.

Store in a cool place, protect from light.

To be used only under medical supervision.

DO NOT FREEZE

FOR EXTERNAL USE ONLY

Puncture nozzle seal with piercing point of the cap

CREAM 15 gms



GLOBENZ

Composition :
Benzocaine BP 20% w/w
Cream base q.s.

Directions: Use as and when required.
Do not use in infants.

NOT TO BE SWALLOWED.
KEEP TUBE CAPPED WHEN NOT IN USE.
Store at a temperature not exceeding 25°C.

CREAM 10 gms



FUNGITER

Composition:
Terbinafine Hydrochloride BP 1%W/W

Keep all medicines out of reach of children.

Store at a temperature not exceeding 25°C.
Do not freeze.

Use as directed by the physician.

For External use only

Puncture nozzle seal with piercing point of cap

CREAM 15 gms



KEZOLE CREAM

Composition :
Each gram contains :
Ketoconazole USP 20 mg
in a cream base.

Direction for use: See package insert.
Replace the cap tightly after use.

Store at a temperature not exceeding 25°C Do not freeze.

FOR EXTERNAL USE ONLY

Puncture nozzle seal with piercing point of cap

CREAM 15 gms



CLONATE CREAM

Composition :
Clobetasol propionate BP 0.05% w/w
Cream base.

FOR EXTERNAL USE ONLY

Puncture nozzle seal with piercing point of cap

Store below 30°C, Protect from light.
Do not freeze.

OINTMENT 15 gms



POLYBAC-N

Composition :

Each gram contains :

Polymyxin B Sulfate USP 5000 Units
Bacitracin Zinc USP 400 Units
Neomycin Sulfate BP 3400 Units

Store below 30°C. Do not freeze
Keep this medicine out of reach of children.
Not for use in eyes

Use as directed by the Physician.
Do not use if allergy to any of its ingredients.

FOR EXTERNAL USE ONLY

OINTMENT 15 gms



GLOSIL-C

Composition :

Silver Sulphadiazine USP 1% w/w
Chlorhexidine Gluconate BP 0.2% w/w
Ointment Base q.s.

Indications : Cuts, Wounds, Ulcers, Burns and bacterial infections of the skin

FOR EXTERNAL USE ONLY

Puncture nozzle seal with piercing point of cap

Store at a temperature not exceeding 25°C
Do not freeze.

CREAM 15 gms



GLOFLAM

Composition:

Hydrocortisone BP 1% w/w
Cream Base Q.S.

Dosage & Directions for use:

Should be applied to the affected area two to four times daily depending on the severity of the condition.

FOR EXTERNAL USE ONLY

Store below 30°C,
Protect from light.

DO NOT FREEZE

OINTMENT 10 gms



GLOBETA-M

Composition:

Povidone Iodine USP 5% w/w (0.5% w/w available Iodine)
Metronidazole BP 1% w/w
Ointment Base. Q.S.

Use as directed by the Physician.

Puncture nozzle seal with piercing point of the cap.

FOR EXTERNAL USE ONLY

Store below 30°C,
Protected from light.
DO NOT FREEZE



CREAM

30 gms



TOLNAFTATE

s-Cream

Drug Facts

ACTIVE INGREDIENT

Tolnaftate 1%

PURPOSE

Antifungal

INACTIVE INGREDIENTS

Disteardimonium hectorite, Fragrance, Isobutane, Isopropyl myristate, SD alcohol 40-B, Sodium bicarbonate.

USES

■ Cures most athlete's foot (tinea pedis) and ringworm (tinea corporis) ■ If used daily, can prevent recurrence of athlete's foot ■ Relieves symptoms of athlete's foot, including itching, burning and cracking.

WARNINGS

For external use only

Flammable:

Do not use while smoking or near heat or flame. Do not puncture or incinerate. Contents under pressure. Do not store at temperature above 120°F

When using this product

- Do not get into eyes or mouth, if products get into eyes, rinse eyes thoroughly with water
- Use only as directed
- Intentional misuse by deliberately concentrating and inhaling contents can be harmful or fatal.

Stop use and ask a doctor if

- Irritation occurs
- No improvement within 4 weeks

Keep out of the reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts
(continued)

Do not use on children under 2 years of age unless directed by a doctor.

DIRECTIONS

Wash affected area and dry thoroughly

Spray a thin layer over affected area twice daily (morning and night)

Supervise children in the use of this product

For athlete's foot: pay special attention to spaces between the toes, wear well-fitting, ventilated shoes and change shoes and socks at least once daily

Use daily for 4 weeks; if conditions persist, consult a doctor.

Apply once or twice daily (morning and/or night).

OTHER INFORMATION

Store between 20° and 30°C (68°F and 86°).

CREAM

30 gms



MICONAZOLE NITRATE

s-Cream

Drug Facts

ACTIVE INGREDIENT

Miconazole Nitrate USP 2%

PURPOSE

Antifungal

INACTIVE INGREDIENTS

Cetostearyl alcohol, Cetomacrogol 1000, Paraffin wax, Micro crystalline wax, Light liquid paraffin, Polyethylene Glycol, Chlorocresol and purified water

USES

■ Effective in the treatment of most athlete's foot (tinea pedis), jock itch (tinea cruris), and ringworm (tinea corporis) ■ For effective relief of itching, scaling, cracking, burning and discomfort that can accompany these conditions.

WARNINGS

For external use only

Do not use on children under 2 years of age unless directed by a doctor.

When using this product avoid contact with eyes.

Stop use and ask a doctor if

- Irritation occurs
- Condition persists
- There is no improvement of athlete's foot or ringworm within 4 weeks or jock itch within 2 weeks.

Keep out of the reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

DIRECTIONS

- Clean the affected area and dry thoroughly
- Apply a thin layer of the product over affected area twice daily (morning and night) or as directed by a doctor
- Supervise children in the use of this product

Drug Facts
(continued)

■ For athlete's foot: pay special attention to spaces between the toes; wear well-fitting, ventilated shoes, and change shoes and socks at least once daily ■ For athlete's foot and ringworm: use daily for 4 weeks ■ For jock itch: use daily for 2 weeks ■ Not effective on the scalp or nails

OTHER INFORMATION

■ Do not use if seal is damaged or is not visible. ■ To open, unscrew cap, pull tab to remove foil seal

■ store at temperature not exceeding 30°C (86°F). Do not freeze. ■ see carton or tube crimp for lot number and expiration date

CREAM

30 gms



ZINC OXIDE

s-Cream

Drug Facts

ACTIVE INGREDIENT

Zinc Oxide 13%

PURPOSE

Skin Protectant

INACTIVE INGREDIENTS

Water, Mineral oil, Petrolatum, Beeswax, Dimethicone, Sorbitan sesquiolate, Microcrystalline wax, PEG-30 Dipolyhydroxystearate, Aloe barbadensis leaf extract, Glycerin, Tropolone, Tocopheryl acetate, 1,2-Hexanediol, Caprylyl glycol, Magnesium sulfate, Potassium hydroxide, Phenoxethanol

USES

- Helps treat and prevent diaper rash
- Protects chafed skin due to diaper rash and helps seal out moisture
- Insect bites
- Burns minor abrasions
- Skin irritation

WARNINGS

For external use only

When using this product

Do not get into eyes

Stop use and ask a doctor if

- Condition worsens
- Symptoms last more than 7 days or clear up and occur again within a few days

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

DIRECTIONS

- Change wet and soiled diapers promptly
- Cleanse the diaper area
- Allow to dry

Drug Facts
(continued)

■ Apply cream liberally as often as necessary, with each diaper change, especially at bedtime or anytime when exposure to wet diapers may be prolonged

OTHER INFORMATION

store at 20° to 25°C (68°F to 77°F)

Protect from freezing

May stain clothing



CREAM 30 gms



CLOTRIMAZOLE s-Cream

Drug Facts

ACTIVE INGREDIENT

Clotrimazole 1%

PURPOSE

Antifungal

INACTIVE INGREDIENTS

Benzyl alcohol (1%), Cetostearyl alcohol, Cetyl esters wax, 2-Octyldodecanol, Polysorbate 60, Purified water, Sorbitan monostearate

USES

Athlete's foot, jock itch and Ringworm, Relieves itching, burning, cracking, scaling and discomfort which accompany these conditions

WARNINGS

For external use only

Do not use on children under 2 years of age unless directed by a doctor.

When using this product avoid contact with eyes.

Stop use and ask a doctor if

Irritation occurs; There is no improvement within 4 weeks (for athlete's foot or ringworm) or 2 weeks (for jock itch).

Keep out of the reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

DIRECTIONS

- Wash affected area and dry thoroughly
- Apply a thin layer over affected area twice daily (morning and night)
- Supervise children in the use of this product

Drug Facts (continued)

- For athlete's foot, pay special attention to spaces between the toes, wear well-fitting, ventilated shoes and change socks at least once daily
- For athlete's foot and ringworm, use daily for 4 weeks; for jock itch, use daily for 2 weeks
- If condition persists longer, seek medical advice
- This product is not effective on the scalp or nails

OTHER INFORMATION

To open: unscrew cap, use the pointed end of cap to puncture seal. Store between 20° - 25°C (68° - 77°F) See carton or tube crimp for lot number and expiration date.

OINTMENT 30 gms



TRIPLE ANTIBIOTIC s-Oint

Drug Facts

ACTIVE INGREDIENT

Bacitracin Zinc 400 units
Neomycin Sulfate 3.5mg
Polymyxin B Sulfate 5,000 units

PURPOSE

First aid
Antibiotic

USES

First aid to help prevent infection in minor:

- Cuts ■ Scrapes ■ Burns

WARNINGS

For external use only

Do not use if you are allergic to any of the ingredients

Ask a doctor before use if you have

- Serious burns ■ Deep or puncture wounds ■ Animal bites

When using this product

- Do not use in the eyes
- Do not apply over large areas of the body

Stop use and ask a doctor if

- You need to use more than 1 week ■ Condition persists or gets worse ■ Symptoms persist for more than 1 week or clear up and occur again within a few days, rash or other allergic reaction develops

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

DIRECTIONS

adults and children 2 years of age or older:

clean the affected area

apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily

Drug Facts (continued)

may be covered with a sterile bandage children under 2 years of age: ask a doctor

OTHER INFORMATION

Store at 59° to 77°F

INACTIVE

INGREDIENTS

Cetyl alcohol, Glycerin, Light mineral oil, Myristoyl / palmitoyl oxostearamide / arachamide MEA, Steareth-w, Squalane, Stearic acid, Tocopheryl acetate, White petrolatum

OINTMENT 30 gms



BACITRACIN s-Oint

Drug Facts

ACTIVE INGREDIENT

Bacitracin 500 units

PURPOSE

First aid antibiotic

USES

Helps prevent infection in minor cuts, scrapes and burns

WARNINGS

For external use only

Do not use

- if you are allergic to any of the ingredients ■ in the eyes ■ over large areas of the body ■ longer than 1 week unless directed by a doctor.

Stop use and ask a doctor if

- The condition persists or gets worse ■ A rash or other allergic reaction develops

Keep out of the reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

DIRECTIONS

- Clean the affected area
- Apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- May be covered with a sterile bandage

Drug Facts (continued)

OTHER INFORMATION

Store at room temperature

INACTIVE INGREDIENT

Light mineral oil, White petrolatum



OINTMENT 30 gms



HYDROCORTISONE s-Oint

Drug Facts

ACTIVE INGREDIENT
Hydrocortisone USP 1%
PURPOSE
Steroid Cream

USES

■ For relief of itching associated with minor skin irritations and rashes due to: ■ Eczema ■ Insect bites ■ Soaps and detergents ■ Cosmetics ■ Jewelry ■ Seborrheic dermatitis ■ Psoriasis ■ Poison ivy, oak or sumac ■ For external genital, feminine and anal itching ■ Other uses of this product should be only under the advice and supervision of a doctor

WARNINGS

For external use only

Do not use ■ In children under 2 years of age ■ If you have a vaginal discharge ■ For the treatment of diaper rash

Ask a doctor before use if you have ■ External genital or feminine itching ■ External anal itching ■ Bleeding

When using this product ■ Avoid contact with eyes ■ Do not exceed the recommended daily dosage unless directed by a doctor ■ Do not put this product into the rectum by using fingers or any mechanical device or applicator

Stop use and ask a doctor ■ If condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, stop use and do not begin use of any other hydrocortisone product, unless you have consulted a doctor.

Keep out of the reach of children.

► If swallowed, get medical help or contact a Poison Control Center.

Drug Facts
(continued)

DIRECTIONS

For minor skin irritations and rashes, adults and children 2 years of age and older: apply to affected area not more than 3 to 4 times daily. For external anal itching: ■ Adults: when practical, cleanse the affected area with mild soap and warm water and rinse thoroughly ■ Gently dry by patting or blotting with toilet tissue or soft cloth before application of this product ■ Children: under 12 years of age, consult a doctor.

OTHER

INFORMATION

■ Do not use if seal is damaged or is not visible. To open, unscrew cap, pull tab to remove foil seal. ■ Store at room temperature ■ See crimp of tube or carton for Lot Number and Expiration Date

INACTIVE

INGREDIENT

Paraffin wax, Micro wax, Light liquid paraffin

OINTMENT 30 gms



POVIDONE-IODINE s-Oint

Drug Facts

ACTIVE INGREDIENT
Povidone-iodine, 5%
(0.5% available iodine)

PURPOSE

First aid antiseptic

INACTIVE INGREDIENTS

Cetostearyl alcohol, glycerin, Mineral oil, PEG-50 stearate, Polysorbate 60, Potassium iodate, Sodium hydroxide, Sorbitan monostearate, Water, White petrolatum

USES:

First aid to help prevent infection in ■ Minor Cuts ■ Scrapes ■ Burns

WARNINGS

Warnings: For external use only

Do not use in the eyes

Over large areas of the body

If you are allergic to povidone-iodine or any other ingredients in this preparation

Ask a doctor before use if you have

■ Deep or puncture wounds
■ Serious burns
■ Animal bites

Stop use and ask a doctor if

The condition persists or gets worse

You need to use this product for more than 1 week

Keep out of reach of children.

► If swallowed, get medical help or contact a Poison Control Center right away

Drug Facts
(continued)

DIRECTIONS

Clean the affected area

Apply a small amount of product (equal to the surface area of a fingertip) to the area 1 to 3 times daily may be covered with a sterile bandage

OTHER

INFORMATION

store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

OINTMENT 30 gms



MENTHOL & ZINC OXIDE s-Oint

Drug Facts

ACTIVE INGREDIENT
Menthol 0.44% w/w
Zinc Oxide 20.625% w/w

PURPOSE

External Analgesic
Skin Protectant

INACTIVE INGREDIENTS

Calamine, Glycerine, Phenol, Lanolin, Heavy Liquid Paraffin, Sodium Bicarbonate, Thymol, White Soft Paraffin

PURPOSE:

A moisture barrier that prevents and helps heal skin irritations from;

■ Urine ■ Diarrhea ■ Perspiration ■ Fistula drainage
■ Feeding tube site leakage ■ Wound drainage (peri-wound skin) ■ Minor burns ■ Cuts ■ Scrapes ■ Itching

WARNINGS

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

INDICATIONS & USAGE

■ Not for deep or puncture wounds. ■ Avoid contact with eyes. ■ If condition worsens or does not improve within 7 days, consult a doctor

DOSAGE & ADMINISTRATION:

Clean skin with mild skin cleanser

Pat dry or allow to air dry

Apply a thin layer of Calasoothe to reddened or irritated skin 2-4 times daily, or after each incontinent episode or diaper change to promote comfort and long lasting protection

Drug Facts
(continued)

DIRECTIONS

Cleanse skin gently with mild skin cleanser. Pat dry or allow to air dry. Apply a thin layer of Diaperene Adult Ointment to reddened or irritated skin 2-4 times daily, or after each incontinent episode or diaper change to promote comfort and long lasting protection.

OTHER

INFORMATION

Store between:
15°-30°C (59-86°F)



OINTMENT

30 gms



LANOLIN & PETROLATUM s-Oint

Drug Facts

ACTIVE INGREDIENT

Lanolin 15.5%
Petrolatum 53.4%

PURPOSE
Diaper rash ointment
Skin Protectant

INACTIVE INGREDIENTS

Cod liver oil (Contains Vitamin A & vitamin D), Fragrance, Light mineral oil, Microcrystalline wax, Paraffin

USES

Helps treat and prevent diaper rash;
Temporarily protects & Soothes:

- Cuts ■ Scrapes ■ Burns
- Temporarily protects and helps relieve chapped, chafed or cracked skin and lips ■ Protect chafed skin due to diaper rash and helps seal out wetness

WARNINGS

For external use only

- When using this product
- Do not get into eyes

Stop use and ask a doctor if

Condition worsens

Symptoms last more than 7 days or clear up and occur again within a few days

Do not use on

- Deep or puncture wounds ■ Animal bites ■ Serious burns.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts
(continued)

DIRECTIONS

For skin protectant use apply as needed

For diaper rash use:

Change wet and soiled diapers promptly
Cleanse the diaper area, and allow to dry
Apply ointment liberally as often as necessary, with each diaper change, especially at bedtime or anytime when exposure to wet diapers may be prolonged

OTHER

INFORMATION

Store between 20°C to 25°C (68° to 77°F)

Store upright when not in use

OINTMENT

30 gms



ZINC OXIDE s-Oint

Drug Facts

ACTIVE INGREDIENT

Zinc Oxide, 40%

PURPOSE
Skin protectant

USES

Helps treat and prevent diaper rash, Protects chafed skin due to diaper rash and helps seal out wetness

WARNINGS

For external use only

When using this product

- Avoid contact with the eyes

Stop use and ask a doctor if

Condition worsens or does not improve after 7 days

Keep out of reach of children.

To prevent accidental ingestion. If swallowed, seek medical help or call Poison Control Center immediately.

DIRECTIONS

Change wet and soiled diaper immediately, cleanse the diaper area and allow to dry, apply ointment liberally as often as necessary with each diaper change and especially when exposed to wet diapers for a prolonged period of time, such as bedtime

Drug Facts
(continued)

OTHER
INFORMATION

Storage temperature: not to exceed 30°C (86°F),

Remove foil seal from the tube's tip,

Use with infants, Children and Adults,

Will stain clothing and fabric

INACTIVE
INGREDIENTS

Castor Oil, Mineral Oil, Paraffin, Peruvian Balsam, White Petrolatum

SYRUP

118 ml



Diphenhydramine hydrochloride 12.5 mg **L-Oral**

ALLERGEX SYRUP

DRUG FACTS

ACTIVE INGREDIENT (IN EACH 5 ML)
Diphenhydramine HCl 12.5 mg

PURPOSE
Antihistamine

USES

■ Relieves symptoms due to hay fever or other upper respiratory allergies ■ Sneezing ■ Runny nose ■ Itchy, watery eyes ■ Itching of the nose or throat

WARNINGS

Do not use

- With any other product containing diphenhydramine, even one used on the skin ■ To make a child sleepy

Ask a doctor before use if the child has ■ Breathing problems such as chronic bronchitis ■ Glaucoma ■ Sodium-restricted diet

Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers.

When using this product ■ Marked drowsiness may occur

■ Sedatives and tranquilizers may increase drowsiness ■ Excitability may occur, especially in children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

DIRECTIONS

■ Do not take more than 6 doses in 24 hours ■ Take every 4 to 6 hours, or as directed by a doctor ■ Measure only with dosing cup provided. Do not use any other dosing device. ■ mL=milliliter ■ Keep dosing cup with product ■ Find the right dose on the chart below

Age (yr)	Dose (mL)
Children 6 to 11 years	5 mL to 10 mL
Children 2 to 5 years	Do not use unless directed by a doctor
Children under 2 years	Do not use

INACTIVE INGREDIENTS

Glycerin, Methylparaben Sodium, Propylparaben Sodium, Sodium Benzoate, Sucralose, sorbitol, Citric acid monohydrate, D&C red #33, FD&C red #40, flavor & purified water.

OTHER INFORMATION ■ Each 5 mL contains: Sodium 0.03 mg ■ Store between 20-25°C (68-77°F). Do not refrigerate. ■ Protect from light. Store in outer carton until contents are used.



SYRUP



118 ml

Bismuth subsalicylate 525 mg

L-Oral

STOMACH RELIEF SYRUP

DRUG FACTS	
ACTIVE INGREDIENT (IN EACH 15 ML DOSE)	PURPOSE
Bismuth subsalicylate 525 mg	Upset stomach reliever and Anti-diarheal
USES	
<ul style="list-style-type: none"> Diarrhea Upset stomach due to overindulgence in food and drink, including: <ul style="list-style-type: none"> Heartburn Indigestion Nausea Gas Belching Fulness 	
WARNINGS	
<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Contains salicylate. Do not take if you are: <ul style="list-style-type: none"> Allergic to salicylates (including aspirin) Taking other salicylate products. </p>	
Do not use if you have	
<ul style="list-style-type: none"> An ulcer A bleeding problem Bloody or black stool. 	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> Fever Mucus in the stool. 	
Ask a doctor or pharmacist before use if you are taking any drug for	
<ul style="list-style-type: none"> Anticoagulation (thinning the blood) Diabetes Gout Arthritis. 	
When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> Symptoms get worse or last more than 2 days Ringing in the ears or loss of hearing occurs Diarrhea lasts more than 2 days. 	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
DIRECTIONS	
<ul style="list-style-type: none"> Shake well before use Only use dose cup provided Adults and children 12 years and over: 15 mL (1 dose) every 1/2 hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea 15mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) Do not exceed 8 doses (120 mL) in 24 hours Use until diarrhea stops but not more than 2 days Children under 12 years: ask a doctor Drink plenty of clear fluids to help prevent dehydration caused by diarrhea. 	
OTHER INFORMATION	
<ul style="list-style-type: none"> Each 30 mL dose contains: magnesium 3 mg, sodium 2 mg Salicylate 230 mg Low sodium Protect from freezing Avoid excessive heat (34°F to 40°C) 	
TAMPER EVIDENT: Do not use if printed shrinkband is missing or broken.	
INACTIVE INGREDIENTS	
Benzoic acid, D&C Red No. 22, D&C Red No. 28, Flavor, Gellan gum, Magnesium aluminum silicate, Methylcellulose, Salicylic acid, Sodium salicylate, Sorbic acid, Sucrose, Water.	

SYRUP



118 ml

PARACETAMOL SYRUP

DRUG FACTS		
Active ingredient	Purpose	
Paracetamol 160 mg	Pain reliever - Fever reducer	
INACTIVE INGREDIENTS		
Artificial flavour, Methylparaben, Propylparaben, Potassium sorbate, Citric Acid, Monohydrate, Sorbitol, Polyethylene glycol, Sucrose, Povidone K30, Glycerin, FD&C Red #40, D&C Red #33 & Purified water.		
Temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> The common cold Flu Headache Sore throat Toothache Temporarily reduces fever. 		
WARNINGS		
<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: <ul style="list-style-type: none"> More than 5 doses in 24 hours, which is the maximum daily amount With other drugs containing acetaminophen </p> <p>Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> Skin reddening Blisters Rash </p>		
If a skin reaction occurs, stop use and seek medical help right away.		
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.		
Do not use <ul style="list-style-type: none"> With any other product containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist If your child is allergic to acetaminophen or any of the inactive ingredients in this product. 		
Ask a doctor before use if your child has liver disease		
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin		
When using this product, do not exceed recommended dose (see overdose warning)		
Stop use and ask a doctor if		
<ul style="list-style-type: none"> Pain gets worse or lasts for more than 5 days Fever gets worse or lasts for more than 3 days New symptoms occur Redness or swelling is present These could be signs of a serious condition.		
Keep out of the reach of children.		
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.		
DIRECTIONS		
<ul style="list-style-type: none"> This product does not contain directions or complete warnings for adult use Find right dose on chart. If possible, use weight to dose; otherwise, use age mL = milliliter If needed, repeat dose every 4 hours Do not use more than 5 doses in 24 hours Use enclosed dosing cup only. Do not use any other device. 		
Weight	Age	Dose
under 24 lbs	Under 2 years	Ask a doctor
24 to 35 lbs	2 to 3 years	5mL
36 to 47 lbs	4 to 5 years	7.5mL
48 to 59 lbs	6 to 8 years	10mL
60 to 71 lbs	9 to 10 years	12.5mL
72 to 95 lbs	11 years	15mL
OTHER INFORMATION		
<ul style="list-style-type: none"> TAMPER EVIDENT: Do not use this product if inner foil seal over mouth of the bottle is cut, torn, broken, or missing. Store at 20°-25°C (68°-77°F) This product is not the same concentration as Infants' Drops. For accurate dosing, follow the dosing instructions on this label. 		

SYRUP



118 ml

Acetaminophen 325mg
Dextromethorphan hydrobromide 10 mg
Phenylephrine hydrochloride 5 mg

L-Oral

COLD & FLU RELIEF SYRUP

DRUG FACTS	
ACTIVE INGREDIENT (IN EACH 5 ML)	PURPOSE
Acetaminophen 325 mg Dextromethorphan HBr 10 mg Phenylephrine HCl 5 mg	Pain reliever/fever reducer Cough suppressant Nasal decongestant
USES	
<ul style="list-style-type: none"> Temporarily relieves common cold/flu symptoms: <ul style="list-style-type: none"> Nasal congestion Cough due to minor throat and bronchial irritation Fever Headache Minor aches and pains Fatigue 	
WARNINGS	
<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if: <ul style="list-style-type: none"> Adult takes more than 4,000 mg of acetaminophen in 24 hours Child takes more than 5 doses in 24 hours Taken with other drugs containing acetaminophen Adult has 2 or more alcoholic drinks every day while using this product. </p> <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> Skin reddening Blisters Rash </p>	
If a skin reaction occurs, stop use and seek medical help right away.	
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Do not use <ul style="list-style-type: none"> With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (part of drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. If you have ever had an allergic reaction to this product or any of its ingredients. 	
Ask a doctor before use if you have <ul style="list-style-type: none"> Liver disease Heart disease Thyroid disease Diabetes High blood pressure Trouble urinating due to an enlarged prostate gland Cough that occurs with too much phlegm (mucus) Persistent or chronic cough such as occurs with smoking, asthma, or emphysema. 	
Ask a doctor or pharmacist before use if you are <ul style="list-style-type: none"> Taking the blood thinning drug warfarin 	
When using this product do not use more than directed	
Stop use and ask a doctor if <ul style="list-style-type: none"> You get nervous, dizzy or sleepless Pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults) Fever gets worse or lasts more than 3 days Redness or swelling is present New symptoms occur Cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. 	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
DIRECTIONS	
<ul style="list-style-type: none"> Take only as directed - see Overdose warning Only use the dose cup provided Do not exceed 4 doses per 24 hrs. 	
Adults & children 12 yrs & over	30 mL every 4 hrs
Children 6 to under 12 yrs	15 mL every 4 hrs
Children 4 to under 6 yrs	Ask a doctor
Children under 4 yrs	Do not use
OTHER INFORMATION	
<ul style="list-style-type: none"> Each 15mL contains: sodium 7mg Store at 20-25°C (68-77°F) 	
INACTIVE INGREDIENTS <ul style="list-style-type: none"> Sulfolated hydroxyanisole, Edetate disodium, FD&C yellow no. 6, Flavor, Glycerin, Mannitol, Monobasic sodium phosphate, Polyethylene glycol, Propylene glycol, Purified water, Saccharin sodium, Sucrose, Xanthan gum 	



SYRUP



118 ml

Acetaminophen 650mg
Dextromethorphan hydrobromide 30 mg
Doxylamine succinate 12.5 mg **L-Oral**

COLD & FLU RELIEF SYRUP-DS

DRUG FACTS	
ACTIVE INGREDIENTS (IN EACH 30 ML) Acetaminophen 650 mg Dextromethorphan HBr 30 mg Doxylamine succinate 12.5 mg	PURPOSE Pain relief / Fever reducer Cough suppressant Antihistamine
USES Temporarily relieves common cold/flu symptoms. ■ Cough due to minor throat and bronchial irritation ■ Sore throat ■ Headache ■ Minor aches and pains ■ Fever ■ Runny nose and sneezing	
WARNINGS Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ More than 4,000 mg of acetaminophen in 24 hours ■ With other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ Skin redness ■ Bleets ■ Rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Do not use ■ With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ If you have ever had an allergic reaction to this product or any of its ingredients	
Ask a doctor before use if you have ■ Liver disease ■ Glaucoma ■ Cough that occurs with too much phlegm (mucus) ■ A breathing problem such as emphysema or chronic bronchitis ■ Trouble urinating due to an enlarged prostate gland ■ Persistent or chronic cough as occurs with smoking, asthma, or emphysema ■ A sodium-restricted diet	
Ask a doctor or pharmacist before use if you have ■ Taking sedatives or tranquilizers ■ Taking the blood thinning drug warfarin	
When using this product ■ Excitability may occur, especially in children ■ Marked drowsiness may occur ■ Avoid alcoholic drinks ■ Be careful when driving a motor vehicle or operating machinery ■ Alcohol, sedatives, and tranquilizers may increase drowsiness	
Stop use and ask a doctor if ■ Pain or cough gets worse or lasts more than 7 days ■ Fever gets worse or lasts more than 3 days ■ Redness or swelling is present ■ New symptoms occur ■ Cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
DIRECTIONS ■ Take only as directed—see Overdose warning ■ Use dose cap ■ Do not exceed 6 doses per 24 hrs Adults & children 12 yrs & over 30 mL every 6 hrs Children 4 to under 12 yrs Ask a doctor Children under 4 yrs Do not use	
OTHER INFORMATION ■ Each 30 mL contains: sodium 38 mg ■ Store at 20-25°C (68-77°F)	
INACTIVE INGREDIENTS Alcohol, Anhydrous citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, Flavor, High fructose corn syrup, Polyethylene glycol, Propylene glycol, Purified water, Saccharin sodium, Sodium citrate	

SYRUP



118 ml

Dextromethorphan hydrobromide 10 mg
Guafenesin 100mg **L-Oral**

COUGH RELIEF SYRUP

DRUG FACTS	
ACTIVE INGREDIENT Each 5 mL (1 teaspoonful) contains: Dextromethorphan HBr10 mg Guafenesin100 mg	PURPOSE Cough Suppressant Expectorant
INACTIVE INGREDIENTS Acesulfame potassium, artificial cherry & vanilla flavor, aspartame, hydroxypropylcellulose, menthyl, methylparaben, potassium sorbate, purified water. Citric acid may be used to adjust pH.	
USES ■ Temporarily relieves cough ■ Helps loosen phlegm (mucus) and thin bronchial secretions to rid bronchial passageways of bothersome mucus	
WARNINGS Do not use If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have ■ A cough that occurs with too much phlegm (mucus) ■ A chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema	
Stop use and ask a doctor if ■ A cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. ■ These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Professional note: Guafenesin has been shown to produce a color interference with certain clinical laboratory determinations of 5-hydroxyindoleacetic acid (5-HIAA) and vanillylmandelic acid (VMA).	
DIRECTIONS ■ Take every 4 hours ■ Do not exceed 6 doses in 24 hours	
Age	Dose
Adults & children 12 years & over	10 mL (2 teaspoonfuls)
Children 6 years to under 12 years	5 mL (1 teaspoonful)
Children 2 years to under 6 years	2.5 mL (1/2 teaspoonful)
Children under 2 years	Ask a doctor
How Supplied: Dextromethorphan HBr and Guafenesin Oral Solution is a clear viscous liquid with a light cherry odor supplied in the following oral dosage forms: 5 mL unit dose, 10 mL unit dose in trays of 10 and 4 L oz. (118 mL) bottle	
Phenylethanolamine: contains phenylethanolamine 8.4 mg per teaspoonful (5 mL)	
Storage: Keep tightly closed. Store at controlled room temperature 20-25°C (68-77°F). [See USP] Protect from light.	

LOTION



118 ml

L-Topical


BENZOYL PEROXIDE LOTION-5

DRUG FACTS	
Active ingredient Benzoyl Peroxide 5%	Purpose Acne Treatment
Use For the treatment of acne. Dries, clears and helps keep skin clear of new acne pimples, blackheads and whiteheads.	
Warnings For external use only	
Ask a doctor before use if you are ■ Using other topical acne medications at the same time or immediately following use of this product may increase dryness or irritation of the skin. If this occurs, only one medication should be used unless directed by a doctor.	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Do not use if you ■ Have very sensitive skin. ■ Are sensitive to Benzoyl Peroxide.	
When using this product ■ Avoid unnecessary sun exposure and use a sunscreen. ■ Avoid contact with eyes, lips, and mouth. ■ Avoid contact with hair and dyed fabrics, which may be bleached by this product. ■ Skin irritation may occur, characterized by redness, burning, itching, peeling, or possibly swelling. Irritation may be reduced by using the product less frequently or in a lower concentration.	
Stop use and ask a doctor if ■ Irritation becomes severe.	
Directions ■ Wash hands with a non-medicated soap (contains no Benzoyl Peroxide). ■ Cleanse and dry face thoroughly 5 minutes before use. ■ Using fingertips apply evenly to affected areas. ■ Use both morning and night or as directed by a physician. ■ If dryness or peeling occurs, reduce application to once a day or every other day. ■ Fair-skinned individuals should begin with one application. ■ For more severe cases, and/or if substantial improvement is not apparent within 3 to 4 weeks, consult a health care practitioner.	
Inactive ingredients C13-14 Isoparaffin, Ethylhexylglycerin, Lanolin Oil, Laureth-4, Laureth-7, Magnesium Aluminum Silicate, Phenoxethanol, Polyacrylamide, Propylene Glycol, Water/Eau, Xanthan Gum.	
Other Information ■ Store at room temperature (59-86°F / 15-30°C)	



LOTION

118 ml



L-Topical

BENZOYL PEROXIDE LOTION-10

DRUG FACTS	
Active ingredient Benzoyl Peroxide 10%	Purpose Acne Treatment
Use For the treatment of acne.	
Warnings For external use only	
When using this product ■ Skin irritation and dryness is more likely to occur if you use another topical acne medication at the same time. ■ If irritation occurs, only use 1 topical acne medication at a time. ■ Avoid unnecessary sun exposure and use a sunscreen. ■ Avoid contact with the eyes, lips and mouth. ■ Avoid contact with hair or dyed fabrics, which may be bleached by this product. Skin irritation may occur, characterized by redness, burning, itching, peeling, or possibly swelling. ■ Irritation may be reduced by using the product less frequently or in a lower concentration.	
Do not use if you have very sensitive skin or sensitive to benzoyl peroxide.	
OTHER INFORMATION Keep tightly closed. Protect from excessive heat (40°/ 140° F) and protect from freezing	
STOP USE Stop use and ask a doctor if irritation or sensitivity develops.	
KEEP OUT OF REACH OF CHILDREN Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
DIRECTIONS Important to use on a clean face. Cover the affected area with a thin layer 1 to 3 times per day. Because of excessive drying of the skin may occur, start with 1 application per day then gradually increase to 2 applications per day 1 morning and one evening if needed, or as directed by a doctor. If dryness or peeling occurs, reduce use to once a day or once every other day. If going outside, use a sunscreen. If irritation or sensitivity develops stop using both products and ask a doctor.	
INACTIVE INGREDIENTS Water (Aqua), Glycerin, Propylene Glycol, Hydroxyethyl Acrylate/Sodium Acryloyldimethyl Taurate Copolymer, Sulfur, PEG-8 SMDI Copolymer, Lactobacillus Ferment, Glycolic Acid, Lactic Acid, Magnesium Aluminum Silicate, Citric Acid, Xanthan Gum, Potassium Hydroxide, Phenoxethanol, Ethylhexylglycerin, Hydrolyzed Algin, Zinc Sulfate, Disodium EDTA, Citrus Aurantium Bergamia (Bergamot) Fruit Oil, Citrus Grandis (Grapefruit) Peel Oil, Citrus Aurantium Dulcis (Orange) Peel Oil, Citrus Tangerina (Tangerine) Peel Oil	

LOTION

118 ml



L-Topical

CALAMINE & ZINC OXIDE LOTION

DRUG FACTS	
Active ingredient Calamine 8% and Zinc Oxide 8%	Purpose Skin protectant
INDICATIONS & USAGE Dries the oozing and weeping o poison ivy, poison oak, and poison sumac.	
WARNINGS ■ For external use only. Use only as directed. ■ Avoid contact with eyes and mucous membranes. ■ Ask a doctor before using on children 6 months of age.	
KEEP OUT OF REACH OF CHILDREN In case of accidental ingestion, seek professional assistance or contact a Poison Control center immediately.	
DOSAGE & ADMINISTRATION ■ Adults and children 2 years of age and older: shake well before using. Cleanse the skin with soap and water and let it dry before each use. Apply lotion to the affected area using a cotton or soft cloth, as often as needed for comfort. ■ Children under 6 months of age: Consult a doctor before use.	
INACTIVE INGREDIENT Bentonite magna, Calcium hydroxide, Glycerin, Purified water.	
Other information Store at room temperature 15-30°C (59-86°F).	

LOTION

118 ml



L-Topical

CALAMINE & PRAMOXINE HCL LOTION

DRUG FACTS	
Active ingredient Calamine 8% Pramoxine HCl 1%	Purpose Skin protectant Topical analgesic
USES ■ Temporarily relieves pain and itching associated with: ■ Rashes due to poison ivy, ■ Poison oak or poison sumac ■ Insect bites ■ Minor skin irritation ■ Minor cuts ■ Dries the oozing and weeping of poison ivy, poison oak and poison sumac	
WARNINGS For external use only. When using this product do not get into eyes Stop use and ask a doctor if ■ Condition worsens or does not improve within 7 days ■ Symptoms persist for more than 7 days or clear up and occur again within a few days	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
DIRECTIONS ■ Shake well before use ■ Adults and children 2 years of age and older: apply to affected area not more than 3 to 4 times daily ■ Children under 2 years of age: ask a doctor	
OTHER INFORMATION Store at 20° to 25°C (68° to 77°F)	
INACTIVE INGREDIENTS SD Alcohol 38-B, Camphor, Diazolidinyl urea, Fragrance, Hydroxyethylcellulose, Methylparaben, Polysorbate 80, Propylene glycol, Propylparaben, Purified water, Xanthan gum	



GEL

118 ml



L-Topical

**WHITE
PETROLATUM GEL**

DRUG FACTS	
Active ingredient WHITE PETROLATUM USP (30%)	Purpose Skin protectant
Uses ■ Temporarily protects minor Burns ■ Temporarily protects and helps relieve chapped or cracked skin ■ Helps protect from the drying effects of wind and cold weather	
WARNINGS For External Use Only When using this product do not get into eyes.	
Stop use and ask a doctor if ■ Condition worsens ■ Symptoms last more than 7 days or clear up and occur again within a few days.	
Do not use on ■ Deep or puncture wounds ■ Animal bites ■ Serious burns	
Keep out of reach of children. If swallowed get medical help or contact a Poison Control Center right away.	
Directions Apply as needed.	
Questions? Call toll-free 1-800-572-6632, Weekdays 7:00 AM - 5:30 PM EST.	
Inactive ingredients Water (Aqua), Caprylic/Capric Triglyceride, Stearic Acid, Glycerin, Sodium Hydroxypropyl Starch Phosphate, Glycol Stearate, PEG-100 Stearate, Glyceryl Stearate, Cetyl Alcohol, Phenoxyethanol, Methylparaben, Isopropyl Myristate, Propylparaben, Stearamide AMP, Acrylates/C10-30 Alkyl Acrylate Crosspolymer, Disodium EDTA, Potassium Hydroxide, Cedrol, Tocopheryl Acetate, Titanium Dioxide (CI 77891).	

GEL

118 ml



L-Topical

**POVIDONE-
IODINE SCRUB**

DRUG FACTS	
Active ingredient Povidone-iodine, 7.5% (0.75% available iodine)	Purpose Antiseptic
Uses ■ For preparation of the skin prior to surgery ■ Helps to reduce bacteria that potentially can cause skin infection ■ For handwashing to reduce bacteria on the skin ■ Significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care	
Warnings For external use only Do not use this product in the eyes	
When using this product ■ Prolonged exposure to wet solution may cause irritation or, rarely, severe skin reactions ■ In pre-operative prepping avoid "pooling" beneath the patient	
Stop use and ask a doctor ■ If irritation and redness develop ■ In rare instances of local irritation or sensitivity	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions A. Surgical hand scrub: ■ Wet hands with water ■ Spread about 5 cc (1 teaspoonful) of Scrub over both hands and forearms ■ Without adding more water, scrub thoroughly for 2 1/2 to 3 minutes ■ Use a sponge if desired. Clean thoroughly under fingernails. ■ Add a little water and develop copious suds. Rinse thoroughly under running water. ■ Repeat the entire procedure using another 5 cc of Scrub B. Antiseptic hand wash: ■ Wet hands with water and pour about 5 cc of Scrub on hands ■ Rub hands vigorously together for at least 15 seconds, covering all surfaces ■ Rinse and dry with a disposable towel C. Patient pre-operative skin preparation: ■ Single use only ■ Wet skin with water ■ Apply Scrub (1 cc is sufficient to cover an area of 20-30 square inches), develop lather and scrub thoroughly for about 5 minutes ■ Rinse off using sterile gauze saturated with water ■ The area may then be painted with BETADINE solution and allowed to dry	
Other information ■ Store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ Store in original container	
Inactive ingredients Ammonium nonoxonyl-4 sulfate, nonoxonyl-9, purified water, sodium hydroxide	
Questions? Call toll-free 1-800-572-6632, Weekdays 7:00 AM - 5:30 PM EST.	

DROPS

15 ml



L-Oral

Simethicone 20mg

GAS RELIEF DROPS

DRUG FACTS		
ACTIVE INGREDIENT (IN EACH 0.3 ML)	PURPOSE	
Simethicone 20 mg	Anti-flatulence	
USES Relieves the symptoms of gas frequently caused by air swallowing or certain formulas or foods		
WARNINGS When using this product Do not exceed 12 doses per day Keep out of reach of children. In case of overdose get medical help or contact a poison control center immediately.		
DIRECTIONS ■ Shake well before using ■ All dosages may be repeated as needed, after meals and at bedtime ■ Fill enclosed dropper to recommend dosage level ■ Dispense liquid slowly into baby's mouth, toward the inner cheek ■ May mix with 1 oz. of cool water, infant formula or other suitable liquids ■ Clean dropper after each use and close the bottle to maintain child resistance		
Age (yr)	Weight (lb)	Dose
Infants under 2	Under 24	0.3 mL
Children over 2	Over 24	0.6 mL
OTHER INFORMATION Store at room temperature		
INACTIVE INGREDIENTS Benzoic acid, Flavor, Magnesium aluminum silicate, Purified water, Simethicone emulsion, Sorbitol, Xanthan gum		



Rubbing Alcohol

447 ml



Carboxy methylcellulose sodium 10mg

L-Topical

ISO PROPYL ALCOHOL

Drug Facts	
Active ingredient	Purpose
Isopropyl alcohol (50%).....	First Aid Antiseptic
Uses first aid to help prevent the risk of infection in minor cuts, scrapes and burns	
Warnings For external use only	
<ul style="list-style-type: none"> ■ Flammable, keep away from fire and flame ■ If taken internally, serious gastric disturbances will result 	
Ask a doctor before use in you have ■ deep or puncture wounds, animal bites or serious burns	
When using this product ■ do not get into eye ■ do not apply over large areas of the body ■ do not use longer than one week unless directed by a doctor ■ do not use otherwise than directed	
Stop use and ask a doctor if	
■ Condition persists or gets worse	
Keep out of reach of children. In case of ingestion, get medical help or contact a Poison Control Center immediately.	
Directions ■ clean affected area ■ apply a small amount of this product on the area 1 to 3 times daily ■ if bandaged, let dry first ■ may be covered with a sterile bandage	
Other information ■ store at room temperature ■ does not contain, nor is intended as a substitute for grain or ethyl alcohol	
Inactive ingredient Purified water	

Sterile Wipe



SWIPE

Drug Facts
Active ingredients: Isopropyl Alcohol 70%, Chlorhexidine 2%
INDICATIONS & PURPOSE : For preparation of skin prior to an injection; for cleaning medical devices.
DIRECTIONS: Tear the packet open, Wipe injection site vigorously and discard.
WARNINGS: For External Use Only
DO NOT USE ■ With electrocautery procedures ■ In the eyes (If contact occurs, flush eyes with water) Stop Use and ask a doctor if irritation and redness develop.
Keep out of reach of children If swallowed, get medical help or contact a Poison Control Center right away.
Store at room temperature
Manufactured by : Global Pharma Healthcare Pvt. Ltd., A-9, SIDCO Pharmaceutical Complex, Alathur-603 110 - INDIA. www.global-pharma.com

EYE DROPS

15 ml



POLYVINYL ALCOHOL

i-Drops

DRUG FACTS

ACTIVE INGREDIENT PURPOSE
Polyvinyl Alcohol 1.4% Eye lubricant

Uses

For use as a lubricant to prevent further irritation or to relieve dryness of the eye(s).

Warnings

- Do not use if seal on the bottle neck is broken or missing.
- Do not use if solution changes color or becomes cloudy.
- To avoid contamination, do not touch tip of container to any surface.
- Replace cap after using.

Stop use and ask a doctor if you experience

- Eye pain, ■ Changes in vision,
- Continued redness or irritation of the eye, or if the condition worsens or persists for more than 72 hours.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center.

DRUG FACTS

(continued)

Directions

Instill 1 or 2 drops in the affected eye(s) as needed.

Other information

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].
Store away from heat.
Protect from freezing.
Keep tightly closed. **RETAIN THIS CARTON FOR FUTURE REFERENCE.**

Inactive ingredients

Benzalkonium Chloride 0.005%, Edetate Disodium, Sodium Chloride, Sodium Phosphate Dibasic, Sodium Phosphate Monobasic, Purified water, Sodium Hydroxide and/or Hydrochloric Acid to adjust pH.

Questions?

Call toll-free 1-800-572-6632, weekdays, 7:00 AM - 5:30 PM EST.



EYE DROPS

15 ml



CARBOXY METHYLCELLULOSE SODIUM i-Drops

DRUG FACTS

ACTIVE INGREDIENT PURPOSE
Carboxymethylcellulose Eye lubricant
Sodium 10mg in 1 ml.

Uses

For use as a protectant against further irritation or to relieve dryness of the eye.
For the relief of discomfort due to minor irritations of the eye or to exposure to wind, sun or screen time.

Warnings

For external use only.
Do not use product if solution changes color or becomes cloudy.
Stop use and ask a doctor if you experience
■ Eye pain, ■ Changes in vision, ■ Continued redness or irritation of the eye or if the condition worsens or persists for more than 72 hours.
When using this product:
■ To avoid contamination, do not touch tip of container to any surface.
■ Replace cap after using. Keep container tightly closed.
■ Remove contact lens before using ▶

DRUG FACTS

(continued)
Keep out of reach of children. If accidentally swallowed, get medical help or contact a Poison Control Center immediately.

Directions

Instill 1 or 2 drops in the affected eye(s) as needed.

Other information

Tamper Evident. Do not use this product if neckband is missing or broken.
RETAIN THIS CARTON FOR FUTURE REFERENCE.
Store at 15°-30°C (59° to 86°F).

Inactive ingredients

Boric Acid, Potassium Chloride, Sodium Chloride, Calcium Chloride, Dihydrate, Magnesium Chloride, Sodium Chloride, Sodium Hydroxide and Water for Injection.

Questions?

Call toll-free 1-800-572-6632, weekdays, 7:00 AM - 5:30 PM EST.

EYE DROPS

15 ml



POLYVINYL ALCOHOL & POVIDONE i-Drops

DRUG FACTS

ACTIVE INGREDIENT PURPOSE
Polyvinyl alcohol 0.5% Lubricant
Povidone 0.6% Lubricant

Uses

■ For use as a protectant against further irritation or to relieve dryness of the eye
■ For the temporary relief of discomfort due to minor irritations of the eye, or to exposure to wind or sun

Warnings

For external use only.
Do not use this product if
■ Solution changes color or becomes cloudy
When using this product
■ Remove contact lens before using
■ To avoid contamination, do not touch tip of container to any surface
■ Replace cap after using. Keep container tightly closed.
Stop use and ask a doctor if you experience
■ Eye pain, ■ Changes in vision, ■ Continued redness or irritation of the eye or if the condition worsens ▶

DRUG FACTS

(continued)
or persists for more than 72 hours
Keep out of the reach of children.
If accidentally swallowed, get medical help or contact a Poison Control Center.

Directions

Instill 1 or 2 drops in the affected eye(s) as needed.

Other information

Store at 15°-30°C (59°-86°F)
Inactive ingredients
Benzalkonium chloride, Dextrose, Edetate disodium, Potassium chloride, Purified water, Sodium bicarbonate, Sodium chloride, Sodium citrate, Sodium phosphate dibasic, Sodium phosphate monobasic.

Questions?

Call toll-free 1-800-572-6632, weekdays, 7:00 AM - 5:30 PM EST.

EYE DROPS

15 ml



POLYETHYLENE GLYCOL & PROPYLENE GLYCOL i-Drops

DRUG FACTS

ACTIVE INGREDIENT PURPOSE
Polyethylene Glycol 400 0.4% Lubricant
Propylene Glycol 0.3% Lubricant

Uses

For the temporary relief of burning and irritation due to dryness of the eye.

Warnings

For external use only.
Do not use
if this product changes color or becomes cloudy
if you are sensitive to any ingredient in this product
When using this product
■ Do not touch tip of container to any surface to avoid contamination
■ Replace cap after each use
Stop use and ask a doctor if
■ You feel eye pain
■ Changes in vision occur
■ Redness or irritation of the eye(s) gets worse, persists or lasts more than 72 hours. ▶

DRUG FACTS

(continued)
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center.

Directions

Instill 1 or 2 drops in the affected eye(s) as needed.

Other information

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].
Inactive ingredients
Purified Water, Boric Acid, Hypromellose, Sodium Chloride, Potassium Chloride, Zinc Chloride, Calcium Chloride Dihydrate, Magnesium Chloride Hexahydrate. May contain Hydrochloric Acid and/or Sodium Hydroxide to adjust pH; Preservative, Benzalkonium Chloride.

Questions?

Call toll-free 1-800-572-6632, weekdays, 7:00 AM - 5:30 PM EST.



EYE DROPS 15 ml



GLYCERIN *i-Drops*

DRUG FACTS

ACTIVE INGREDIENT	PURPOSE
Glycerin.....1.0%	lubricant

Uses

- For use as a lubricant to prevent further irritation or to relieve dryness of the eye.
- For the temporary relief of discomfort due to minor irritations of the eye or to exposure to wind or sun.

Warnings

For external use only

Do not use if solution changes color or becomes cloudy

When using this product

- To avoid contamination, do not touch tip of container to any surface
- replace cap after using

Stop use and ask a doctor if

- You experience eye pain
- You experience changes in vision
- You experience continued redness or irritation of the eye
- The condition worsens or persists for more than 72 hours.

Keep out of reach of children.

If product is swallowed, get medical help or contact a Poison Control Center.

DRUG FACTS

(continued)

Directions

Instill 1 or 2 drops in the affected eye(s) as needed.

Other information

store in the refrigerator do not freeze store in dark, cool place away from sunlight once open, discard container and any contents after 30 days use before expiration date marked on container prior to drop instillation, contact lenses should be removed and reinserted after 10 minutes

Inactive ingredients

Boric acid, Citric acid, Hypromellose, Potassium bicarbonate, Purified benzyl alcohol, Sterile water.

Questions?

Call toll-free 1-800-572-6632, weekdays, 7:00 AM - 5:30 PM EST.

EYE DROPS 15 ml



GLYCERIN, HYPROMELLOSE & POLYETHYLENE GLYCOL *i-Drops*

DRUG FACTS

ACTIVE INGREDIENT	PURPOSE
Glycerin 0.2%	Lubricant
Hypromellose 0.2%	Lubricant
Polyethylene Glycol 400 1%	Lubricant

Uses

- For use as a protectant against further irritation or to relieve dryness of the eye.
- For the temporary relief of discomfort due to minor irritations of the eye, or to exposure to wind or sun.

Warnings

Do not use this product if solution changes color or becomes cloudy

Stop use and ask a doctor if you experience

- Eye pain,
- Changes in vision,
- Continued redness or irritation of the eye, or if the condition worsens or persists for more than 72 hours.

When using this product

To avoid contamination, do not touch tip of container to any surface. Replace cap after using. Keep container tightly. Remove contact lens before using

DRUG FACTS

(continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center.

Directions

Instill 1 or 2 drops in the affected eye(s) as needed.

Other information

Do not use if the tamper-resistant bottle cap is broken or opened. RETAIN THIS CARTON FOR FUTURE REFERENCE store between 15° to 30°C (59°F to 86°F).

Inactive ingredients

Aminoacetic acid, Ascorbic acid, Benzalkonium chloride, Boric acid, Disodium phosphate, Magnesium chloride, Potassium chloride, Purified water, Sodium borate, Sodium chloride, Sodium citrate, Sodium lactate.

Questions?

Call toll-free 1-800-572-6632, weekdays, 7:00 AM - 5:30 PM EST.

Corporate Office:

No. 2A, 4th Street, Ganga Nagar, Kodambakkam, Chennai - 600 024. INDIA +91 44 2472 6001

Factory:

A-9, SIDCO Pharmaceutical Complex, Alathur, Thiruporur - 603 110. INDIA. +91 44 2744 4408

E-mail: venkatesh@global-pharma.com | prarthana@global-pharma.com

www.global-pharma.com