



EUROPEAN UNION  
**GMP**  
APPROVED



**Globela<sup>®</sup> Pharma Pvt. Ltd.**

*"Every Life Is Precious"*

# PRODUCT CATALOGUE



## Experienced Manufacturer for Finished & Semi Finished Formulation WHO GMP, ISO Certified, MNC Accredited Manufacturing Company

- ✔ Dedicated **Oral Solid Dosage** Facilities (Tablets, Capsules, Dry Oral Powder)
- ✔ Dedicated Facilities for **Anti-Cancer** (Cytotoxic) Oral finished formulations
- ✔ Dedicated **Cephalosporins** Facilities (Tablets, Capsules, Dry Oral Powder)
- ✔ **Direct Compressed Granules** and Pellets Unit equipped with latest Technology
- ✔ DSIR Approved & in House **R&D** Center
- ✔ **API & Intermediate** Manufacturing

📍 Plot-357-358, Road-3, Sachin G.I.D.C, [info@globelapharma.com](mailto:info@globelapharma.com)  
Surat - 394 230, Gujarat, India. [sales@globelapharma.com](mailto:sales@globelapharma.com)

☎ +91 - 261- 6158000-99

🌐 [www.globelapharma.com](http://www.globelapharma.com)



## SERVICES OFFERED

### Finished Solid Oral Dosage

We at Globela Pharma are committed for high quality pharmaceutical formulation in various forms with a team of qualified & experienced technocrats in R & D, Q.A, Q.C & Production division. We aim to meet our client expectations by our services.



01

Contract Manufacturing & Formulation Development

02

Exhibit batch & F&D batch formulation

03

Technology Transfer & Dossier Preparation

04

API / Vitamins manufacturer & Intermediates manufacturer



### Oncology (Anti Cancer)

We have dedicated WHO-GMP approved facility for Oncology solid oral dosages along with dedicated utilities and team of qualified & experienced technical staff.

### Taste Masked and Direct Compressed Granules

Globela Group is one of the few companies with the Manufacturing capability of Pharmaceutical Formulation Intermediates (PFIs). We offer our customers all three components API, DCs & FDs of the pharmaceutical Manufacturing to enhance flexibility and efficiency

### Direct Compressed Granules

Savings in capital expenditures & technical resources.

Increased asset utilisation

Lower testing costs

Save Development cost & Proven Technology back up

### Production Capacity/Month

● Tablets (Conventional)	300 Million
● Tablets (Effervescent)	50 Million
● Capsules	150 Million
● ORS (Sachet)	15 Million
● Dry Syrup	8 Million
● Pellets	70 MT

## THERAPEUTIC SEGMENTS

- Anti-Cancer (Oncology)
- Antibiotics & Antibacterials
- Analgesic & Antispasmodic
- Anti Hypertensive
- Immunosuppressants
- Antimalarials
- Anti Cold & Anti Histamine
- Anti-Diabetics
- Nephrology & Urology
- Anti-Fungal & Anthelmintics
- Vitamins, Minerals & Electrolytes
- Gastrology, GI Tract & Many more...

## CATEGORY



ANALGESIC & ANTISPASMODICS	1
GENERAL ANTIBIOTICS (ANTI BACTERIAL)	6
ANTI COLD & ANTI HISTAMINE	10
ANTI DIABETICS	13
ANTI MALARIAL	17
ANTI PSYCHOTIC & CNS	19
ANTIFUNGAL & ANTHELMINTICS	22
ARV PRODUCT	24
CARDIOVASCULAR	30
ELECTROLYTES & MINERALS	38
ERECTILE DYSFUNCTION	39
GASTROLOGY	40
NEPHROLOGY, UROLOGY & HEPATIC	44
ONCOLOGY & IMMUNOSUPPRESSANTS	46
STEROIDAL ANTI INFLAMMATORY	55
VITAMINS & MINERALS	56
LAXATIVE	58
EFFERVESCENT TABLETS & NUTRACEUTICAL	59
CEPHALOSPORINS & BETALACTAM	63



## ANALGESIC & ANTISPASMODICS



### ACECLOGLOB

**Composition :** ACECLOFENAC TABLETS 100 MG

**Indication :** Aceclofenac Tablet is indicated for Rheumatoid arthritis, ankylosing spondylitis, osteoarthritis and peri-arthritis of scapulo-humeral, lumbago, ischiodynia, pain caused by non-articular rheumatism.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### ACECLOGLOB-P

**Composition :** ACECLOFENAC & PARACETAMOL TABLETS 100+500 MG

**Indication :** Aceclofenac and paracetamol tablet is used for fever, joint pain, toothache, cold, headache, ear pain & other conditions.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### DICLOGLOBE

**Composition :** DICLOFENAC SODIUM TABLETS 50/100 MG

**DICLOFENAC POTASSIUM TABLETS 50/100 MG**

**Indication :** Relief of all grades of pain and inflammation such as rheumatoid arthritis, osteoarthritis, ankylosing spondylitis.

**Pack :** 1 x 10, 10 x 10 and 1 x 12 Blister



### DICLOGLOB P

**Composition :** PARACETAMOL & DICLOFENAC POTASSIUM TABLETS 325+50 MG

**Indication :** Tooth extraction, pericoronitis, ligament tear, mild to moderate pains.

**Pack :** 1 x 10, 10x10 Blister



### DICLOGLOB B

**Composition :** DICLOFENAC VIT. B1 + VIT. B6 + VIT. B12 TABLETS

**Indication :** Relief of all grades of pain and inflammation such as rheumatoid arthritis, osteoarthritis, ankylosing spondylitis.

**Pack :** 1 x 10, 2 x 10 and 3 x 10 Blister



### IBUGLOB

**Composition :** IBUPROFEN TABLETS 200/400/600/800MG  
**Indication :** Analgesics, antipyretics, nonsteroidal anti-inflammatory drugs, drugs used to treat out and disease-modifying antirheumatic drugs.  
**Pack :** 1 x 10, 2 x 10 and 10 x 10 Blister



### IBURICH

**Composition :** IBUPROFEN PARACETAMOL & CAFFEINE CAPSULES 200+325+30 MG  
**Indication :** Analgesics, antipyretics, nonsteroidal anti-inflammatory drugs, drugs used to treat out and disease-modifying antirheumatic drugs.  
**Pack :** 1 x 10, 2 x 10 and 10 x 10 Blister



### ACECLOGLOB-PC

**Composition :** ACECLOFENAC + CHLORZOAZONE + PARACETAMOL TABLETS 100+250+500 MG  
**Indication :** Acetaminophen or paracetamol is used for fever by the patient, or the doctor prescribes it for a headache, body pain and fever.  
**Pack :** 1 x 10, 3 x 10, 10x1 and 20x10 Blister



### GLOCOXIB

**Composition :** CELECOXIB CAPSULES BP 100/200/400 MG  
**Indication :** Celecoxib is used to treat pain or inflammation caused by many conditions such as arthritis, ankylosing spondylitis and menstrual pain.  
**Pack :** 3 x 10 and 10 x 10 Blister



### DECHLOZ MR

**Composition :** DICLOFENAC POTASSIUM + PARACETAMOL + CHLORZOAZONE TABLET 50+325+250 MG  
**Indication :** It is used to treat muscle aches, backaches, dental pain, menstrual cramps and sports injuries. It also reduces pain, swelling and joint stiffness caused by arthritis.  
**Pack :** 1x10, 2x10, 3x10 and 10 x 10 Blister



### DROTAGLOB

**Composition :** DROTAVERINE HCL TABLETS 40/80 MG  
**Indication :** The medication works by inhibiting phosphodiesterase in the body with no anticholinergic effects. It can be used to treat pain and dysfunction caused by smooth muscle spasms. It is mainly prescribed in the treatment of intestinal biliary and renal colics, irritable bowel syndrome and uterine spasms.  
**Pack :** 10 x 10 Blister



## ETORIGLOB

**Composition :** ETORICOXIB TABLETS 60/90/120 MG

**Indication :** Arthritis, spondylitis, post surgical pain management & acute low back pain.

**Pack :** 2x10, 3x10 and 10x10 Blister



## INDOMETHACIN

**Composition :** INDOMETHACIN CAPSULES 25/50 MG

**Indication :** It is used to reduce swelling and to treat pain. It can also be used for painful joint and muscular problems such as arthritis, tendinitis, bursitis and out.

**Pack :** 2 x 10 and 10x10 Blister



## ASPIRIN

**Composition :** ASPIRIN DISPERSIBLE TABLETS 75/81/100/300 MG

**Indication :** Aspirin is a type of nonsteroidal anti-inflammatory drug (NSAID) that can treat mild to moderate pain, inflammation or arthritis. It also lowers your risk of heart attack, stroke or blood clot.

**Pack :** 1x10, 3 x 10 and 10x10 Blister/Strip



## ORPHEGLOB

**Composition :** ORPHENADRINE CITRATE & PARACETAMOL TABLETS 35+350 MG

**Indication :** Orphenadrine is used to relax muscles and relieve pain. Discomfort caused by muscle injuries may cause drowsiness. Do not drive until you know how it affects you.

**Pack :** 1x10, 3x10 and 10x10 Blister



## P-FLU

**Composition :** PARACETAMOL & CAFFEINE TABLETS 500+30/500+65 MG

**Indication :** Paracetamol & caffeine Tablets are a mild analgesic and antipyretic formulated to give extra pain relief.

**Pack :** 3x10 and 10x10 Blister



### TAPZGLO

**Composition :** TAPENTADOL HYDROCHLORIDE TABLETS 50/75/100 MG  
**Indication :** It is indicated for the relief of moderate to severe acute pain in adults, which can be adequately managed only with opioid analgesics.  
**Pack :** 2x10 and 10x10 Blister



### GLONIM-MR

**Composition :** NIMESULIDE & PARACETAMOL TABLETS 100+325 MG  
**Indication :** Osteoarthritis/rheumatoid arthritis, musculo skeletal injuries, tendonitis & bursitis low back pain/ankylosing spondylitis, dysmenorrhea  
**Pack :** 10x 10 Blister



### IBUGLOB-P

**Composition :** IBUPROFEN & PARACETAMOL TABLETS 400+325 MG  
**Indication :** Analgesics, antipyretics, nonsteroidal anti-inflammatory drugs, drugs used to treat out and disease-modifying antirheumatic drugs.  
**Pack :** 1x10 and 10x10 Blister



### GLONIM-100

**Composition :** NIMESULIDE TABLETS 50/100 MG  
**Indication :** Low back pain / ankylosing spondylitis, antipyretic, dysmenorrhea.  
**Pack :** 10x10 and 50x10 Blister



### ACETAMINOPHEN, ASPIRIN & CAFFEINE

**Composition :** ACETAMINOPHEN, ASPIRIN & CAFFEINE TABLETS 250+250+65 MG  
**Indication :** It is used for the temporary relief of pain from conditions such as muscle aches, toothaches, menstrual cramps, or headaches (including migraine).  
**Pack :** 1x10, 10x10 Blister



### MELOXICAM

**Composition :** MELOXICAM TABLETS 7.5/15 MG  
**Indication :** Meloxicam is used to relieve pain, tenderness, swelling, and stiffness caused by osteoarthritis (arthritis caused by a breakdown of the lining of the joints) and rheumatoid arthritis (arthritis caused by swelling of the lining of the joints).  
**Pack :** 3x10 and 10x10 Blister



## TRAMADOL

**Composition :** TRAMADOL CAPSULES 100/200 MG

**Indication :** Tramadol is an opioid agonist that may be used to treat moderate to moderately severe chronic pain in adults. It acts in the central nervous system (CNS) to relieve pain.

**Pack :** 3x10 and 10 x 10 Blister



## SERRATIOPEPTIDASE

**Composition :** SERRATIOPEPTIDASE TABLETS 5/10/20/40 MG

**Indication :** Serratiopeptidase tablets are indicated for the reduction of inflammation, edema, and pain associated with post-surgical recovery, traumatic injuries, and inflammatory conditions such as sinusitis, arthritis, and chronic bronchitis.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister/Strip



## MEFANEMIC

**Composition :** MEFANEMIC TABLETS 250 MG/500 MG

**Indication :** Mefenamic acid is a type of NSAIDs. Mefenamic acid is used to relieve mild to moderate pain, including menstrual pain (pain that happens before or during a menstrual period). It works by stopping the body's production of a substance that causes pain, fever, and inflammation.

**Pack :** 1x10, 3x10 and 10x10 Blister





## GENERAL ANTIBIOTICS (ANTI BACTERIAL)



### AZBACT SUSPENSION

**Composition :** AZITHROMYCIN FOR ORAL SUSPENSION 200 MG / 5 ML

**Indication :** To treat bacterial infections, acute bacterial sinusitis, acute otitis media, community-acquired pneumonia, pharyngitis / tonsillitis, uncomplicated skin & skin structure, urethritis, cervicitis.

**Pack :** 15 ml / 30 ml HDPE Bottle

**Flavor :** Banana / Apple / Strawberry



### TERBINAFINE

**Composition :** TERBINAFINE TABLETS 125/250 MG

**Indication :** Terbinafine tablets are indicated for the treatment of onychomycosis (fungal infections of the toenails or fingernails) and dermatophyte infections such as tinea pedis (athlete's foot), tinea corporis (ringworm of the body), and tinea cruris (jock itch).

**Pack :** 2 x 7 Blister



### AZBACT

**Composition :** AZITHROMYCIN TABLETS 100 / 250 / 500 MG

**Indication :** Upper & lower RTIs, SSTIs, urogenital infections, gonorrhoea.

**Pack :** 1 x 3, 1 x 6 and 1 x 10 Blister



### BACIFLOX OZ

**Composition :** OFLOXACIN & ORNIDAZOLE TABLETS 200 + 500 MG

**Indication :** Community-acquired pneumonia, uncomplicated skin & skin structure infection, uncomplicated & complicated UTI.

**Pack :** 10 x 10 Blister



### DOXYGLOBE

**Composition :** DOXYCYCLINE HYCLATE CAPSULES 50 / 100 MG

**Indication :** Respiratory tract infection, pneumonia, chronic bronchitis, sinusitis, urinary tract infection, sexually transmitted diseases, dermatological & ophthalmic infections, rickettsial infections.

**Pack :** 1 x 10 and 10 x 10 Blister



## GATIFLOXACIN

**Composition :** GATIFLOXACIN TABLETS 200 / 400MG

**Indication :** Gatifloxacin is indicated in patients >18 years of age for the treatment of the following infections when caused by susceptible bacteria community acquired pneumonia caused by streptococcus pneumonia, haemophilus influenzae, moraxella catarrhalis.

**Pack :** 1 x 10 and 10x10 Blister



## G-LEVO

**Composition :** LEVOFLOXACIN TABLETS 250 / 500 / 750 MG

**Indication :** Community-acquired pneumonia uncomplicated skin & skin structure infection, Uncomplicated & complicated UTI.

**Pack :** 1 x 10 Blister



## GLOMYCIN

**Composition :** ERYTHROMYCINE STEARATE TABLETS 250 / 500 MG

**Indication :** Treatment of URTI, LRTI, Ear, Oral, Eye, Skin & Soft tissue, GIT infections, etc.

**Pack :** 1 x 10 Blister and 10 x 10 Blister



## LINCOCLOB

**Composition :** LINCOMYCIN CAPSULES 250 / 500 MG

**Indication :** In treatment of infection from Steptococci, Pneumococci and Staphylococci.

**Pack :** 1 x 10 Blister



## BACIFLOX

**Composition :** OFLOXACIN TABLETS USP 200 / 400 MG

**Indication :** Chronic bronchitis, Community-acquired pneumonia, Acute pelvic inflammatory disease (PID), Acute, Uncomplicated cervical and urethral gonorrhoea, Urethritis, Cervicitis, Uncomplicated cystitis.

**Pack :** 1 x 10 Blister



## SPARFLOXACIN

**Composition :** SPARFLOXACIN TABLETS 100/200/400 MG

**Indication :** Haemophilus influenzae, Haemophilus para-influenzae, Moraxella catarrhalis, Mycoplasma pneumoniae or Streptococcus pneumonia.

**Pack :** 1 x 10 Blister



### PYLOGLOB KIT

**Composition :** OMEPRAZOLE, CLARITHROMYCIN & TINIDAZOLE 20/250/500 MG  
 ESOMEPRAZOLE, CLARITHROMYCIN & TINIDAZOLE 40/250/500 MG  
 LANSOPRAZOLE, CLARITHROMYCIN & TINIDAZOLE 30/250/500 MG

**Indication :** a.1) Helicobacter pylori, Tonsillitis, Pharyngitis, Amoebiasis, Giardiasis, Duodenal ulcer 2. Gastric ulcer 3. Zollinger - Ellison syndrome 4. Gastroesophageal reflux disease b.) Acid reflux , Ulcer, Zollinger- Ellison syndrome 4. Gastroesophageal reflux disease.

**Pack :** Combi Kit



### ROXYGLOB

**Composition :** ROXITHROMYCINE TABLETS 150 / 300MG

**Indication :** It is a semi-synthetic macrolide antibiotic used to treat a variety of bacterial infections. It is used to prevent and treat bacterial infection of the lungs, airways, urinary tract and inner layers of the skin.

**Pack :** 10x10 Blister



### BACINOR

**Composition :** NORFLOXACIN TABLETS 400 MG

**Indication :** Trichomoniasis, bacterial vaginosis, vaginal discharge due to candidiasis, infection or mixed infections with chlamydia and neisseria.

**Pack :** 10x10 and 30x10 Blister



### BACIPRO

**Composition :** CIPROFLOXACIN TABLETS 250 / 500 MG

**Indication :** Exacerbations of chronic obstructive pulmonary disease, broncho-pulmonary infections in cystic fibrosis or in bronchiectasis , chronic suppurative otitis media, urinary tract.

**Pack :** 1x10 and 2x7 Blister



### BACIPRO-TZ

**Composition :** CIPROFLOXACIN + TINIDAZOLE TABLETS 500 + 600 MG

**Indication :** Vaginal or stomach infection, gums infection, skin infections, lungs infection, bacterial infections, treatment of bacterial infections, parasitic infections, infections of the blood, chest infections.

**Pack :** 1x10 Blister



### CLARIGLOB

**Composition :** CLARITHROMYCIN TABLETS 250 / 500 MG

**Indication :** Pharyngitis/Tonsillitis, acute bacterial exacerbation of chronic bronchitis, community-acquired pneumonia, acute maxillary sinusitis, acute otitis media, uncomplicated skin and skin structure infections.

**Pack :** 1x10, 2x7 and 10x10 Blister



### CLINDAMYCIN

**Composition :** CLINDAMYCIN HCL CAPSULES 150/300/600 MG

**Indication :** Serious infections caused by anaerobic bacteria, including intra-abdominal infections, skin and soft tissues infections, tonsillitis, dental infections.

**Pack :** 1x10 Blister



### MOXIGLOB

**Composition :** MOXIFLOXACIN HYDROCHLORIDE TABLETS 400 MG

**Indication :** Acute bacterial sinusitis, Acute exacerbations of chronic bronchitis, Community acquired pneumonia, Mild to moderate pelvic inflammatory disease, Complicated skin and skin structure infections.

**Pack :** 1x5, 1x10 Blister



### AQUAFLUCON

**Composition :** FLUCONAZOLE, AZITHROMYCIN & SECNIDAZOLE TABLETS 250+1000+1000 MG

**Indication :** Trichomoniasis, bacterial vaginosis, vaginal discharge due to candidiasis, infection or mixed infections with chlamydia and neisseria.

**Pack :** Blister Kit



### TINIDAZOLE

**Composition :** TINIDAZOLE TABLETS 500 MG

**Indication :** Vaginal infection (Bacterial vaginosis, Trichomoniasis), Parasite infections (Giardiasis, Amebiasis).

**Pack :** 1x10 Blister



### COTRIM-G

**Composition :** CO-TRIMOXAZOLE TABLETS 480/960 MG

**Indication :** Cotrim-G is used to treat certain bacterial infections, such as pneumonia (a lung infection), bronchitis (infection of the tubes leading to the lungs), and infections of the urinary tract, ears, and intestines.

**Pack :** 10x10 Blister

Coming Soon.....

VORICONAZOLE TABLETS 50 MG AND 200 MG

GEMIFLOXACIN TABLETS 320 MG

LOMEFLOXACIN TABLETS 400 MG

LEVOFLOXACIN AND ORNIDAZOLE TABLETS

### RELIABLE PROTECTION





## ANTI COLD &amp; ANTI HISTAMINE



## CHLORPHENIRAMINE

**Composition :** CHLORPHENIRAMINE TABLETS 2/4/8 MG

**Indication :** Allergy, High fever, Common Cold, Rash, Watery Eyes, Itch Eyes, Nose, Throat, Skin, Cough, Runny nose, Sneezing.

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



## DESLOGLOB

**Composition :** DESLORATADINE TABLETS 5/10 MG

**Indication :** Allergic rhinitis nasal congestion, Acne, Allergic skin conditions (e.g. Chronic idiopathic urticaria)

**Pack :** 3 x 10 and 10x10 Blister



## FEXOglob

**Composition :** FEXOFENADINE HCL TABLETS 120/180 MG

**Indication :** Relief of symptoms associated with seasonal allergic rhinitis, Chronic idiopathic urticaria

**Pack :** 1 x 10 Blister



## GLOCET

**Composition :** CETIRIZINE TABLETS 5/10 MG

**Indication :** Allergic rhinitis such as sneezing, Rhinorrhoea, Itching, Nasal congestion & lacrimation

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



## CETGLO

**Composition :** LEVOCETIRIZINE HYDROCHLORIDE TABLETS 5/10 MG

**Indication :** Relief of nasal and ocular symptoms of seasonal and perennial allergic rhinitis, - The relief of symptoms of chronic idiopathic urticaria.

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



### LORAGLOB

**Composition :** LORATADINE TABLETS 10 MG  
**Indication :** Symptomatic treatment of allergic rhinitis and chronic idiopathic urticaria.  
**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



### MONTELUKAST

**Composition :** MONTELUKAST SODIUM TABLETS 5/10 MG  
**Indication :** To relieve the symptoms of seasonal allergies, Asthma symptoms and seasonal allergic rhinitis or high fever.  
**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



### PERFECT

**Composition :** CETIRIZINE HCL, PARACETAMOL, PHENYLEPHRINE TABLETS/  
 CETIRIZINE HCL, PARACETAMOL, PHENYLEPHRINE ZINC  
 GLUCONATE TABLETS  
**Indication :** Nasal congestion, Allergy, Common cold, Rhinitis, Sinusitis, Fever & CNS stimulant  
**Pack :** 1x10 and 10x10 Blister



### PERFECT D

**Composition :** PARACETAMOL, DEXTROMETHORPHAN HYDROBROMIDE,  
 PHENYLEPHRINE HCL & CPM TABLETS KIT  
**Indication :** Relief of coughs and upper respiratory tract symptoms, including nasal congestion, associated with allergy or common cold.  
**Pack :** 1 x 10 Blister



### ZUKAST

**Composition :** MONTELUKAST SODIUM | LEVOCETIRIZINE HCL 10/5 MG  
 MONTELUKAST SODIUM | LEVOCETIRIZINE HCL 5/2.5 MG  
 MONTELUKAST SODIUM | LEVOCETIRIZINE HCL 4/2.5 MG  
**Indication :** Inflammation of the nose due to an allergic rhinitis, Chronic idiopathic urticaria, Exercise - induced bronchospasm prevention  
**Pack :** 1 x 10, 3 x 10, 10x10 and 20 x 10 Blister



### DELCON FORTE

**Composition :** PHENYLEPHRINE HYDROCHLORIDE, CHLORPHENAMINE MALEATE AND PARACETAMOL TABLETS 10+2+500 MG  
**Indication :** It helps relieve symptoms of cold and flu such as headache, sore throat, runny nose, muscular pain, and fever.  
**Pack :** 1x10, 5x20 Blister



### DIPHENHYDRAMINE

**Composition:** DIPHENHYDRAMINE TABLETS 25/50 MG  
**:Indication :** indicated for the symptomatic relief of allergy symptoms, including rhinitis and urticaria, as well as for the prevention and treatment of motion sickness.  
**Pack :** 3 x 10 and 10 x 10 Blister



### AMBROXOL

**Composition:** AMBROXOL TABLETS 30/60/75/100 MG  
**:Indication :** Ambroxol tablets are indicated for the treatment of acute and chronic respiratory conditions associated with excessive mucus production, such as chronic bronchitis, chronic obstructive pulmonary disease (COPD), and asthma. They work as a mucolytic agent to enhance the clearance of mucus from the respiratory tract by thinning and loosening mucus.  
**Pack :** 3 x 10 and 10 x 10 Blister



### HISTIGLOB

**Composition :** BETAHISTINE HCL TABLETS 8/16/24 MG  
**Indication :** Betahistine is indicated for treatment of meniere's syndrome symptoms of which may include vertigo, tinnitus, hearing loss & nausea. Initial oral treatment is 8 to 16 mg three time daily. Taken preferably with meals.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister





## ANTI DIABETICS



### ADNIL

**Composition :** GLIBENCLAMIDE TABLETS 2.5/5 MG

**Indication :** Glibenclamide Tablet is a fast acting oral anti-diabetic medicine used to control blood sugar levels in Type II Diabetic patients. It belongs to a group of medicines known as sulfonylureas.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### GLICLAZIDE MR

**Composition :** GLICLAZIDE MR TABLETS 30/60 MG

**Indication :** Non Insulin Dependent Diabetes (Type II) in adults when dietary measures, physical exercise and weight loss alone are not sufficient to control blood glucose.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### EMPAGLIFLOZIN+METFORMIN

**Composition :** EMPAGLIFLOZIN+METFORMIN TABLETS 10+500/10+1000/25+500/25+100 MG

**Indication :** Empagliflozin and Metformin tablets are indicated for the management of type 2 diabetes mellitus in adults. This combination therapy aids in glycemic control by reducing blood glucose levels through the enhancement of insulin sensitivity and the reduction of glucose reabsorption in the kidneys.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### GLOPRIDE

**Composition :** GLIMEPIRIDE & EXTENDED RELEASE METFORMIN HCL  
GLIMEPIRIDE & PIOGLITAZONE & METFORMIN 2+15+500 MG

**Indication :** Treatment of Type II Diabetes Mellitus.  
Additive anti-hyperglycaemic effect without increasing the adverse effects of either pharmacological class.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### PIORIDE 30

**Composition :** PIOGLITAZONE HYDROCHLORIDE TABLETS 15/30 MG

**Indication :** Controlling high blood sugar helps prevent kidney damage, blindness, nerve problems, loss of limbs and sexual function problems. Proper control of diabetes may also your risk of a heart attack or stroke.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### STAGLOB M

**Composition :** METFORMIN & SITAGLIPTIN PHOSPHAT 500+50/1000+50 MG

**Indication :** Metformin & Sitagliptin are Oral Diabetes medicines that helps control blood sugar levels. Metformin works by decreasing glucose (sugar) production in the liver and decreasing absorption of glucose by the intestines.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### STAGLOB

**Composition :** SITAGLIPTIN PHOSPHATE TABLETS 25/50/100 MG

**Indication :** Sitagliptin is used with a proper diet and exercise program and possibly with other medications to control high blood sugar. Controlling high blood sugar helps prevent kidney damage, blindness, nerve problems, loss of limbs & sexual function problems.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### GLOFORMIN-ER

**Composition :** METFORMIN HYDROCHLORIDE EXTENDED-RELEASE TABLETS 500/750/1000 MG

**Indication :** Metformin Tablet is an effective first choice medicine used to control high blood glucose levels in patients with Type II Diabetes. Patients are advised to maintain a strict exercise & diet regimen along with this medicine to get the best possible effect.

**Pack :** 1x10, 3X10, 10x10 Blister



### GLOFORMIN

**Composition :** METFORMIN TABLETS 500/850/1000 MG

**Indication :** Metformin Tablet is an effective first choice medicine used to control high blood glucose levels in patients with Type II Diabetes. Patients are advised to maintain a strict exercise & diet regimen along with this medicine to get the best possible effect.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### REPAGLO

**Composition :** REPAGLINIDE TABLETS 0.5/1/2 MG

**Indication :** Repaglinide tablets are used alone or with other medications to control high blood sugar along with a proper diet and exercise program. Controlling high blood sugar helps prevent kidney damage, blindness, nerve problems, loss of limbs and sexual function problems.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### SAXAGLOB

**Composition :** SAXAGLIPTIN TABLETS 2.5/5 MG

**Indication :** Saxagliptin is used along with diet and exercise to lower blood sugar levels in patients with type 2 diabetes. Saxagliptin is in a class of medications called dipeptidyl peptidase-4 (DPP-4) inhibitors.

**Pack :** 2x14 Blister, 1x10 Alu Alu



### EMPAGLOB

**Composition :** EMPAGLIFLOZIN TABLETS 10/25 MG

**Indication :** Empagliflozin is a type of medicine known as a sodium-glucose co-transporter-2 (SGLT2) inhibitor. It reduces blood sugar levels in people with diabetes.

**Pack :** 10x10 Blister



### CANAGLOB

**Composition :** CANAGLIFLOZIN TABLETS 100/300 MG

**Indication :** Canagliflozin is indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus.

**Pack :** 3x10 Blister



### DAPAGLIFLOZIN + METFORMIN

**Composition :** DAPAGLIFLOZIN + METFORMIN TABLETS 5+500/5+1000/10+500/10+1000 MG

**Indication :** Dapagliflozin and Metformin tablets are indicated for the treatment of type 2 diabetes mellitus in adults. This combination therapy works to improve glycemic control by utilizing dapagliflozin's mechanism as an SGLT2 inhibitor to reduce renal glucose reabsorption and metformin's action to enhance insulin sensitivity and decrease hepatic glucose production.

**Pack :** 3 x 10 and 10 x 10 Blister



### VILDAGLIPTIN + METFORMIN

**Composition :** VILDAGLIPTIN+METFORMIN TABLETS 50+500/50+1000/100+500/100+1000/100+2000 mg

**Indication :** Vildagliptin and Metformin tablets are indicated for the management of type 2 diabetes mellitus in adults. This combination therapy enhances glycemic control by increasing insulin secretion (through vildagliptin, a DPP-4 inhibitor) and improving insulin sensitivity and reducing hepatic glucose production (through metformin).

**Pack :** 3 x 10 and 10 x 10 Blister



### DAPAGLOB

**Composition :** DAPAGLIFLOZIN TABLETS 5/10 MG

**Indication :** Dapagliflozin is an antidiabetic medication. It works by removing excess sugar from your body through urine.

**Pack :** 1x14 Blister



### GLOPRIDE

**Composition :** GLIMEPIRIDE TABLETS 1/2/4 MG

**Indication :** Treatment of Type II Diabetes Mellitus.

**Pack :** 1 x10, 3 x 10 and 10 x 10 Blister



### VILDAGLIPTIN

**Composition :** VILDAGLIPTIN TABLETS 50/100 MG

**Indication :** Vildagliptin tablets are indicated for the treatment of type 2 diabetes mellitus in adults. They work as a DPP-4 (dipeptidyl peptidase-4) inhibitor, enhancing glycemic control by increasing insulin secretion and decreasing glucagon levels in response to elevated blood glucose levels.

**Pack :** 3 x 10 and 10 x 10 Blister



### GLIBENCLAMIDE + METFORMIN

**Composition :** GLIBENCLAMIDE + METFORMIN TABLETS  
2.5+500/2.5+1000/5+500/5+1000

**Indication :** Vildagliptin tablets are indicated for the treatment of type 2 diabetes mellitus in adults. They work as a DPP-4 (dipeptidyl peptidase-4) inhibitor, enhancing glycemic control by increasing insulin secretion and decreasing glucagon levels in response to elevated blood glucose levels.

**Pack :** 3 x 10 and 10 x 10 Blister



## ANTI MALARIAL



## CHLOROQUINE PHOSPHATE

**Composition :** CHLOROQUINE PHOSPHATE TABLETS 100/150/250/500 MG

**Indication :** Chloroquine Phosphate Tablets are indicated for suppressive treatment and for acute attacks of Malaria due to P. Vivax, P. Malariae, P. Ovale and Susceptible strains of P. Falciparum.

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



## ARTESUNATE &amp; MEFLOQUINE

**Composition :** ARTESUNATE + MEFLOQUINE TABLETS 300+375 MG

**Indication :** Artesunate & Mefloquine is a medication used to treat Malaria. The combination is specifically recommended to treat uncomplicated P. Falciparum Malaria.

**Pack :** 1 x 6 Blister



## GLOQ-300

**Composition :** QUININE SUPHATE TABLETS 150/300/600 MG

**Indication :** The treatment of Plasmodium Falciparum Malaria. Treatment and prevention of nocturnal leg cramps in adults and the elderly, when cramps cause regular disruption of sleep.

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



## GLOSUNAT PLUS 200 KIT

**Composition :** COMBI KIT OF ARTESUNATE + PYRIMETHAMINE & SULPHADOXINE TABLETS 25+500+200 MG

**Indication :** As first line treatment of confirmed P. Falciparum Malaria infection as first line or after failure of Chloroquine Therapy.

**Pack :** 1 and 10 Blister Kit



## GLOSUNATE PLUS 50

**Composition :** ARTESUNATE + AMODIAQUINE HCL 50+153.1 MG

**Indication :** As first line treatment of confirmed P. Falciparum Malaria infection as first line or after failure of Chloroquine Therapy.

**Pack :** 24 Blister Kit



## LUMAGLOB

**Composition :** ARTEMETHER + LUMEFANTRINE 80+480 MG

ARTEMETHER + LUMEFANTRINE 40+240 MG

ARTEMETHER + LUMEFANTRINE 20+120 MG

**Indication :** Treatment of acute uncomplicated Plasmodium Falciparum Malaria.

**Pack :** 1 x 3, 1 x 6, 4 x 6, 1 x 12 and 1 x 24 Blister



## LUMAGLOB SUSPENSION

**Composition :** ARTEMETHER + LUMEFANTRINE SUSPENSION 180+1080 MG  
**Indication :** Treatment of Malaria in children.  
**Pack :** 60 ML  
**Flavor :** Orange, Banana



## PIPART

**Composition :** DIHYDROARTEMISININ + PIPERAQUINE PHOSPHATE SUSPENSION 80+640 MG  
**Indication :** Treatment of Malaria in children.  
**Pack :** 80 ML  
**Flavor :** Banana



## PYMAL

**Composition :** DIHYDROARTEMISININ TABLETS + PIPERAQUINE PHOSPHATE 40+320/120+960 MG  
**Indication :** Treatment of acute uncomplicated Plasmodium Falciparum Malaria.  
**Pack :** 1 x 3 and 1 x 9 Blister



## ARTEMISININ + PIPERAQUINE

**Composition :** ARTEMISININ + PIPERAQUINE TABLETS 62.5 + 375 MG  
**Indication :** This combination therapy provides rapid parasite clearance and extended protection against malaria through the synergistic effects of artemisinin and piperaquine.  
**Pack :** 1 x 4 Blister



## ANTI PSYCHOTIC &amp; CNS



## LAMOTRIGINE

**Composition :** LAMOTRIGINE TABLETS 25/50/100/200 MG

**Indication :** Lamotrigine tablets are indicated for the management of epilepsy and bipolar disorder. In epilepsy, it is used as an adjunctive treatment for partial-onset seizures, For bipolar disorder, it is used as mood episodes, depressive episodes.

**Pack :** 3 x 10 and 10 x 10 Blister



## CITICOLINE

**Composition :** CITICOLINE TABLETS 500 MG

**Indication :** Citicoline on it's own can serve to enhance several different mental processes related to memory, learning, logical reasoning and focus.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## ESCITALOPRAM

**Composition :** ESCITALOPRAM TABLETS USP 5/10/15/20 MG

**Indication :** Treatment of - major depressive episodes, panic disorder with or without agoraphobia, social anxiety disorder (social phobia), generalised anxiety disorder, obsessive-compulsive disorder.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## GLOPENTIN

**Composition :** GABAPENTIN TABLETS / CAPSULES 100/300/600/800 MG

**Indication :** Gabapentin is used with other medications to prevent and control seizures. It is also used to relieve nerve pain following shingles (a painful rash due to herpes zoster infection) in adults.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## LEVETIRACETAM

**Composition :** LEVETIRACETAM TABLETS 250/500/1000 MG

**Indication :** It is used in combination with other seizure-control medications to treat and prevent seizures associated with epilepsy.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## PAROXETINE

**Composition :** PAROXETINE TABLETS 20/40 MG

**Indication :** Paroxetine tablets are indicated for the treatment of major depressive disorder.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## QUETIAPINE FUMARATE

**Composition :** QUETIAPINE FUMARATE TABLETS 25/50/100/200/300/400 MG

**Indication :** This medication is used to treat certain mental/mood conditions (such as schizophrenia, bipolar disorder, sudden episodes of mania or depression associated with bipolar disorder).

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## CITALOPRAM

**Composition :** CITALOPRAM TABLETS 10/20 MG

**Indication :** Citalopram is known as a selective serotonin reuptake inhibitor (SSRI). Citalopram is used to treat depression. It may improve your energy level and feelings of well-being.

**Pack :** 10x10 Blister




## ARIPIRAZOLE

**Composition :** ARIPIRAZOLE TABLETS 2/5/10/15/20/30 MG

**Indication :** Aripiprazole is used to treat certain mental/mood disorders (such as bipolar disorder, schizophrenia, Tourette's syndrome, and irritability associated with autistic disorder).

**Pack :** 10x10 Blister



## BRIVARACETAM

**Composition :** BRIVARACETAM TABLETS 25/50 MG

**Indication :** Brivaracetam is used alone and in combination with other medications to control partial onset seizures in adults, children, and infants 1 month of age and older.

**Pack :** 10x10 Blister



## VIGAGLOB 500

**Composition :** VIGABATRIN TABLETS USP 500 MG

**Indication :** Vigabatrin is indicated as adjunctive therapy in the treatment of refractory complex partial seizures in patients 2 years of age and older who have had inadequate responses to multiple previous treatments.

**Pack :** 10x10 Blister



## OXCARBAZEPINE

**Composition :** OXCARBAZEPINE TABLETS 150/300/600 MG

OXCARBAZEPINE EXTENDED RELEASE TABLETS 300/450 MG

**Indication :** Oxcarbazepine tablets are indicated for the treatment of partial-onset seizures, with or without secondary generalization, in adults and children with epilepsy. They can be used as monotherapy or adjunctive

**Pack :** therapy to achieve better seizure control.  
1 x 10, 3 x 10 and 10 x 10 Blister



## ANTI PSYCHOTIC &amp; CNS



## CARBAMAZEPINE

**Composition :** CARBAMAZEPINE 100/200/400 MG

**Indication :** Carbamazepine Tablets are indicated for the treatment of epilepsy and neuropathic pain. It is used as monotherapy or adjunctive therapy for partial seizures, generalized tonic-clonic seizures, and mixed seizure types.

**Pack :** 3 x 10 and 10 x 10 Blister



## QUETIAPINE XR

**Composition :** QUETIAPINE XR TABLETS 50/100/200/300/400 MG

**Indication :** Quetiapine XR Tablets are indicated for the treatment of schizophrenia and bipolar disorder in adults and adolescents. It is used to manage acute manic, mixed, and depressive episodes in bipolar disorder, as well as to maintain stability and prevent relapse. Quetiapine XR is also prescribed for the adjunctive treatment of major depressive disorder in combination with other antidepressants.

**Pack :** 3 x 10 and 10 x 10 Blister



## PREGABALIN

**Composition :** Pregabalin Capsules 25/50/75/100/150/200/300 MG

**Indication :** Pregabalin Capsules are indicated for the management of neuropathic pain associated with diabetic peripheral neuropathy, postherpetic neuralgia, and spinal cord injury. Additionally, they are used as adjunctive therapy for partial-onset seizures and for the treatment of generalized anxiety disorder.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## ANTIFUNGAL &amp; ANTHELMINTICS



## ALBAGLOB

**Composition :** ALBENDAZOLE TABLETS 400 MG

**Indication :** Treatment of Parenchymal Neurocysticercosis, Parasitic Infections.  
Treatment of Cystic Hydatid Disease of Liver, Lung & Peritoneum.

**Pack :** 1 x 1, 1 x 10, 3 x 10 and 10 x 10 Blister



## CLOTRIMAZOLE

**Composition :** CLOTRIMAZOLE VAGINAL TABLETS 100/400 MG

**Indication :** Clotrimazole is an antifungal medication that fights infections caused by fungus. Clotrimazole vaginal (for use in the vagina) is used to treat vaginal candida (yeast) infections.

**Pack :** 1 x 6, 3 x 10 and 10 x 10 Blister And Strip



## CYSTOGLY-N

**Composition :** DILOXANIDE FUROATE TABLETS 500 MG

**Indication :** Diloxanide is a medication used to treat amoeba infections. It is a second line treatment after paromomycin when no symptoms are present in places where infections are not common. For people who are symptomatic.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## FLUCAT

**Composition :** FLUCONAZOLE TABLETS/CAPSULES 50/150/200/400 MG

**Indication :** Coccidioidomycosis, Invasive Candidiasis, Mucosal Candidiasis including Oropharyngeal & Oesophageal candidiasis, Candiduria & Chronic Mucocutaneous Candidiasis.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## GLOMET

**Composition :** METRONIDAZOLE TABLETS 200/250/400/500 MG

**Indication :** Prophylaxis and treatment of identified / suspected anaerobic bacterial infections such as Bacteroides, Fusobacteria, Clostridia, Eubacteria, Anaerobic Cocci and Gardnerella Vaginalis, Trichomonas.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## GRISOMED

**Composition :** GRISEOFULVIN TABLETS 250/500 MG

**Indication :** Griseofulvin is an effective antifungal medicine used to treat infection of the skin, scalp, nails, feet, groin & other parts of the body.

**Pack :** 1 x 10, 3 x 10, 1 x 20 and 10x10 Blister



### ITRACONAZOLE

**Composition :** ITRACONAZOLE CAPSULES 100/200 MG

**Indication :** Itraconazole is used to treat a variety of fungal infections. It belongs to a class of drugs known as azole antifungals. It works by stopping the growth of fungi.

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



### KETAGLO

**Composition :** KETOCONAZOLE TABLETS 200 MG

**Indication :** Treatment of systemic fungal infections in patients with failure or intolerance to other therapies: Blastomycosis, Coccidioidomycosis, Histoplasmosis, Chromomycosis & Paracoccidioidomycosis.

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



### NITAZOXANIDE

**Composition :** NITAZOXANIDE TABLETS 500 MG

**Indication :** This medication is used to treat diarrhoea caused by certain parasite infections of the intestines (Cryptosporidium parvum and Giardia Lamblia).

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



### BACTROSEC

**Composition :** SECNIDAZOLE TABLETS 500 MG

**Indication :** Secnidazole is an effective medicine to treat infections caused by protozoa and certain strains of anaerobic bacteria. It is used to treat infections of the stomach, intestine, urinary tract and genital area.

**Pack :** 1 x 2, 1 x 4, 1 x 10, 3 x 10 and 10 x 10 Blister



### MEBAMED

**Composition :** MEBENDAZOLE TABLETS 100/200/500 MG

**Indication :** Mebendazole is a type of medicine for treating worms. It is used mainly for infections of the gut such as threadworms and other less common worm infections.

**Pack :** 1 x 6 Blister



### FLUCAT SUSPENSION

**Composition :** FLUCONAZOLE SUSPENSION

**Indication :** Coccidioidomycosis, Invasive Candidiasis, Mucosal Candidiasis including Oropharyngeal & Oesophageal candidiasis, Candiduria & Chronic Mucocutaneous Candidiasis.

**Pack :** 60 ML





## ARV PRODUCTS



## ABACAVIR SULFATE &amp; LAMIVUDINE

**Composition :** ABACAVIR 300/600 MG + LAMIVUDINE 150/300 MG TABLETS

**Indication :** Abacavir/Lamivudine is indicated in antiretroviral combination therapy for the treatment of Human Immunodeficiency Virus (HIV) infection in adults, adolescents and children weighing at least 25 kg before initiating treatment with abacavir.

**Pack :** 30/60 Tablets Bottle



## ATAZANAVIR &amp; RITONAVIR

**Composition :** ATAZANAVIR & RITONAVIR TABLETS 300+100 MG

**Indication :** Atazanavir/Ritonavir is indicated in antiretroviral combination therapy for the treatment of Human Immunodeficiency Virus (HIV-1) infections.

**Pack :** 30 Tablets Bottle



## DARUNAVIR

**Composition :** DARUNAVIR (DRV) 300/400/600/800 MG TABLETS

**Indication :** Darunavir co-administrated with low dose ritonavir is indicated in combination with other antiretroviral medicinal products for the treatment of patients with human immunodeficiency virus (HIV-1) infections.

**Pack :** 30/60 Tablets Bottle



## EFAVIRENZ

**Composition :** EFAVIRENZ CAPSULES 200/600 MG

**Indication :** This drug is used with other HIV medications to help control HIV infection. It helps to increase the amount of HIV in your body so your immune system can work better. This lowers your chance of getting HIV complications (such as new infections, cancer) & improves your quality of life.

**Pack :** 30/90 Capsules Bottle



## EFAVIRENZ, LAMIVUDINE &amp; TENOFOVIR DISOPROXIL FUMARATE

**Composition :** EFAVIRENZ & LAMIVUDINE & TENOFOVIR 600+300+300 MG

**Indication :** This combination product is used by itself to help control HIV infection. It helps to decrease the amount of HIV in your body so your immune system can work better.

**Pack :** 30/60 Tablets Bottle



## EMTRICITABINE & TENOFOVIR

**Composition :** TENOFOVIR & EMTRICITABINE TABLETS 300+200 MG

**Indication :** This product is used with other HIV medications to help control HIV infection. It helps to decrease the amount of HIV in your body so your immune system can work better. This lowers your chance of getting HIV complications (such as new infections, cancer) & improves your quality of life.

**Pack :** 30/60 Tablets Bottle



## RALTEGLOB

**Composition :** RALTEGRAVIR TABLETS USP 400/600 MG

**Indication :** Raltegravir is indicated in combination with other antiretroviral medicinal products for the treatment of human immunodeficiency virus (HIV-1) infection.

**Pack :** 30/60 Tablets Bottle



## RITONAVIR

**Composition :** RITONAVIR TABLETS 100 MG

**Indication :** Ritonavir is indicated in combination with other antiretroviral agents for the treatment of HIV-1 infected patients (adults and children of 2 years of age and older).

**Pack :** 30/60 Tablets Bottle



## SOFOSBUVIR

**Composition :** SOFOSBUVIR TABLETS 150/200/300/400 MG

**Indication :** Sofosbuvir is indicated for the treatment of chronic Hepatitis C virus (HCV) infection as a component of a combination antiviral regimen for patients with HCV mono-infection and HCV/HIV-1 coinfection.

**Pack :** 28 Tablets Bottle



## TENOFOVIR & LAMIVUDINE

**Composition :** TENOFOVIR + LAMIVUDINE TABLETS 300+300 MG

**Indication :** Tenofovir/Lamivudine is indicated in anti-retroviral combination therapy for the treatment of human immunodeficiency virus (HIV) infection.

**Pack :** 30/60 Tablets Bottle



## TENOFOVIR

**Composition :** TENOFOVIR TABLETS 300 MG

**Indication :** Tenofovir Disoproxil 245 mg film-coated tablets are indicated in combination with other anti-retroviral medicinal products for the treatment of HIV-1 infected adults.

**Pack :** 30/60 Tablets Bottle



## LOPINAVIR & RITONAVIR

**Composition :** LOPINAVIR & RITONAVIR TABLETS 200+50/400+100 MG

**Indication :** Lopinavir/Ritonavir tablets are indicated for the treatment of infected adults and children weighing 25 kg or more who are infected with human immunodeficiency virus type 1 (HIV-1).

**Pack :** 120 Tablets Bottle



## NEVIRAPINE

**Composition :** NEVIRAPINE TABLETS USP 200 MG

**Indication :** Nevirapine is indicated in combination with other antiretroviral medicinal products for the treatment of HIV-1 infected adults, adolescents, and children of any age. Most of the experience with Nevirapine is in combination with nucleoside reverse transcriptase inhibitors (NRTIs).

**Pack :** 60 Tablets Bottle



## LAMIVUDINE & ZIDOVUDINE

**Composition :** LAMIVUDINE & ZIDOVUDINE TABLETS 150 + 300 MG / 30 + 60 MG

**Indication :** Lamivudine & Zidovudine Tablet is used with other anti retroviral medicines to treat HIV infection in adults and children. It reduces the amount of HIV in your body & keeps it at a low level. It also increases CD4 cell count in your blood.

**Pack :** 60 Tablets Bottle



## LAMIVUDINE, ZIDOVUDINE & NEVIRAPINE

**Composition :** LAMIVUDINE, ZIDOVUDINE & NEVIRAPINE TABLETS

**Indication :** 150+200+300MG

The combination of lamivudine, Zidovudine & Nevirapine is used along with other medications to treat human immunodeficiency virus (HIV) infection. Lamivudine and zidovudine are in a class of medications called nucleoside reverse transcriptase inhibitors (NRTIs). They work by decreasing the amount of HIV in the blood.

**Pack :** 60 Tablets Bottle



## ATAZANAVIR

**Composition :** ATAZANAVIR CAPSULES 100/150 MG

**Indication :** Atazanavir is a prescription medicine approved by the U.S. Food and Drug Administration (FDA) for the treatment of HIV infection in adults and children 3 months of age and older who weigh at least 11 lb (5 kg). Atazanavir is always used in combination with other HIV medicines.

**Pack :** 60 Tablets Bottle



## ABACAVIR

**Composition :** ABACAVIR CAPSULES 300 MG

**Indication :** Abacavir is indicated in antiretroviral combination therapy for the treatment of Human Immunodeficiency Virus (HIV) infection in adults, adolescents and children.

**Pack :** 60 Capsules Bottle



## EMTRICITABINE & TENOFOVIR ALAFENNAMIDE

**Composition :** EMTRICITABINE & TENOFOVIR ALAFENNAMIDE TABLETS 200,10 MG / 200,25 MG

**Indication :** Combination of Emtricitabine (FTC) and Tenofovir alafenamide (TAF), both of them are HIV nucleosides analog reverse transcriptase inhibitors (NRTIs), and is indicated for the HIV-1 infection

**Pack :** 30 Tablets Bottle



## RIBAVIRIN

**Composition :** RIBAVIRIN CAPSULES 200 MG

**Indication :** Ribavirin is a nucleoside analogue and antiviral agent used to treat RSV infection, hepatitis C and some viral hemorrhagic fevers. For hepatitis C, it is used in combination with other medications such as simeprevir, sofosbuvir, and peg interferon alfa-2b or peg interferon alfa-2a.

**Pack :** 40 Capsules Bottle



## DOLUTEGRAVIR

**Composition :** DOLUTEGRAVIR TABLETS 50 MG

**Indication :** Dolutegravir is a human immunodeficiency virus type 1 (HIV-1) integrase strand transfer inhibitor (INSTI) indicated in combination with other antiretroviral agents for the treatment of HIV-1 infection in adults and children aged 12 years and older and weighing at least 40 kg.

**Pack :** 30 Tablets Bottle

**DOLUTEGRAVIR & LAMIVUDINE****Composition :** DOLUTEGRAVIR & LAMIVUDINE TABLETS 50/300 MG**Indication :** Combination of dolutegravir (integrase strand transfer inhibitor [INSTI]) and lamivudine (nucleoside analogue reverse transcriptase inhibitor [NRTI]) is indicated as a complete regimen for the treatment of human immunodeficiency virus type 1 (HIV-1) infection**Pack :** 30 Tablets Bottle**ABACAVIR, DOLUTEGRAVIR & LAMIVUDINE****Composition :** ABACAVIR, DOLUTEGRAVIR & LAMIVUDINE TABLETS 600+50+300 MG**Indication :** Combination of dolutegravir (integrase strand transfer inhibitor [INSTI]), abacavir, and lamivudine (both nucleoside analogue reverse transcriptase inhibitors) is indicated for the treatment of HIV-1 infection in adults and in pediatric patients weighing at least 40 kg.**Pack :** 30 Tablets Bottle**FAMCICLOVIR****Composition :** FAMCICLOVIR TABLETS 250 MG**Indication :**

- For the treatment of herpes zoster and ophthalmic zoster in immunocompetent adults
- recurrent outbreaks of genital herpes caused by HSV and to prevent outbreaks from reoccurring (called chronic suppressive therapy)
- shingles (also known as herpes zoster) caused by VZV

**Pack :** 1 x 6 Blister**GLOVUDINE****Composition :** LAMIVUDINE TABLETS 100/150/300 MG**Indication :** Compensated liver disease with evidence of active viral replication, persistently elevated serum alanine aminotransferase (ALT) levels and histological evidence of active liver inflammation and/or fibrosis. Initiation of lamivudine treatment should only be considered when the use of an alternative antiviral agent with a higher genetic barrier is not available or appropriate**Pack :** 1 x 10 Blister**VCYCLOR****Composition :** VALACYCLOVIR TABLETS 500/1000 MG**Indication :** Valacyclovir is used to treat infections caused by certain types of viruses. In children, it is used to treat cold sores around the mouth (caused by herpes simplex) and chickenpox (caused by varicella zoster).**Pack :** 10 x 10 Blister



### ZIDOVUDINE

**Composition :** ZIDOVUDINE 100/300 MG Capsules

**Indication :** Zidovudine is indicated in anti-retroviral combination therapy for Human Immunodeficiency Virus (HIV) infected adults and children. Zidovudine chemoprophylaxis is indicated for use in HIV-positive pregnant women (over 14 weeks of gestation) for prevention of maternal-foetal HIV transmission and for primary prophylaxis of HIV infection in newborn infants.

**Pack :** 10x10 Blister



### FAVIPRAVIR

**Composition :** FAVIPRAVIR TABLETS 200/400 MG

**Indication :** Favipiravir is an antiviral medication used to treat influenza. It is also being studied to treat a number of other viral infections. Like the experimental antiviral drugs (T-1105 and T-1106), it is a pyrazinecarboxamide derivative.

**Pack :** 1x10 Blister



### OSELTAMIVIR

**Composition :** OSELTAMIVIR PHOSPHATE CAPSULES USP 75 MG

**Indication :** Osetamivir is in a class of medications called neuraminidase inhibitors. It works by stopping the spread of the flu virus in the body.

**Pack :** 1x10 Blister



### GLOCYTE

**Composition :** VALGANCICLOVIR TABLETS USP 450 MG

**Indication :** Valganciclovir is indicated for the induction and maintenance treatment of cytomegalovirus (CMV) retinitis in adult patients with acquired immunodeficiency syndrome (AIDS).

**Pack :** 60 tablets in HDPE Bottle



### TENOFOVIR + LAMIVUDINE + DOLUTEGRAVIR

**Composition :** TENOFOVIR + LAMIVUDINE + DOLUTEGRAVIR 300+300+50 MG

**Indication :** Valganciclovir is indicated for the induction and maintenance treatment of cytomegalovirus (CMV) retinitis in adult patients with acquired immunodeficiency syndrome (AIDS).

**Pack :** 60 tablets in HDPE Bottle



## CARDIOVASCULAR DRUGS



## ATORVASTATIN

**Composition :** ATORVASTATIN TABLETS 10/20/40 MG

**Indication :** Atorvastatin is used along with a proper diet to help lower “bad” cholesterol and fats (such as LDL, triglycerides) and raise “good” cholesterol (HDL) in the blood. It belongs to a group of drugs known as “statins”. It works by reducing the amount of cholesterol made by the liver.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## ATORVASTATIN &amp; FENOFIBRATE

**Composition :** ATORVASTATIN & FENOFIBRATE TABLETS 10+160 MG

**Indication :** Atorvastatin uses include treatment of dyslipidemia and the prevention of cardiovascular disease. Atorvastatin reduces level of cholesterol, HDL and triglycerides which are responsible for cardiovascular disorders, Fenofibrate a lipid-lowering agent, prescribed for hypercholesterolemia.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## ATORVASTATIN &amp; EZETIMIBE

**Composition :** ATORVASTATIN & EZETIMIBE 20+10/40+10/80+10 MG

**Indication :** Atorvastatin is used along with a proper diet to help lower “bad” cholesterol and fats (such as LDL, triglycerides) and raise “good” cholesterol (HDL) in the blood. It belongs to a group of drug known as “statins”. It works by reducing the amount of cholesterol made by the liver.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## BISOPROLOL FUMARATE

**Composition :** BISOPROLOL FUMARATE TABLETS 2.5/ 5 / 10 MG

**Indication :** Treatment of hypertension, chronic stable angina pectoris, stable chronic heart failure with reduced systolic ventricular function in addition to ACE inhibitors, diuretics and optionally cardiac glycosides.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## BOSENTAN

**Composition :** BOSENTAN TABLETS 62.5/ 125 MG

**Indication :** Bosentan belongs to a class of medication called endothelin receptor antagonists hypertension.

**Pack :** 1 x 10, 2X10, 3 x 10 and 10 x 10 Blister



## CAPTOGLO

**Composition :** CAPTOPRIL TABLETS 25/50 MG

**Indication :** Captopril is an ACE inhibitor. ACE stands for Angiotensin Converting Enzyme. Captopril is used to treat high blood pressure (hypertension), congestive heart failure, kidney problems caused by diabetes, and improve survival after heart attack.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### CLOPIDOGREL + ASPIRIN

**Composition :** CLOPIDOGREL 75 + ASPIRIN TABLETS 75 MG  
**Indication :** Clopidogrel and Aspirin are anti-platelet agent that is a drug that inhibits the ability of platelet to clump together as a part of a blood clot.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### DIPIGLO

**Composition :** AMLODIPINE BESYLATE TABLETS 5/10 MG  
**Indication :** Hypertension, Chronic Stable Angina Pectoris  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### DIPIGLO-L

**Composition :** AMLODIPINE & LISINOPRIL 5/10 MG & 10/10 MG  
**Indication :** Amlodipine & Lisinopril works well to lower BP and prevent chest pain. You can't miss does and you might get persistent swelling. Lisinopril is a good blood pressure-lowering medicine that protects kidney function.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### DIPIGLO-V & DIPIGLO-VH

**Composition :** AMLODIPINE & VALSARTAN TABLETS 5/160 MG & 5/320 MG  
 AMLODIPINE, VALSARTAN & HCT TABLETS 10/160/12.5 MG  
**Indication :** This product is used to treat high blood pressure. Lowering high blood pressure helps strokes, heart attacks and kidney problems.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### ATENOLOL

**Composition :** ATENOLOL TABLETS 25/50/100 MG  
**Indication :** Atenolol Tablets are indicated for the management of cardiovascular conditions Hypertension, Angina Pectoris, Acute Myocardial Infarction, Chronic Stable Angina.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### ENAGLOB

**Composition :** ENALAPRIL TABLETS 5/10/20 MG  
**Indication :** Enalapril tablet is used in the treatment of high blood pressure and certain other heart conditions. It is important to maintain a proper diet and exercise regularly when taking this medicine. Do not take this medicine if you have a know allergy to it.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### EPLERENONE

**Composition :** EPLERENONE TABLETS 25/ 50 MG

**Indication :** This medicine is used alone or in combination with other medicines to treat high blood pressure. It works by blocking a chemical (Aldosterone) in your body which in turn lowers the amount of sodium and water the body retains.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### FENOFIBRATE

**Composition :** FENOFIBRATE TABLETS 120/160/200 MG

**Indication :** Fenofibrate is used along with a proper diet to help lower “bad” cholesterol and fats (such as LDL, Triglycerides) and raise “good” cholesterol (HDL) in the blood.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### GLOSARTAN

**Composition :** TELMISARTAN TABLETS 20/40/80/160 MG

**Indication :** Treatment of essential hypertension in adults, cardiovascular prevention (patients with H/O CHD, Stroke/PAD, Type II DM with target organ damage).

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### GLOSARTAN H

**Composition :** TELMISARTAN + HCT TABLETS 40/12.5 MG & 80/12.5

**Indication :** Treatment of essential hypertension.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### IRBESARTAN

**Composition :** IRBESARTAN TABLETS 150/300 MG

**Indication :** Irbesartan is used to treat high blood pressure (Hypertension) and to help protect the kidneys from damage due to Diabetes. Lowering high blood pressure helps prevent strokes, heart attacks and kidney problems

**Pack :** 1 x 10, 2 x 7, 3 x 10 and 10 x 10 Blister



### IRBESARTAN & HCT

**Composition :** IRBESARTAN & HYDROCHLORTHIAZIDE 50+12.5/150+12.5/300+25 MG

**Indication :** Irbesartan Hydrochlorothiazide tablets are indicated for the treatment of hypertension. Patients whose blood pressure is not adequately controlled on monotherapy.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## IVABRADINE

**Composition :** IVABRADINE TABLETS 5/7.5 MG

**Indication :** Ivabradine tablets are indicated to reduce the risk of hospitalization for worsening heart failure in patients with stable, symptomatic chronic heart failure with left ventricular ejection fraction with left ventricular ejection fraction  $\leq 35\%$ .

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## LISINOPRIL

**Composition :** LISINOPRIL TABLETS 5/10 MG

**Indication :** Lisinopril is used to treat high blood pressure. Lowering high blood pressure helps prevent strokes, heart attacks and kidney problems. It is also used to treat heart failure and to improve survival after a heart attack.

**Pack :** 2 x 14, 1 x 10, 3 x 10 and 10 x 10 Blister



## LOSARGLOB

**Composition :** LOSARTAN POTASSIUM TABLETS 25/50 MG

**Indication :** Losartan is used to treat high blood pressure (hypertension) and to help protect the kidneys from damage due to Diabetes. It is also used to lower the risk strokes in patients with high blood pressure and an enlarged heart. Lowering high blood pressure strokes, heart kidney problems.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## LOSARTAN HCT

**Composition :** LOSARTAN POTASSIUM + HCT TABLETS 50+12.5/100+12.5 MG

**Indication :** Losartan + HCT Tablets are indicated to reduce blood pressure in patients with essential hypertension and improve cardiovascular outcomes by managing hypertension, which may reduce the risk of stroke, myocardial infarction, and renal complications.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## METHYLDOPA

**Composition :** METHYLDOPA TABLETS 250/500 MG

**Indication :** Methyldopa is used to treat high blood pressure. Lowering high blood pressure helps prevent strokes, heart attacks and kidney problems. It is also used to treat heart failure and to improve survival after a heart attack.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## NEBIVOLOL

**Composition :** NEBIVOLOL TABLETS 2.5/5 MG

**Indication :** Nebivolol is used to treat high blood pressure. Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## GLODIPINE

**Composition :** NIFEDIPINE EXTENDED RELEASE 5/10/20 /30/60/90 MG

**Indication :** This medication is used to prevent certain types of chest pain (Angina). It may allow you to exercise more and decrease the frequency of Angina attacks. Nifedipine belongs to a class of medications known as Calcium Channel Blocker.

**Pack :** 3 x 10 and 10 x 10 Blister



## OLMESARTAN MEDOXOMIL

**Composition :** OLMESARTAN MEDOXOMIL TABLETS 20/40 MG

**Indication :** Olmesartan is used to treat high blood pressure (Hypertension). Lowering high blood pressure helps prevent strokes, heart attacks and kidney problems. Olmesartan belongs to a class of drugs called Angiotensin Receptor Blocker (ARB).

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## PERINDOPRIL ARGININE

**Composition :** PERINDOPRIL ARGININE TABLETS 2.5/5/10 MG

**Indication :** Perindopril shares the indications of ACE inhibitors as a class, including essential hypertension, stable coronary artery disease (reduction of risk of cardiac events in patients with a history of myocardial infarction and / or revascularization) and treatment of heart disease or heart failure.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## PLAVIDO

**Composition :** CLOPIDOGREL TABLETS 75 MG

**Indication :** Clopidogrel is used to reduce the rate of myocardial infarction and strokes.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## RAMIGLOB

**Composition :** RAMIPRIL TABLETS 2.5/5/10 MG

**Indication :** Ramipril 5 mg tablet is an Angiotensin Converting Enzyme (ACE) inhibitor. Ramipril 5 mg tablet is used for the treatment of Hypertension or high blood pressure.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## ROSUGLOB

**Composition :** ROSUVASTATIN TABLETS 5/10/20/40 MG

**Indication :** Treatment of hypercholesterolaemia, prevention of cardiovascular events.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## ROSUGLOB-EZ

- Composition :** ROSUVASTIN & EZETIMIBE TABLETS 20/10 MG  
**Indication :** Rosuvastatin is used to reduce LDL, Triglycerides - "bad" cholesterol and fats and raise HDL - "good" cholesterol in the blood.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## SIMVASYN

- Composition :** SIMVASTATIN 5/10/20 MG  
**Indication :** Simvastatin is used along with a proper diet to help lower "bad" cholesterol and fats (such as LDL, Triglycerides) and raise "good" cholesterol (HDL) in the blood. Simvastatin are used to lower bad cholesterol and raise good cholesterol (HDL).  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## ROSUGLOB-F

- Composition :** ROSUVASTATIN & FENOFIBRATE TABLETS 10+160 MG  
**Indication :** Rosuvastatin is used to reduce LDL, Triglycerides ("bad") cholesterol and fats and raise "good" cholesterol (HDL).  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## VALSARTAN

- Composition :** VALSARTAN TABLETS 40/80/160/320 MG  
**Indication :** Valsartan is used to treat high blood pressure and heart failure. It is also used to improve the chance of living longer after a heart attack. In people with heart failure, it may also lower the chance of having to go to the hospital for heart failure.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## VALSARTAN HCT

- Composition :** VALSARTAN & HCT 320/160/80 & 12.5 MG  
**Indication :** Valsartan and Hydrochlorothiazide is indicated for the treatment of hypertension, to lower blood pressure. Lowering blood pressure reduces the risk of fatal and nonfatal cardiovascular events, primarily strokes.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## CANDESARTAN



**Composition :** CANDESARTAN CILEXETIL TABLETS 4/8/16/32 MG

**Indication :** Candesartan Tablets are indicated for the treatment of hypertension and heart failure. They are used to lower blood pressure in adults and children ( $\geq 1$ -year-old) and to manage symptomatic heart failure in adults, improving symptoms and reducing the risk of hospitalization.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister

## CANDESARTAN + HYDROCHLOROTHIAZIDE



**Composition :** CANDESARTAN + HYDROCHLOROTHIAZIDE TABLETS 8+12.5/16+12.5/16+25/32+25 mg

**Indication :** Candesartan + Hydrochlorothiazide (HCT) Tablets are indicated for the treatment of hypertension (high blood pressure). This combination therapy helps to lower blood pressure by utilizing the complementary effects of candesartan, an angiotensin II receptor blocker (ARB), and hydrochlorothiazide, a thiazide diuretic. It is used in patients who require multiple medications to achieve effective blood pressure control.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister

## DABIGATRAN



**Composition :** DABIGATRAN CAPSULES 75/110/150 MG

**Indication :** Dabigatran Capsules are indicated for the prevention and treatment of thromboembolic events i.e. Prevention of stroke, deep vein thrombosis (DVT), venous thromboembolism (VTE) following hip replacement surgery.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister

## METOPROLOL



**Composition :** METOPROLOL TABLETS 25/50/100 MG

**Indication :** Metoprolol Tablets are indicated for the management of cardiovascular conditions i.e. Hypertension, Angina Pectoris, Heart Failure, Acute Myocardial Infarction.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister

## METOPROLOL + HCT



**Composition :** METOPROLOL + HCT TABLETS 50+12.5/100+12.5/100+25 MG

**Indication :** Metoprolol + HCT Tablets are indicated for the management of hypertension. This combination therapy is used to: Reduce blood pressure in patients with essential hypertension, improving overall cardiovascular health and reducing the risk of stroke, myocardial infarction, and renal complications.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## HYDROCHLOROTHIAZIDE

**Composition :** HYDROCHLOROTHIAZIDE TABLETS 12.5/25/50 MG

**Indication :** Hydrochlorothiazide is a thiazide diuretic used to treat hypertension (high blood pressure) and edema (fluid retention) associated with heart failure, kidney disorders, or liver cirrhosis. It helps reduce excess fluid in the body, lowering blood pressure and preventing complications like strokes and heart attacks.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## SPIRONOLACTONE

**Composition :** SPIRONOLACTONE TABLETS 25/50/100 MG

**Indication :** Spironolactone Tablets are indicated for the management of various conditions associated with fluid retention and electrolyte imbalances. Specifically, they are used to treat fluid retention in conditions such as congestive heart failure, liver cirrhosis, and nephrotic syndrome, Hypertension.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## ELECTROLYTES & MINERALS



### DIOCTAHEDRAL SMECTITE

**Composition :** DIOCTAHEDRAL SMECTITE POWDER 3.0 GM

**Indication :** DiOctahedral Smectite is used for Acute and Chronic Diarrhea, Irritable bowel syndrome in adults, Gastroesophageal reflux in infant and other conditions.

**Pack :** 10 Sachets x 3.0 gm



### GITRAL M

**Composition :** ORAL REHYDRATION SALTS 4.2/20.5/27.9 GM

**Indication :** Treatment of Diarrhoea by reversing the process of dehydration, acidosis and loss of electrolytes associated with diarrhoea, whether caused by bacteria, viruses, cryptosporidia or inappropriate nutrition.

**Pack :** 3, 10, 25, 50 and 100 Sachets



### GITRAL M PLUS

**Composition :** ORAL REHYDRATION SALTS 20.5G & ZINC SULPHATE TABLETS 20MG

**Indication :** Zinc is critical to body's ability to fight disease without enough zinc, the immune system is compromised and the gut is less to keep fluids in the body zinc supplementation can prevent diarrhoea and pneumonia through several mechanisms, notably through its ability to restore immunity in children who have zinc deficiency. Together, zinc + ORS are proven to stop diarrhoea, restore strength and even save lives.

**Pack :** Combikit of 2 Sachets + 1x10 Blister



### POTASSIUM CHLORIDE

**Composition :** POTASSIUM CHLORIDE ER TABLETS 600/750 MG

**Indication :** Potassium Chloride XR Tablets are indicated for the treatment and prevention of potassium deficiency in patients with hypokalemia. They are used to maintain normal potassium levels in the body, especially in patients experiencing potassium loss due to conditions such as chronic diarrhea, vomiting, or the use of certain diuretics.

**Pack :** 10x10 Blister & Bottle Pack



## ERECTILE DYSFUNCTION

### FOR MEN



**Composition :** SILDENAFIL & DAPOXETINE TABLETS 50/60 MG

**Indication :** The Sildenafil 50 mg+ Dapoxetine 60 mg Tablet is the highly recommended distinctive solution to get over the problem of premature ejaculation. The main objective of the drug is to delay the process of ejaculation resulting in enhanced sexual pleasure to a man suffering from the problem of pre-ejaculation.

**Pack :** 1 x 4, 1 x 10, 1 x 1 Blister



### LOVEGLOB

**Composition :** SILDENAFIL CITRATE TABLETS 50/100 MG

**Indication :** Sildenafil citrate in men with erectile dysfunction in an international, double-blind, placebo-controlled study: effect on the sexual experience and reducing feelings of anxiety the next intercourse attempt.

**Pack :** 1 x 4, 1 x 10, 1 x 1 Blister



### TADALAFIL

**Composition :** TADALAFIL TABLETS & TADALAFIL CHEWABLE TABLETS 5/10/20 MG

**Indication :** Tadalafil used to treat for erectile dysfunction. The recommended starting does of Tadalafil for use as needed in most patients is 10 mg, taken prior to anticipated sexual activity. The dose may be increased to 20 mg or decreased to 5 mg, based on individual efficacy and tolerance.

**Pack :** 1 x 10, 1 x 4, 1 x 1 Blister



### VARDENAFIL

**Composition :** VARDENAFIL TABLETS 5/10/20 MG

**Indication :** Vardenafil is used to treat male sexual function problems (Impotence or Erectile Dysfunction - ED). In a combination with sexual stimulation, Vardenfil works by increasing blood flow to the penis to help a man get and keep an erection.

**Pack :** 1 x 10, 1 x 4, 1 x 1 Blister



## GASTROLOGY



## AMEGHADOM

- Composition :** DOMPERIDONE TABLETS 10 MG  
**Indication :** For the relief of the symptoms of nausea and vomiting.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## BISOTIDINE

- Composition :** CIMETIDINE TABLETS 200/400 MG  
**Indication :** Gastroesophageal Reflux Disease (GERD), Heartburn, Acid Dyspepsia, Zollinger-Ellison Syndrome, Erosive Esophagitis.  
**Pack :** 1 x 10, 1 x 20, 3 x 10 and 10 x 10 Blister,



## DISETRON

- Composition :** ONDANSETRON ORALLY DISINTEGRATING TABLETS 4/8 MG  
**Indication :** Nausea and vomiting induced by Cytotoxic Chemotherapy and Radiotherapy, & for the prevention & treatment of Post-Operative Nausea & Vomiting (PONV).  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## ESOMEPRAGLOBE

- Composition :** ESOMEPRAZOLE MAGNESIUM TABLETS 20/40 MG  
**Indication :** Treatment of Erosive Reflux Oesophagitis, Prevention of Re-bleeding of Peptic Ulcers, Treatment of Zollinger Ellison Syndrome.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister




## ESOMEPRAGLOBE D

- Composition :** ESOMEPRAZOLE + DOMPERIDONE CAPSULES 20+10 MG  
**Indication :** Acid reflux, ulcers, heal acid damage to the stomach and oesophagus.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### RABEGLOB D

**Composition :** RABEPRAZOLE + DOMPERIDONE 20+10 MG  
**Indication :** Duodenal or Benign Gastric Ulcers, GERD.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### GLOMEZ-D

**Composition :** OMEPRAZOLE + DOMPERIDONE CAPSULES 20+10 MG  
**Indication :** GERD, NSAID related Ulcers, Peptic Ulcer associated Bleeding Zollinger Ellison Syndrome.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### GLOCID

**Composition :** RANITIDINE TABLETS 150/300 MG  
**Indication :** Ulcer in stomach & duodenum, hyper secretory upset stomach.  
**Pack :** 1 x 10, 3 x 10, 10 x 10 and 30 x 10 Blister



### GLOCID-D

**Composition :** RANITIDINE HCL + DOMEPRIDONE TABLETS 150+10 MG  
**Indication :** Intractable nausea and vomiting, benign duodenal ulcer, gastric ulcer, inflammation of the esophagus, acid peptic disease, treatment for symptoms associated with idiopathic or diabetic gastroparesis and other conditions.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### LANSOGLOB

**Composition :** LANSOPRAZOLE DELAYED RELEASE CAPSULES 30 MG  
**Indication :** Treatment of duodenal and gastric ulcer, symptomatic gastro-esophageal reflux disease, Zollinger-Ellison Syndrome.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## MAGNESIUM TABLETS

**Composition :** MAGNESIUM TABLETS 100/400 MG

**Indication :** Low amounts of magnesium in the blood. Normal functioning of cells, nerves, muscles, bones and the heart.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## OMEPRAGLOBE

**Composition :** OMEPRAZOLE DELAYED RELEASE CAPSULES 20/40 MG

**Indication :** Prevent ulcers of the swallowing tube (Esophagus). Gastroesophageal reflux disease (GERD, Acid Reflux).

**Pack :** 1 x 10, 1 x 14, 1 x 15, 2 x 7, 3 x 10 and 10 x 10 Blister & Strip



## PANGLO-DSR

**Composition :** PANTOPRAZOLE SODIUM (EC) & DOMPERIDONE (SR) CAPSULES 40+10 MG

**Indication :** Duodenal or Benign Gastric Ulcers, Acid Reflux Disease, Reflux Esophagitis, Dyspepsia & Bloating.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## PENTAGLOB

**Composition :** PANTOPRAZOLE TABLETS 20/40 MG

**Indication :** GERD, Ulcers, Peptic Ulcer, Bleeding Zollinger Elison Syndrome.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## RABEGLOB

**Composition :** RABEPRAZOLE SODIUM ENTERIC COATED TABLETS 10/20/40 MG

**Indication :** Healing of Erosive or Ulcerative GERD in adults. Healing of Duodenal Ulcers in adults.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## HYOSCINE BUTYLBROMIDE

- Composition :** HYOSCINE BUTYLBROMIDE TABLETS 10 MG  
**Indication :** Crampy abdominal pain remove bolds, esophageal spasms, renalcolic & bladder spasms, improve respiratory secretions.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## LOPERAGLOBE

- Composition :** LOPERAMIDE CAPSULES 2 MG  
**Indication :** Acute nonspecific diarrhea, mild traveler's diarrhea, irritable bowel syndrome, chronic diarrhea due to bowel resection and chronic diarrhea secondary to inflammatory bowel disease.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## URSOGLOB

- Composition :** URSODEOXYCHOLIC ACID TABLETS 150/300 MG  
**Indication :** Treatment of Primary Biliary Cirrhosis (PBC) and for the dissolution of small to medium sized radiolucent, cholesterol- rich gall-stones.  
**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## ALUMINIUM HYDROXIDE + MAGNESIUM HYDROXIDE

- Composition :** ALUMINIUM HYDROXIDE + MAGNESIUM HYDROXIDE TABLETS 200+200/400+400/500+500 MG  
**Indication :** Aluminum Hydroxide + Magnesium Hydroxide Tablets are indicated for the symptomatic relief of dyspepsia and heartburn. This combination therapy is used to neutralize gastric acid and reduce symptoms associated with acid reflux, gastritis, and peptic ulcers.  
**Pack :** 10 x 10 Blister



## NEPHROLOGY, UROLOGY &amp; HEPATIC



## DEFFERGLOB

**Composition :** DEFERASIROX TABLETS 100/250/400/500 MG

**Indication :** This medication is used to treat ongoing high levels of iron in the body caused by multiple blood transfusions. It is also used to treat high levels of iron in people with a certain blood disorder who do not require blood transfusions (non-transfusion-dependent thalassemia).

**Pack :** 4 x 7, 1 x 10, 3 x 10 and 10x10 Blister



## FEBUGLOB

**Composition :** FEBUXOSTAT TABLETS 40/80 MG

**Indication :** Febuxostat is indicated for the prevention and treatment of hyperuricaemia in adult patients undergoing chemotherapy for haematologic malignancies at intermediate to high risk of Tumor Lysis Syndrome (TLS).

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



## RENAGLOB

**Composition :** KETOANALOGUES PLUS ESSENTIAL AMINO ACID TABLET

**Indication :** Ketoanalogues/Essential Amino Acid is used for the treatment, control, prevention and improvement of inefficient protein metabolism in chronic renal insufficiency and Uremia.

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



## SELAGLOB

**Composition :** SEVELAMER HYDROCHLORIDE TABLETS 400/800 MG  
SEVELAMER CARBONATE TABLETS 400/800 MG

**Indication :** Sevelamer is used to lower high blood phosphorus (phosphate) levels in patients who are on dialysis due to severe kidney disease. Dialysis removes some phosphate from your blood.

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



## URIGLOB / URIGLOB D

**Composition :** TAMSULOSIN HCL MODIFIED RELEASE CAPSULES 400 MG

**Indication :** TAMSULOSIN & DUTASTERIDE CAPSULES 0.4+0.5 MG

Tamsulosin 400 mg contains an active ingredients in the capsules which is a so called Alpha 1A-blocker that reduces muscle contraction in the Prostate and Urethra. This facilitates the flow of urine through the Urethra and Aids Urination.

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



## KALARA

**Composition :** CALCIUM POLYSTYRENE SULFONATE 15 G

**Indication :** Calcium Polystyrene Sulfonate is used for reducing blood levels of potassium. It is used for the treatment of Hyperkalemia (High Potassium Levels) associated with kidney conditions such as anuria, severe oliguria, chronic kidney disease, etc. It is also used to reduce high potassium levels in patients who are undergoing regular dialysis.

**Pack :** 10 Sachets



## CINACALCET

**Composition :** CINACALCET TABLETS 30/60/90 MG

**Indication :** Cinacalcet Tablets are indicated for the treatment of conditions related to elevated parathyroid hormone (PTH) levels. Specifically, they are used to manage secondary hyperparathyroidism in patients with chronic kidney disease (CKD) on dialysis, treat hypercalcemia and control hyperparathyroidism in patients with parathyroid carcinoma.

**Pack :** 3 x 10 and 10 x 10 Blister



## ALFUZOSIN

**Composition :** ALFUZOSIN EXTENDED-RELEASE TABLETS 10 MG

**Indication :** Alfuzosine Tablets are indicated for the treatment of benign prostatic hyperplasia (BPH). Specifically, they are used to improve urinary symptoms associated with BPH, such as urinary frequency, urgency, and nocturia. Relax smooth muscle in the prostate and bladder neck, thereby improving urine flow and reducing bladder outlet obstruction.

**Pack :** 3 x 10 and 10 x 10 Blister



## ALLOPURINOL

**Composition :** ALLOPURINOL TABLETS 100/300 MG

**Indication :** Allopurinol Tablets are indicated for the management of hyperuricemia associated with conditions such as gout and certain types of kidney stones. Reduce elevated serum uric acid levels in patients with chronic gout, thereby preventing gout attacks and the formation of uric acid crystals. Prevent the formation of uric acid stones in patients with a history of recurrent kidney stones.

**Pack :** 3 x 10 and 10 x 10 Blister



## FINASTERIDE

**Composition :** FINASTERIDE TABLETS 1/5 MG

**Indication :** Finasteride Tablets are indicated for the management of conditions related to the prostate and hair loss. Benign Prostatic Hyperplasia (BPH): To reduce the size of the prostate and improve urinary symptoms such as difficulty in urination, urinary frequency, and urgency. Androgenetic Alopecia: To treat male pattern baldness by promoting hair regrowth and slowing hair loss in men.

**Pack :** 3 x 10 and 10 x 10 Blister



## ONCOLOGY &amp; IMMUNOSUPPRESSANTS PRODUCTS



## ABIRAGLOB

**Composition :** ABIRATERONE ACETATE TABLETS 250/500 MG

**Indication :** For the treatment of metastatic castration resistant prostate cancer in adult men who are asymptomatic or mildly symptomatic after failure of androgen deprivation therapy.

**Pack :** 60/120 Tablets Bottle



## DASAGLOB

**Composition :** DASATINIB TABLETS 20/50/70 MG

**Indication :** Newly diagnosed Philadelphia chromosome positive (Ph+) chronic myelogenous leukaemia (CML) in the chronic phase. Chronic, accelerated or blast phase CML with resistance or intolerance to prior therapy including imatinib mesylate.

**Pack :** 30 Tablets Bottle



## ERLOGLOB

**Composition :** ERLOTINIB TABLETS 100/150 MG

**Indication :** Locally advanced or metastatic non-small cell lung cancer (NSCLC) with EGFR activating mutations, either as 1 line or as switch maintenance treatment and stable disease after 1 line chemotherapy. Also used for metastatic pancreatic cancer.

**Pack :** 30 Tablets Bottle



## ETOGLOB

**Composition :** ETOPOSIDE CAPSULES 50/100 MG

**Indication :** Treatment of small cell lung cancer (as first line in combination with other agents), refractory testicular tumors, also used for.. Hodgkin's disease, Malignant (Non-hodgkin's) lymphomas, Acute Non-lymphocytic leukemia

**Pack :** 4 & 8 Capsules Bottle



## LENAGLOB

**Composition :** LENALIDOMIDE CAPSULES 5/10/15/25 MG

**Indication :** For the treatment of newly diagnosed Multiple Myeloma, Myelodysplastic Syndromes, Mantle Cell Lymphoma.

**Pack :** 30 Tablets Bottle



## GLOSUNIB

**Composition :** SUNITINIB MALATE CAPSULES 12.5/25/50 MG

**Indication :** The treatment of gastrointestinal stromal tumor (GIST) after disease progression on or intolerance to imatinib mesylate the treatment of advanced renal cell carcinoma (RCC).

**Pack :** 30 Capsules Bottle



## EVEROLIMUS

**Composition :** EVEROLIMUS TABLETS 0.75/1/5/10 MG

**Indication :** Treatment of adult patients with renal angiomyolipoma associated with Tuberous Sclerosis Complex (TSC).  
Treatment of patients with subependymal giant cell (SEGA) associated with Tuberous Sclerosis Complex (TSC).

**Pack :** 30 Tablets Bottle & 1x10 Blister



## AZAGLOB

**Composition :** AZATHIOPRINE TABLETS 50/100 MG

**Indication :** To enhance the survival of organ transplants, such as renal transplants, cardiac transplants and hepatic transplant and to reduce the corticosteroid requirements of renal transplant recipients.

**Pack :** 10x10 Blister



## MERKGLOB

**Composition :** MERCAPTOPYRINE TABLETS 50 MG

**Indication :** For the treatment of Acute Lymphoblastic Leukaemia (ALL) / Acute Promyelocytic Leukaemia (APL) / Acute Myeloid Leukaemia M3 (AML M3).

**Pack :** 10x10 Blister



## GLOUREA

**Composition :** HYDROXYUREA CAPSULES 200/500 MG

**Indication :** Chronic Myeloid Leukaemia, cancer of the Cervix & other solid type tumors in conjunction with Radio therapy, Sickle Cell Anemia (only approved drug), reduces frequency of painful crises, reduce the need for blood transfusions.

**Pack :** 10x10 Blister



## GLOTEMOZ

**Composition :** TEMOZOLOMIDE CAPSULES 20/100/250 MG

**Indication :** For the treatment of adult patients with newly-diagnosed glioblastoma multiforme, for management of malignant glioma.

**Pack :** 5 Tables Bottle & 1 x 5 Blister



### DEXACEE

**Composition :** DEXAMETHASONE TABLETS 4/8 MG

**Indication :** Treatment of cancer Mammae, Ovarii, Prostate or Testis, Prophylaxis of Emesis induced by emetogenic chemotherapy, cerebral oedema or increased intracranial pressure due to brain tumor.

**Pack :** 10x10 & 5x8 Blister



### LAPAGLOB

**Composition :** LAPATINIB TABLETS 250 MG

**Indication :** Lapatinib Tablets 250 mg is indicated for the treatment of patients with advanced or metastatic breast cancer whose tumors overexpress human epidermal growth factor receptor 2 (HER2) and who have received prior therapy including an anthracycline, a taxane, and trastuzumab.

**Pack :** 3x10, 1x10 Blister



### MYCOFET

**Composition :** MYCOPHENOLATE MOFTIL TABLETS 250/500 MG

**Indication :** For the prophylaxis of acute transplant rejection in patients receiving allogeneic renal, cardiac or hepatic transplants (in combination with cyclosporin and corticosteroids).

**Pack :** 1x10, 10x10 Strip



### ELTROGLOB

**Composition :** ELTROMBOPAG TABLETS 25/50 MG

**Indication :** Eltrombopag Tablets are indicated for the treatment of Chronic Immune Thrombocytopenia (ITP): To increase platelet counts in patients. Severe Aplastic Anemia: In combination with other therapies for the treatment of severe aplastic anemia in adults and pediatric patients who have not responded to immunosuppressive therapy.

**Pack :** 3x10, 10x10 Blister & 14 Tab Bottle pack



### ENZAGLOB 40

**Composition :** ENZALUTAMIDE CAPSULES 40/80 MG

**Indication :** Enzalutamide is used to treat prostate cancer. This medication belongs to a class of drugs known as anti-androgens (anti-testosterone).

**Pack :** 1x10, 1x14 Blister



## ANAGLOB

**Composition :** ANASTROZOLE TABLETS 1 MG

**Indication :** Adjuvant treatment of hr-p early breast cancer, first line treatment of hr-p or hr-unknown locally advanced or metastatic breast cancer, second line treatment of advanced breast cancer.

**Pack :** 3 x 14 and 3 x 10 Blister



## GLOTRAZ

**Composition :** LETROZOLE TABLETS 2.5 MG

**Indication :** Adjuvant treatment of HR-P invasive early breast cancer, Extended adjuvant Treatment of Hormone dependent invasive breast cancer, First line treatment for hormone dependent advanced breast cancer.

**Pack :** 3 x 10 Blister



## TREXGLO

**Composition :** METHOTREXATE TABLETS 2.5/5/10 MG

**Indication :** For the treatment of neoplastic disease, Treatment of severe cases of uncontrolled psoriasis. For treatment of adults with severe, active, classical or definite rheumatoid arthritis.

**Pack :** 3 x 10 Blister



## FLUTAGLOB

**Composition :** FLUTAMIDE TABLETS 250 MG

**Indication :** Flutamide is indicated for the treatment of advanced prostatic carcinoma in which suppression of testosterone effects is indicated. flutamide may be used in combination with an LHRH agonist, both on commencement of treatment or as an adjunctive therapy in patients already receiving an LHRH agonist. Flutamide may also be used in surgically castrated patients.

**Pack :** 3 x 10 Blister



## THALIGLOB

**Composition :** THALIDOMIDE CAPSULES 50/100 MG

**Indication :** As First line Therapy (in combination with melphalan and prednisone) for treatment of patients with untreated of patients with untreated multiple myeloma, aged > 65 years or ineligible for high dose chemotherapy

**Pack :** 3 x 10 Blister



## GLONIL

**Composition :** NILOTINIB CAPSULES 200 MG

**Indication :** Newly Diagnosed Philadelphia Chromosome Positive Chronic Myelogenous leukemia (CML) in the chronic phase.

**Pack :** 1 x 10 Blister



## GLOMATINIB

Composition : **IMATINIB MESYLATE TABLETS 100/400 MG**

Indication : Chronic Myeloid Leukemia, Acute Lymphoblastic Leukemia, Hyper-eosinophilic Syndrome or CEL, Aggressive Systemic Mastocytosis, Dermatofibrosarcoma protuberans, GIST, MDS/MPD.

Pack : 1 x 10 & 10 X 10 Blister



## PROSTAGLOB

Composition : **BICALUTAMIDE TABLETS 50/150 MG**

Indication : Treatment of advanced prostate cancer, alone or in combination with luteinizing hormone releasing hormone (LHRH) analogue.

Pack : 1 x 10 & 3 x 10 Blister



## GLOMELFA

Composition : **MELPHALAN TABLETS 2/5 MG**

Indication : Treatment of multiple myeloma and advanced ovarian adenocarcinoma. Alone or in combination for treatment of a proportion of patients suffering from advanced breast carcinoma.

Pack : 1 x 10, 5 x 10 & 1 x 10 Blister



## GLOCAPX

Composition : **CAPECITABINE TABLETS 150/500 MG**

Indication : Adjuvant treatment of patients following surgery of Stage-III (Dukes' Stage C) Colon Cancer, Metastatic Colorectal Cancer, Locally Advanced or Metastatic Breast Cancer after failure of Cytotoxic Chemotherapy (in combination).

Pack : 1 x 10 Blister



## GLOCARBAZIN

Composition : **PROCARBAZINE CAPSULES 50 MG**

Indication : Matulane is indicated for use in combination with other anticancer drugs for the treatment of Stage III and IV Hodgkin's disease.

Pack : 1 x 10 Blister



## TCOGLOB

Composition : **TACROLIMUS CAPSULES 0.5/1/5 MG**

Indication : Prophylaxis of transplant rejection in liver, kidney or heart allograft recipients.

Pack : 1 x 10 & 10 X 10 Blister



## ZFTIGLO

**Composition :** GEFITINIB TABLETS 250 MG

**Indication :** Treatment of adult patients with locally advanced or metastatic nonsmallcell lung cancer (NSCLC) with activating mutations of EGFR TK.

**Pack :** 30 Tablets Bottle & 1 x 10 Blister



## SIROGLOB

**Composition :** SIROLIMUS TABLETS 1/2 MG

**Indication :** For the prophylaxis of organ rejection in adult patients at low to moderate immunological risk receiving a renal transplant.

**Pack :** 1 x 10 Strips



## TAMGLOB

**Composition :** TAMOXIFEN TABLETS 10/20 MG

**Indication :** For the treatment of breast cancer.  
To stimulate ovulation in anovulatory infertility.

**Pack :** 1 x 10, 3x10 & 10x10 Blister



## GLOMUSTIN

**Composition :** ESTRAMUSTINE PHOSPHATE CAPSULES 140 MG

**Indication :** Carcinoma of the prostate, especially in cases unresponsive to, or relapsing after, treatment by conventional oestrogens (stibioestrol, polyoestradiol phosphate etc.) or by orchidectomy.

**Pack :** 1 x 10 Blister



## GLOSULFAN

**Composition :** BUSULPHAN TABLETS 2 MG

**Indication :** Busulphan is used in pediatrics and adults in combination with cyclophosphamide or fludarabine/clofarabine as a conditioning agent prior to bone marrow transplantation, especially in chronic myelogenous leukemia (CML).

**Pack :** 1 x 10 Blister



## MYCOFET

**Composition :** MYCOPHENOLATE SODIUM TABLETS 180/360 MG

**Indication :** Mycophenolate Sodium Tablets is used in combination with other medications to keep your body from attacking and rejecting your transplanted kidney. It belongs to a class of medications called immunosuppressant.

**Pack :** 1 x 10 Strip



## APALUTAMIDE

**Composition :** APALUTAMIDE TABLETS 60 MG

**Indication :** Apalutamide is used to treat men with prostate cancer. This medication belongs to a class of drugs known as anti-androgens (anti-testosterone). It works by blocking the effects of testosterone to slow the growth.

**Pack :** 30 Tablets Bottle



## BINIMETINIB

**Composition :** BINIMETINIB TABLETS 15 MG

**Indication :** Binimetinib is indicated, in combination with encorafenib, for the treatment of patients with unresectable or metastatic melanoma with a BRAF V600E or V600K mutation

**Pack :** 30 Tablets Bottle



## CABOZANTINIB

**Composition :** CABOZANTINIB TABLETS 20 / 80 MG

**Indication :** For the treatment of patients with advanced renal cell carcinoma (RCC). For the treatment of patients with Hepatocellular Carcinoma (HCC) who have been previously treated with sorafenib.

**Pack :** 30 Tablets Bottle



## IBRUTINIB

**Composition :** IBRUTINIB CAPSULES 140/400 MG

**Indication :** For the treatment of adult patients with: mantle cell lymphoma (MCL) who have received at least one prior therapy. -Chronic Lymphocytic leukemia (CLL) who have received at least one prior therapy. - Chronic lymphocytic leukemia with 17p deletion

**Pack :** 30 Capsule Bottle



## LAROTRECTINIB

**Composition :** LAROTRECTINIB CAPSULES 25 / 100 MG

**Indication :** Larotrectinib is for adults and children with solid tumors that test positive for NTRK genes, which can help cancerous tumors grow.

**Pack :** 30 Capsule Bottle



## NIRAPARIB

**Composition :** NIRAPARIB CAPSULES 100 MG

**Indication :** Niraparib is used as first-line maintenance treatment in patients with advanced ovarian cancer, fallopian tube cancer, or primary peritoneal cancer.

**Pack :** 30 Capsule Bottle







## STERODIAL ANTI INFLAMMATORY



### BETAMETHASONE

**Composition :** BETAMETHASONE TABLETS 0.5 MG

**Indication :** Betamethasone 0.5 mg tablet is similar to the naturally occurring steroidal hormones (hydrocortisone/cortisone) produced by the adrenal glands. It is used to treat internal and external swelling associated with diseases like asthma, colitis, arthritis, severe skin allergies etc.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### DEFLAZACORT

**Composition :** DEFLAZACORT TABLETS 1/6/18/30 MG

**Indication :** Deflazacort is a steroid that reduces inflammation in the body. Deflazacort is not a cure for muscular dystrophy, but this medicine may improve muscle strength and slow the progression of disability.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### DEXACEE

**Composition :** DEXAMETHASONE TABLETS 0.5/1 MG

**Indication :** Dexamethasone is used to treat conditions such as arthritis, blood/hormone/immune system disorders, allergic reactions, certain skin and eye conditions, breathing problems, certain bowel disorders and certain cancers.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### PREDNISOLONE

**Composition :** PREDNISOLONE TABLETS 5/10/20 MG

**Indication :** Prednisolone Tablets are a corticosteroid which is used to relieve inflammation in conditions like arthritis and allergic disorders. Use with caution in the patients with Diabetes and other kidney disease.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### SULBUTIAMINE

**Composition :** SULBUTIAMINE TABLETS 200 MG

**Indication :** Sulbutiamine is man-made chemical, similar to B Vitamin Thiamine. Unlike Thiamine which dissolves in water, Sulbutiamine dissolves in fats. Sulbutiamine is able to increase Thiamine levels in the brain.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## VITAMINS AND MINERALS



## CALCIGLO D3

**Composition :** CALCIUM CARBONATE WITH VITAMIN D3 TABLETS

**Indication :** As an adjunct to therapy for osteoporosis, as therapeutic supplementation of malnutrition e.g. in pregnancy and established Vitamin D dependent osteomalacia. Prevention and treatment of Calcium/ Vitamin D deficiency especially in the housebound and institutionalized elderly subjects.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## DIOSMIN &amp; HESPERIDIN

**Composition :** DIOSMIN & HESPERIDIN TABLETS 50/450 MG

**Indication :** Haemorrhoids Chronic Venous Disease.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## FOLIC ACID

**Composition :** FOLIC ACID TABLETS 5 MG

**Indication :** Folic Acid is used for preventing and treating low blood levels of folate (folate deficiency), as well as its complications, including "tired blood" (Anemia) and the inability of the bowel to absorb nutrients properly. Folic Acid is also used for other conditions commonly associated with folate deficiency, including ulcerative colitis, liver disease, alcoholism and kidney dialysis.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## GLUCOSAMINE

**Composition :** GLUCOSAMINE TABLETS 500 MG

**Indication :** Glucosamine is sugar protein that helps your body build cartilage (the hard connective tissue located mainly on the bones near your joints). Glucosamine is a naturally occurring substance found in bones, bone marrow.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## G-TOTAL

**Composition :** MULTIVITAMIN & MULTIMINERAL CAPSULES

**Indication :** This medication is a multivitamin product used to treat or prevent vitamin deficiency due to poor diet, certain illnesses or during pregnancy. Vitamins are important building blocks of the body and help keep you in good health.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### GTOTAL-F

- Composition :** FERROUS FUMARATE, CYANOCOBALAMIN, FOLIC ACID & ZINC SULPHATE TABLETS
- Indication :** Treatment and prophylaxis of uncomplicated iron deficiency anaemia, vitamin B12 deficiencies, Addisonian anemia, folic acid deficiency and other conditions.
- Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



### HEMGLOB

- Composition :** FERROUS SULPHATE TABLETS 200 MG
- Indication :** Ferrous sulfate (or sulphate) is a medicine used to treat and prevent iron deficiency anaemia. Iron helps the body to make healthy red blood cells, which carry oxygen around the body.
- Pack :** 10x10 Blister



### HEMGLOB PLUS

- Composition :** FERROUS SULPHATE & FOLIC ACID TABLETS 200+0.4 MG
- Indication :** Ferrous sulphate + folic acid tablets are used to treat iron deficiency anemia (a lack of red blood cells caused by having too little iron in the body). Ferrous sulphate is a type of iron and is an essential body mineral whereas folic support the body for new cell generation & maintenance.
- Pack :** 10x10 Blister



### VITAMIN C 1000 CHEWABLE TABLETS

- Composition :** VITAMIN C CHEWABLE TABLETS 500/1000 MG
- Indication :** Vitamin C Tablets is a vitamin used to correct vitamin C deficiency and to increase the intestinal absorption of iron.
- Pack :** 100 Tablets Bottle



## LAXATIVE



## BISAMED

**Composition :** BISACODYL TABLETS 5 MG

**Indication :** Bowel block, belly pain, upset stomach, stimulant laxative

**Pack :** 1 x 10, 3 x 10, 7 x 15 and 10 x 10 Blister



## PSYLLIUM HUSK POWDER

**Composition :** PSYLLIUM HUSK POWDER

**Indication :** This medication is used to treat constipation. It increases the bulk in your stool, an effect that helps to cause movement of the intestines.

**Pack :** HDPE Bottle (36.5 OZ)



## SENNOSIDES

**Composition :** SENNOSIDES TABLETS 8.6 MG

**Indication :** Sennosides are used to treat constipation. Sennosides are known as stimulant laxatives. They work by keeping water in the intestines, which causes movement of the intestines.

**Pack :** 600/800 tablet's Jar



# EFFERVESCENT & NUTRACEUTICAL PRODUCTS



## RAP'DFIZZ BOOSTER +

VITAMIN C EFFERVESCENT TABLETS 500/1000 MG  
ORANGE & ELDERBERRY FLAVOUR  
Pack : 16 Tablets Strip Pack / 20 Tablets Tube Pack



## RAP'DFIZZ IMMUNIFIER

VITAMIN C + ZINC EFFERVESCENT TABLETS 1000/12.5 &  
500/12.5 MG ORANGE & ELDERBERRY FLAVOUR  
Pack : 16 Tablets Strip Pack / 20 Tablets Tube Pack



## RAP'DFIZZ CALCIFY

CALCIUM CARBONATE + VITAMIN D3 EFFERVESCENT TABLETS  
Pack : 16 Tablets Strip Pack / 20 Tablets Tube Pack



## RAP'DFIZZ RESTORE

ELECTROLYTES ANTIOXIDANT CARBOHYDRATES 2:1  
EFFERVESCENT TABLETS  
Pack : 16 Tablets Strip Pack / 20 Tablets Tube Pack



**GITRAL**

**ELECTROLYTE EFFERVESCENT TABLETS**  
Pack : 16 Tablets Strip Pack / 20 Tablets Tube Pack



**GLOPAR-500**

**PARACETAMOL EFFERVESCENT TABLETS 500/1000 MG**  
Pack : 16 Tablets Strip Pack / 20 Tablets Tube Pack



**GLOPAR-C**

**PARACETAMOL & CAFFEINE EFFERVESCENT TABLETS**  
500+65/500+30 MG  
Pack : 16 Tablets Strip Pack / 20 Tablets Tube Pack




**DICLOGLOB - P**

**PARACETAMOL 500 MG & DICLOFENAC SODIUM 50 MG**  
EFFERVESCENT TABLETS  
Pack : 4 x 4 Tablets Strip Pack



**EFFERVESCENT & NUTRACEUTICAL PRODUCTS**



**VIATMIN D3 1000 IU**

VIATMIN D3 1000 IU/ 5000 IU TABLETS  
 PACK : 90 TABLETS BOTTLE



**ZINC 100**

ZINC CAPLETS 100 MG  
 Pack : 60 Caplet Bottle



**SILDENAFIL 50**

SILDENAFIL CITRATE EFFERVESCENT TABLETS 50 MG  
 Pack : 5 x 4 Effervecent Tablets Strip Pack



**MUCOXIT 600**

ACETYLCYSTEINE EFFERVECENT TABLETS 600 MG  
 Pack : 10 x 10 Effervecent Tablets Strip Pack



## COMING SOON...

- Acetaminophen, Chlorpheniramine Maleate, Dextromethorphan HBR, Phenylephrine HCL 250/2/10/5 mg
- Aspirin, Chlorpheniramine Maleate, Dextromethorphan HBR, Phenylephrine Bitartrate 325/2/10/7.5 mg
- Anhydrous Citric Acid, Aspirin, Sodium Bicarbonate 1000/325/1916 mg
- Aspirin, Chlorpheniramine Maleate, Phenylephrine Bitartrate 325/2/7.8 mg
- Calcium Carbonate 500/1000 mg
- Acetaminophen USP, Diphenhydramine HCL 500/25 mg
- ZMA(Zinc Magnesium Aaspartate)
- OMEGA 3,6 & 9
- BCCA (Branched Chain Amino Acid)
- Korean Panax Ginseng
- Probiotic
- Omega-3 Fish Oil
- Vitamin B12 Capsule
- L-Arginine
- Fat Burner
- Biotin
- Vitamin C + Zinc + Vitamin D3 Chewable Tablets
- Ferrous Ascorbate with Folic acid Tablets
- Vitamin B-Complex Tablets
- Silymarin, L-Glutathione, N-Acetylcysteine, L-Carnitine, Choline with Antioxidants & Micronutrients Tablets.
- Digestive Enzyme Capsules
- Ginseng, Lycopene, Multivitamin & Multimineral Capsules
- Biotin, Multivitamin & Multimineral Tablets
- Lutein, Zeaxanthin and Multivitamin Capsules
- Calcium+ Magnesium+ Vitamin D3+ Zinc Tablets
- Glucosamine, Collagen peptides II & Vitamin C Capsule



## CEPHALOSPORINS & BETA LACTAM



### GLOXIM

**Composition :** CEFIXIME TABLETS USP 100 / 200 / 400 MG

**Indication :** Cefixime is indicated for the treatment of certain infections caused by susceptible bacteria Label, 12, 13. These infections include respiratory tract infections, otitis media, skin and skin structure infections, bone infections and genitourinary tract infections.

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



### GLOXIM DT

**Composition :** CEFIXIME DISPERSIBLE TABLETS 50 / 100 / 200 / 400 MG

**Indication :** For use in the treatment of the following infections when caused by susceptible strains of the designated microorganisms: (1) uncomplicated UTI caused by E. coli and P.mirabilis, (2) otitis media caused by H. influenzae (beta-lactamase +ve and -ve strains), M. catarrhalis (most of which are beta-lactamase +ve), and S. pyogenes, (3) pharyngitis and tonsillitis caused by S. pyogenes, (4) acute bronchitis and acute exacerbations of chronic bronchitis caused by S. pneumoniae and H. influenzae (beta-lactamase +ve and -ve strains), and (5) uncomplicated gonorrhea (cervical/urethral) caused by N. gonorrhoeae (penicillinase- and non-penicillinase-producing strains).

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



### GLOXIM CAPSULES

**Composition :** CEFIXIME USP CAPSULES 100 / 200/ 400 MG

**Indication :** Cefixime is indicated for the treatment of certain infections caused by susceptible bacteria Label, 12, 13. These infections include respiratory tract infections, otitis media, skin and skin structure infections, bone infections and genitourinary tract infections.

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



### GLOXIM-O

**Composition :** CEFIXIME 100/200 MG AND OFLOXACIN 100/200 TABLETS

**Indication :** CEFIXIME+OFLOXACIN belongs to a group of medications called 'cephalosporin antibiotics' primarily used for the treatment of a wide range of bacterial infections of the skin, ears, lungs, prostate and urinary tract. Besides this, it also treats typhoid fever and STDs (sexually transmitted diseases) like chlamydia and gonorrhoea. A bacterial infection happens when harmful bacteria enter inside the body and start multiplying. These harmful bacteria are responsible for producing toxins that can cause diseases like strep throat, urinary tract infection (UTI), bacterial food poisoning, gonorrhoea, bacterial meningitis, cellulitis, Lyme disease, and tetanus. CEFIXIME+OFLOXACIN does not work against viral infections like flu and colds

**Pack :** 1x10, 3x10 and 10x10 Blister




### GLOROXIM DT

- Composition :** CEFUROXIME AXETIL DISPERSIBLE TABLET 125 / 250 MG  
**Indication :** Cefuroxime is a cephalosporin antibiotic and is used to treat certain infections caused by bacteria. These include pharyngitis, otitis media, sinusitis, and bronchitis. They also include urinary tract infections, gonorrhea, Lyme disease, and impetigo.  
**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



### GLOROXIM

- Composition :** CEFUROXIME AXETIL TABLETS USP 250/500 MG  
**Indication :** For the treatment of the following infections (skin, UTI, ENT) caused by; S. pneumoniae, H. influenzae, staphylococci, S.pyogenes (Group A beta-hemolytic streptococci), E.coli, P.mirabilis, Klebsiella sp, coagulase-negative staphylococci and Streptococcus pyogenes.  
**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



### CEFUROXIME AXETIL & CLAVULANATE POTASSIUM

- Composition :** CEFUROXIME AXETIL AND CLAVULANATE POTASSIUM TABLETS  
**Indication :** 250+125/500+125 MG  
 CEFUROXIME+CLAVULANIC ACID belongs to the group of medicines called antibiotics used to treat various bacterial infections such as pharyngitis/tonsillitis (throat infections), otitis media (ear infections), sinusitis (infection of the sinuses), gonorrhoea (sexually transmitted disease), Lyme disease  
**Pack :** 1x10, 3x10 and 10x10 Blister



### GLOCEFIM

- Composition :** CEPHALEXIN CAPSULES USP 50 / 100 / 250 / 500 MG  
**Indication :** Cefalexin is a semi synthetic cephalosporin antibiotic for oral administration. Cefalexin is indicated in the treatment of the following infections: Respiratory tract infections; otitis media; skin and soft tissue infections; bone and joint infections; genito-urinary infections, including acute prostatitis and dental infections.  
**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



### GLOLEXIN DT

- Composition :** CEFALEXIN DISPERSIBLE TABLETS 125/250 MG  
**Indication :** Cefalexin is a semi synthetic cephalosporin antibiotic for oral administration. Cefalexin is indicated in the treatment of the following infections: Respiratory tract infections; otitis media; skin and soft tissue infections; bone and joint infections; genito-urinary infections, including acute prostatitis and dental infections.  
**Pack :** 1x10, 3x10 and 10x10 Blister



### GLOLEXIN

**Composition :** CEFALEXIN TABLETS USP 250 / 500 MG

**Indication :** Cefalexin is a semi synthetic cephalosporin antibiotic for oral administration. Cefalexin is indicated in the treatment of the following infections: Respiratory tract infections; otitis media; skin and soft tissue infections; bone and joint infections; genito-urinary infections, including acute prostatitis and dental infections.

**Pack :** 1x10, 3x10 and 10x10 Blister



### DROXIGLO

**Composition :** CEFADROXIL TABLETS USP 500 MG

**Indication :** For the treatment of the following infections (skin, UTI, ENT) caused by; S. pneumoniae, H. influenzae, staphylococci, S.pyogenes (Group A beta-hemolytic streptococci), E.coli, P.mirabilis, Klebsiella sp, coagulase-negative staphylococci and Streptococcus pyogenes.

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister




### DROXIGLO DT

**Composition :** CEFADROXIL DISPERSIBLE TABLETS 250/500 MG

**Indication :** For the treatment of the following infections (skin, UTI, ENT) caused by; S. pneumoniae, H. influenzae, staphylococci, S. pyogenes (group A beta-hemolytic streptococci), E. coli, P. mirabilis, Klebsiella sp, coagulase-negative staphylococci and Streptococcus pyogenes.

**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



### CEFADROXIL AND CLVULANATE POTASSIUM

**Composition :** CEFADROXIL AND CLAVULANATE POTASSIUM TABLETS 500+125

**Indication :** CEFADROXIL+CLAVULANIC ACID belongs to the group of medicines called antibiotics used to treat various bacterial infections such as urinary tract, skin, chest, and throat infections. Bacterial infections are caused due to the multiplication of harmful bacteria inside or on the body.

**Pack :** 1x10, 3x10 and 10x10 Blister



### CEFPODOXIME PROXETIL DISPERSIBLE TABLETS

**Composition :** CEFPODOXIME PROXETIL DISPERSIBLE TABLETS 100/200 MG

**Indication :** Cefpodoxime is used to treat certain infections caused by bacteria such as bronchitis (infection of the airway tubes leading to the lungs); pneumonia; gonorrhea (a sexually transmitted disease); and infections of the skin, ear, sinuses, throat, tonsils, and urinary tract

**Pack :** 1x10, 3x10 and 10x10 Blister



### CEFPODOXIME PROXETIL TABLETS

**Composition :** CEFPODOXIME PROXETIL TABLETS 100/200 MG  
**Indication :** Cefpodoxime is used to treat certain infections caused by bacteria such as bronchitis (infection of the airway tubes leading to the lungs); pneumonia; gonorrhoea (a sexually transmitted disease); and infections of the skin, ear, sinuses, throat, tonsils, and urinary tract  
**Pack :** 1x10, 3x10 and 10x10 Blister



### CEFPODOXIME PROXETIL AND CLAVULANATE POTASSIUM TABLETS

**Composition :** CEFPODOXIME PROXETIL 100/200 MG AND CLAVULANATE POTASSIUM 62.5/125 MG TABLETS  
**Indication :** Cefpodoxime proxetil+Clavulanic acid is used for the treatment of bacterial infections of the skin, ear, respiratory tract, urinary tract, lungs, throat and tonsil and sexually transmitted diseases  
**Pack :** 1x10, 3x10 and 10x10 Blister



### GLOXIM SUSPENSION

**Composition :** CEFIXIME FOR ORAL SUSPENSION USP 50 MG / 100 MG  
**Indication :** For use in the treatment of the following infections when caused by susceptible strains of the designated microorganisms: (1) Uncomplicated UTI caused by E. Coli and P.mirabilis, (2) Otitis media caused by H. influenzae (beta-lactamase +ve and -ve strains), M. Catarrhalis (most of which are beta-lactamase +ve) and S. pyogenes, (3) Pharyngitis and tonsillitis caused by S. pyogenes, (4) Acute bronchitis and acute exacerbations of chronic bronchitis caused by S. pneumoniae and H. influenzae (beta-lactamase +ve and -ve strains) and (5) Uncomplicated gonorrhoea (cervical/urethral) caused by N. gonorrhoeae (penicillinase and non-penicillinase producing strains).  
**Pack :** 30, 50, 60 & 100 ml Bottle



### GLOROXIM SUSPENSION

**Composition :** CEFUROXIME AXETIL FOR ORAL SUSPENSION USP 125/250 MG/ 5 ML  
**Indication :** For the treatment of the following infections (skin, UTI, ENT) caused by; S. pneumoniae, H. influenzae, staphylococci, S.pyogenes (Group A beta-hemolytic streptococci), E.coli, P.mirabilis, Klebsiella sp, coagulase-negative staphylococci and Streptococcus pyogenes.  
**Pack :** 30, 60 & 100 ML



### CEFDINIR CAPSULES

- Composition :** Cefdinir Capsules USP 300 mg  
**Indication :** Cefdinir is an antibiotic indicated in the treatment Community-Acquired Pneumonia, Acute Maxillary Sinusitis, otitis media, strep throat, and cellulitis. Uncomplicated infections caused by staphylococcus aureus.  
**Pack :** 1 x 10, 3 x 10 and 10x10 Blister



### GLOCEFIM

- Composition :** CEFALEXIN ORAL SUSPENSION 125/250 MG  
**Indication :** Cefalexin is a semi synthetic cephalosporin antibiotic for oral administration. Cefalexin is indicated in the treatment of the following infections: Respiratory tract infections; otitis media; skin and soft tissue infections; bone and joint infections; genito-urinary infections, including acute prostatitis and dental infections.  
**Pack :** 30 & 60 ML Bottle



### GLOCEF DINIR

- Composition :** CEFDINIR FOR ORAL SUSPENSION USP 125 MG  
**Indication :** Cefdinir is an antibiotic indicated in the treatment Community-Acquired Pneumonia, Acute Maxillary Sinusitis, otitis media, strep throat, and cellulitis. Uncomplicated infections caused by staphylococcus aureus  
**Pack :** 50 ml Bottle



### CEFADROXIL AND CLAVULANATE POTASSIUM FOR ORAL SUSPENSION

- Composition :** CEFADROXIL AND CLAVULANATE POTASSIUM FOR ORAL SUSPENSION 100+62.5/250+62.5 MG  
**Indication :** CEFADROXIL+CLAVULANIC ACID belongs to the group of medicines called antibiotics used to treat various bacterial infections such as urinary tract, skin, chest, and throat infections. Bacterial infections are caused due to the multiplication of harmful bacteria inside or on the body.  
**Pack :** 30ML



## CEFIXIME 100 AND CLAVULANATE POTASSIUM FOR ORAL SUSPENSION

- Composition :** CEFIXIME AND CLAVULANATE POTASSIUM FOR ORAL SUSPENSION  
**Indication :** 100+62.5 MG  
 Cefixime+Potassium Clavulanate / Clavulanic acid is used to treat severe bacterial infections of the urinary tract, ear, lungs, throat, tonsil and skin or infections like typhoid fever or gonorrhoea  
**Pack :** 30 ML



## CEFPODOXIME PROXETIL FOR ORAL SUSPENSION

- Composition :** CEFPODOXIME PROXETIL FOR ORAL SUSPENSION 50/100 MG  
**Indication :** CEFPODOXIME (sef podo OX eem) is a cephalosporin antibiotic. It treats some infections caused by bacteria. It will not work for colds, the flu, or other viruses. This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions  
**Pack :** 30 ML



## CEFPODOXIME AND CLAVULANATE POTASSIUM FOR ORAL SUSPENSION

- Composition :** CEFPODOXIME AND CLAVULANATE POTASSIUM FOR ORAL SUSPENSION 100+62.5 MG  
**Indication :** Cefalexin is a semi synthetic cephalosporin antibiotic for oral administration. Cefalexin is indicated in the treatment of the following infections: Respiratory tract infections; otitis media; skin and soft tissue infections; bone and joint infections; genito-urinary infections, including acute prostatitis and dental infections.  
**Pack :** 60 ML



## CEFADROXIL FOR ORAL SUSPENSION

- Composition :** CEFADROXIL FOR ORAL SUSPENSION USP 125/250 MG  
**Indication :** For the treatment of the following infections (skin, UTI, ENT) caused by; S. pneumoniae, H. influenzae, staphylococci, S.pyogenes (Group A beta-hemolytic streptococci), E.coli, P.mirabilis, Klebsiella sp, coagulase-negative staphylococci and Streptococcus pyogenes.  
**Pack :** 30 ML



## CEFUROXIME + CLAVULANATE POTASSIUM

**Composition :** CEFUROXIME + CLAVULANATE POTASSIUM POWDER FOR ORAL SUSPENSION

**Indication :** For the treatment of the following infections (skin, UTI, ENT) caused by; *S. pneumoniae*, *H. influenzae*, staphylococci, *S. pyogenes* (Group A beta-hemolytic streptococci), *E. coli*, *P. mirabilis*, *Klebsiella sp.*, coagulase-negative staphylococci and *Streptococcus pyogenes*.

**Pack :** 100 ML



## GLOMOXX

**Composition :** AMOXICILLINE POWDER FOR ORAL SUSPENSION 125/250 MG/5ML

**Indication :** Amoxicillin Oral Suspension is indicated for the treatment Acute bacterial sinusitis, Acute Otitis media, Acute streptococcal tonsillitis and pharyngitis, Acute exacerbation of chronic bronchitis, Community acquired pneumonia, Acute cystitis, Asymptomatic bacteriuria in pregnancy

**Pack :** 100 ML



## AMOXICILLINE + CLAVULANATE POTASSIUM

**Composition :** AMOXICILLINE + CLAVULANATE POTASSIUM POWDER FOR ORAL SUSPENSION

**Indication :** Amoxicillin Oral Suspension is indicated for the treatment Acute bacterial sinusitis, Acute Otitis media, Acute streptococcal tonsillitis and pharyngitis, Acute exacerbation of chronic bronchitis, Community acquired pneumonia, Acute cystitis, Asymptomatic bacteriuria in pregnancy

**Pack :** 100 ML



## GLOAMP CLX SUSPENSION

**Composition :** AMPICILLIN AND CLOXACILLIN FOR ORAL SUSPENSION 250+312.5 MG

**Indication :** Ampicillin & Cloxacillin dry syrup is used to treat many bacterial infections like tonsillitis, bronchitis, pneumonia, gonorrhoea, and infections of the ear, nose, throat, skin and urinary tract.

**Pack :** 100 ml bottle



### GLOAMP

**Composition :** AMPICILLIN ORAL SUSPENSION BP 125 OR 250 MG/ 5ML

**Indication :** Ampicillin 125mg Dry Syrup is an antibiotic medicine with versatile actions. It can treat a multitude of bacterial infections including those of the ears, eyes, nose, tonsils, tooth, throat, lungs, skin, bone, heart, gastrointestinal tract, and urinary tract.

**Pack :** 60ml, 100 ml



### GLOAMP CLX

**Composition :** AMPICILLIN AND CLOXACILLIN CAPSULES

**Indication :** Ampicillin + Cloxacillin is used to treat patients with bacterial infections. It is a semi-synthetic derivative of the drug penicillin. It is used to treat urinary tract and respiratory tract infections, meningitis, gonorrhoea and infections of the stomach or intestine.

**Pack :** 1x10, 10x10 Blister



### GLOBETA

**Composition :** AMOXICILLIN AND POTASSIUM CLAVULANATE TABLETS BP 625/1000 MG

**Indication :** The combination of amoxicillin and clavulanic acid is used to treat certain infections caused by bacteria, including infections of the ears, lungs, sinus, skin, and urinary tract.

**Pack :** 2x7, 1x10 Blister



### G-CLOX

**Composition :** CLOXACILLIN SODIUM CAPSULES 250/500 MG

**Indication :** Cloxacillin is used to treat a wide variety of bacterial infections. This medication is a type of penicillin antibiotic. It works by stopping the growth of bacteria. This antibiotic treats only bacterial infections.

**Pack :** 10x10 Blister




### CEFIXIME + OFLOXACIN + LB

**Composition :** CEFIXIME + OFLOXACIN + LB TABLETS

**Indication :** Cefixime + Ofloxacin + LB Tablets are indicated for the treatment of various bacterial infections. This combination is used to: Treat infections caused by susceptible bacteria including those of the respiratory tract, urinary tract, and skin. Provide a broad-spectrum antibacterial effect due to the combined actions of cefixime (a cephalosporin antibiotic), ofloxacin (a fluoroquinolone antibiotic), and LB (lactic acid bacteria, which helps to restore gut flora disrupted by antibiotic therapy).

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister



## CEFIXIME + CLAVULANIC ACID

**Composition :** CEFIXIME + CLAVULANIC ACID TABLETS 200+125 MG

**Indication :** Cefixime + Clavulanic Acid Tablets are indicated for the treatment of infections caused by susceptible bacteria. This combination is used to treat a variety of bacterial infections, including those of the respiratory tract, urinary tract, and skin.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister




## FLUCLOXACILLIN

**Composition :** FLUCLOXACILLIN CAPSULES 250/500 MG

**Indication :** Flucloxacillin Capsules are indicated for the treatment of infections caused by susceptible Gram-positive bacteria. Specifically, they are used to treat infections such as skin and soft tissue infections, including impetigo, cellulitis, and abscesses. Manage infections of the respiratory tract, bone, and joints, as well as endocarditis and osteomyelitis, when caused by Staphylococcus aureus or other susceptible organisms.

**Pack :** 1 x 10, 3 x 10 and 10 x 10 Blister





## TASTE MASKED & DIRECT COMPRESSION GRANULES

### Pharmaceutical Formulation Intermediates DC (Direct Compression Granules)

#### What is a DC?

DC is the intermediate product between an API and a Finished Dosage. The proportion that DC could be outsourced as per customer requirement without much technology risk at their end. Customer will get ready granules with proven quality and data of stability to meet International standard.

#### Why Direct Compression Granules ?

• The DC granules are taken directly from the drums to a hopper, where they are directly compressed into tablets. • It decreases the overall cost of the manufacturers and which finally results in decreased cost of product to the end consumer.

#### Benefits of Direct Compression Granules.

- Saves Process Time
- Simplifies Supply Chain Management
- Reduces Capital Expenditure
- Reduces Vendor Development Cost
- Lower Testing Cost
- Regulatory Benefits
- Develops Dry-Blending with other APIs

**Desired properties :** Leading to finished dosage manufacturers getting favorable outcomes from their bioequivalence studies.

**Dry Blending :** We possess the capability for some Pharmaceutical Formulation Intermediates to be dry-blended with other APIs without the need for further granulation to obtain homogeneous mixtures.

We back our production line with EU GMP & WHO GMP manufacturing infrastructure. To support our Pharmaceutical Formulation Intermediate, we have prominent tie up with API manufacturer for DMF.

#### Direct Drum to Hopper product with Unique formula.

- Sustain released Direct Compression granules – Metformin SR suitable to 500 mg , 850 mg & 1 gm
- Tablets • Extended released • Rapid released
- Taste masking of formula for paediatric suspension

## PRODUCTS

Amlodipine DC 1.35%

Atorvastatin 6.66%, 13.33%

Ascorbic Acid DC 62.5 %

Aceclofenac DC 41.66 %

Azithromycin DC granules 60%

Betahistin DC 8/16/24%

Ciprofloxacin DC Granules 65%, 75%

Clopidogrel DC 26.78%

Doxycycline Pre-Mix Granules 50%

Glibenclamide DC 1.36%

Finasteride DC 4%

Fluconazole DC 50%

Furosemide DC 71%

Ibuprofen DC Granules 66% / 70% / 90%

Lisinopril DC 5%

Loperamide DC 1%

Metformin HCL DC Granules 90% / 95%

Metformin Hydrochloride SR 75%

Metronidazole DC granules 76.92%

Naproxen DC 66.66%

Ofloxacin DC 57.14% / 83.33 %

Paracetamol DC Granules 66% / 90% / 95%

Rosuvastatin 6.25%

Sevelamer Hydrochloride DC 75%, 84%

Sildenafil Citrate DC 17.24%

Tadalafil DC 3.44%

Tinidazole DC 66.66%

Alpha Ketoanalogue DC Granules

Ketoprofen DC 60 %

Other Product Available as per Request and Specification



PRODUCTS NAME	CATEGORY	PACK SIZE
Beclomethasone Cream BP	Corticosteroid	15 g Aluminum Tube
Betamethasone Dipropionate Cream BP 0.05% w/w	Corticosteroid	15 g Aluminum Tube
Betamethasone Dipropionate Cream IP 0.1%	Corticosteroid	15 g Aluminum Tube
Betamethasone Dipropionate, Clotrimazole, Gentamicin Sulphate Cream	Corticosteroid	15 g Aluminum Tube
Betamethasone Valerate, Gentamicin Sulfate, Tolnaftate and Iodochlorhydroxyquinoline Cream	Corticosteroid	30 g Aluminum Tube
Clindamycin Phosphate Gel	Topical Antibiotic	30 g Lami Tube
Clobetasol Propionate Cream BP	Topical Steroid	30 g Lami Tube
Clobetasol Propionate Cream USP	Topical Steroid	30 g Lami Tube
Clotrimazole and Beclomethasone Cream	Topical Steroid	15/30 g Alu. Tube
Clotrimazole Vaginal Cream BP 1% w/w	Topical Antifungal	30 g Aluminum Tube
Clotrimazole Vaginal Cream USP 1% w/w	Topical Antifungal	30 g Aluminum Tube
Clotrimazole Vaginal Cream USP 2% w/w	Topical Antifungal	30 g Aluminum Tube
Clotrimazole, Beclomethasone and Neomycin Cream	Topical Antifungal	30 g Lami Tube
Diclofenac Diethylamine, Methyl Salicylate, Oleum lini & Menthol Gel	NSAID	20/30/50 g Lami Tube
Diclofenac Gel	NSAID	15 g Aluminum Tube
Hydrocortisone Acetate Cream BP 1.0%w/w	Topical Steroid	20 g Lami Tube
Ketoconazole Cream 2%w/w	Topical Antifungal	30 g Aluminum Tube
Metronidazole Gel BP	Anti-infective	15 g Aluminum Tube
Miconazole Nitrate Cream BP	Anti-infective	
Precipitated Sulfur Cream	Anti-infective	25 g Lami Tube
Salicylic Acid Ointment	NSAID	
Silver Sulfadiazine And Chlorhexidine Gluconate Cream		30/50 g Alu. Tube
Silver Sulfadiazine Cream USP 1.0%w/w	Anti-infective	10/15 g Alu. Tube
Terbinafine Hydrochloride Cream 1% w/w	Topical Antifungal	



### CARDIOVASCULAR DRUGS

Amlodipine	5/10 mg
Amlodipine Besylate + Lisinopril	5+5 mg
Amlodipine + Valsartan + HCT	10 +160+25 mg
Apixaban	2.5/5 mg
Atorvastatin	10/20 mg
Atorvastatin + Ezetimibe	10+10 mg
Atorvastatin + Fenofibrate	10+160 mg
Bisoprolol fumarate	2.5/5/10 mg
Bosentan	62.5/125 mg
Captopril	25/50 mg
Candesartan	4/8/16 mg
Clopidogrel	75 mg
Clopidogrel + Aspirin	75+75 mg
Digoxin	250 mg
Enalapril	2.5/5/10/20 mg
Fenofibrate	100/160/200 mg
Furosemide	20/40/80 mg
Hydrochlorothiazide	12.5/25 mg
Irbesartan	150/300mg
Irbesartan + Hydrochlorothiazide	150+12.5 mg
Ivabradine	5/7.5 mg
Lisinopril	5/10/20 mg
Losartan + Amlodipine	50+5/100+10 mg
Losartan + Hydrochlorothiazide	50+12.5 mg
Losartan Potassium	25/50 mg
Methyldopa	250 mg
Nifedipine ER	5/10/20 mg
Metoprolol Tartrate	50/100 mg
Olmesartan Medoxomil	20/40 mg
Trimetazidine	35 mg
Perindopril	2.5/5/10 mg
Ramipril	5/10mg
Rivaroxaban	2.5/5/10/20 mg
Rosuvastatin	10/20mg
Rosuvastatin + Ezetimibe	20+10 mg
Rosuvastatin + Fenofibrate	10+160 mg
Sacubitril + Valsartan	24+26 mg
Simvastatin	5/10/20 mg
Simvastatin + Ezetimibe	10+10 mg
Telmisartan	20/40/80 mg
Telmisartan + Amlodipine	40+5/80+10 mg
Telmisartan + Hydrochlorothiazide	40+12.5 mg
Valsartan	80/160/320 mg
Valsartan + Hydrochlorothiazide	160+12.5mg
Verapamil	40/80 mg

### ANTI PSYCHOTIC & CNS

Carbamazepine	100/200/500 mg
Citalopram	20/40 mg
Citicoline	500 mg
Escitalopram	5/10/20 mg
Gabapentin	100/300/400/600/750/800/900 mg
Levetiracetam	250/500/750/1000 mg
Oxcarbazepine	150/300/600 ER mg
Paroxetine	20/40 mg
Quetiapine	25/50/100/200/300/400 mg
Vigabatrin	500 mg

### STEROIDAL ANTI INFLAMMATORY

Betamethasone	0.5 mg
Deflazacort	1/6/18/30 mg
Dexamethasone	0.5/ 1 mg/4 mg/ 8 mg
Prednisolone	5/10 mg

### ANTI MALARIAL

Artemether + Lumefantrine	20+120/40+240/ 80+480 mg
Artemether + Lumefantrine Dry Sus.	60 ml
Combipack of Artesunate & Amodiaquine	160+320 mg
Dihydroartemisinin + Piperaquin	40+320 mg
Dihydroartemisinin + Piperaquine Dry Sus..	80+640 mg
Quinine Sulphate	300/600 mg
Artemisinin + Piperaquine	62.5+375 mg

### NEPHROLOGY & UROLOGY

Allopurinol	100/200/300/500 mg
Alpha Ketoanalogues & Essential Amino Acids	15 g
Calcium Polyestylene Sulfonate	15 g
Cinacalcet	30/60/90 mg
Deferasirox	100/400/250/500 mg
Febuxostat	40/80 mg
Finasteride	1/5 mg
Sevelamer Carbonate / Hydrochloride	400/800 mg
Tamsulosin + Dutasteride	400+500 mcg
Tamsulosin HCL	400 mcg

### ARV PRODUCT

Abacavir	300 mg
Abacavir + Lamivudine	300+150/600+300 mg
Abacavir, Dolutegravir & Lamivudine	600+50 +300 mg
Acyclovir	200/400 mg
Atazanavir	100/150 mg
Darunavir	400/600/800 mg
Dolutegravir	50 mg
Dolutegravir & Lamivudine	50+300 mg
Efavirenz	200/600 mg
Efavirenz & Lamivudine & Tenofovir	600+300+300 mg
Emtricitabine + Tenofovir	300+200 mg
Emtricitabine & Tenofovir Alafenamide	200+10 mg
Lamivudine	100/150/300 mg
Lamivudine + Zidovudine	150+300 mg
Favipiravir	150/300 mg
Nevirapine	200 mg
Raltegravir	400/600 mg
Ritonavir	100 mg
Sofosbuvir	150/200/300/400 mg
Tenofovir + Lamivudine	300+300 mg
Tenofovir + Lamivudine + Dolutegravir	300+300+50 mg
Tenofovir	300 mg
Lopinavir & Ritonavir	200+50/400+100 mg
Valacyclovir	500/1000 mg
Valganciclovir	450 mg
Zidovudine	100/300 mg

### VITAMINS AND MINERALS

Ascorbic Acid Tablets	100/500mg
Calcium Carbonate (500 mg) With Vitamin D3	---
Diosmin & Hesperidin	50+450 mg
Folic Acid	5 mg
Ferrous Gluconate	324 mg
Multivitamin & Multimineral	---
Potassium Chloride ER	650/750 mg

# FINISHED FORMULATION



## Dedicated Oral Solid Dosage Facilities

### IMMUNO SUPPRESANTS

Azathioprine Tablets	50/100 mg
Everolimus Tablets	0.25/0.50/0.75/5/10 mg
Mycophenolate Mofetil Tablets	250/500 mg
Mycophenolate Sodium Tablets	180/360 mg
Sirolimus Tablets	1 mg
Tacrolimus Capsules	0.5/1/5 mg

### GASTROLOGY

Bisacodyl	5 mg
Cimetidine	200/400 mg
Compound magnesium ( Antacid)	100/400 mg
Domperidone	10 mg
Esomeprazole & Domperidone	20 /10 mg
Esomeprazole Mag. Di & Trihydrate	20/40 mg
Hyoscine Butylbromide	10 mg
Famotidine	20/40 mg
Lansoprazole	30 mg
Loperamide	2 mg
Magnesium	400 mg
Misoprostol	200 mcg
Omeprazole	20/40 mg
Omeprazole + Domperidone	20+10 mg
Pysillium Husk Powder	---
Pantoprazole	20/40 mg
Pantoprazole + Domperidone	40+10 mg
Rabeprazole + Domperidone	20+10 mg
Rabeprazole Sodium	10/20/40 mg
Sennoside Calcium	8.6 mg
Ursodeoxycholic Acid	150/250/300/500 mg

### ANTI ALLERGIC/ANTI HISTAMINE

Betahistine HCL	8/16/24 mg
Chlorpheniramine	8/4 mg
Cetirizine	10 mg
Desloratadine	5/10 mg
Diphenhydramine	25/50 mg
Fexofenadine Hcl	60/120/180 mg
Levocetirizine	5/10 mg
Levocetirizine hcl + Montelukast	5+10 mg
Loratadine	10 mg
Montelukast	4/5/10 mg
Ondansetron	4 /8 mg
Paracetamol (500 mg) + Phenylephrine hydrochloride (10 mg) + Chlorpheniramine (2 mg)	
Paracetamol, Dextromethorphan Hydrobromide, Phenylephrine Hydrochloride & CPM Tablets Kit	

### ANTIFUNGAL / ANTHELMINTICS

Albendazole	400 mg
Clotrimazole vaginal	100 mg
Diloxanide furoate	500 mg
Fluconazole	50/150/200/400 mg
Griseofulvin	250/500 mg
Itraconazole	100/200 mg
Ketoconazole	200 mg
Mebendazole	200 mg
Nitazoxanide	500 mg
Metronidazole	200/400/500 mg
Secnidazole	500 mg/1gm

### ERECTILE DYSFUNCTION

Sildenafil citrate	50 / 100 mg
Tadalafil/ Tadalafil Chewble	5/10/20 mg
Vardenafil	5/10/20 mg

### ANALGESIC & ANTISPASMODICS

Aceclofenac	100 mg
Aceclofenac + Paracetamol	100+500 mg
Celecoxib	100/200mg
Diclofenac Potassium + Paracetamol	50 +500 mg
Diclofenac Sodium	50/100 mg
Diclofenac Potassium	50/100 mg
Diclofenac Sodium + Vit. B1+ Vit B6 + Vit B12	
Diclofenac Potassium, Chlorzoxazone & Paracetamol	100+500+250 mg
Dexketoprofen	25 mg
Etoricoxib	60/90/120 mg
Ibuprofen	200/400/600 mg
Ibuprofen + Paracetamol	10+325 mg
Ibuprofen + Paracetamol + Caffeine	200+ 325+30 mg
Indometacin	25/50/100 mg
Mefanamic	250/500 mg
Meloxicam	7.5/15 mg
Naproxen	500 mg
Nebumetone	500/750 mg
Nebumetone + paracetamol	500+500 mg
Nimesulide	50/100 mg
Nimesulide + Paracetamol	400+325 mg
Orphenadrine + Paracetamol	35+450 mg
Paracetamol + Caffeine	500+30 mg
Paracetamol + Dicyclomine	500+20 mg
Paracetamol	325/500 mg
Pentoxifylline	400 mg
Piroxicam	20 mg
Serratiopeptidase	10 mg
Tapentadol Hcl	50/75/100 mg

### ANTI DIABETICS

Dapagliflozin	5/10 mg
Empagliflozin	10/25 mg
Glibenclamide	2.5/ 5 mg
Gliclazide/Gliclazide MR	40/80-MR 30/60 mg
Glimepiride	1/2/3/4 mg
Glibenclamide + Metformin	5/500 mg
Glimepiride + Metformin	2 /500 mg
Metformin	500/850/1000 mg
Pioglitazone HCL	15/30 mg
Sitagliptin + Metformin	50+500/100+1000 mg
Sitagliptin	25/50/100 mg
Vildagliptin	50/100 mg
Vildagliptin + Metformin	50+500/100+1000 mg

### ELECTROLYTES

Diocahedral Smectite Powder	3.0 g
Oral Rehydration Salts	27.9/20.5/4.2 gm
Oral Rehydration Salts and Zinc	20.5gm/20mg



### ANTI BACTERIAL

Azithromycin	100/250/500 mg
Azithromycin dry sup.	200/ 5 ml
Ciprofloxacin	250/500 mg
Ciprofloxacin + Tinidazole	500 + 600 mg
Clarithromycin	250/500 mg
Clindamycin	150/300/ 600 mg
Cotrimoxazole	480/960 mg
Doxycycline	50/75/100 mg
Erythromycin Stearate	250/500 mg
Fluconazole	150/200 mg
Fluconazole Suspension	50/5 ml
Griseofulvin	250/500 mg
Levofloxacin	250/500/750 mg
Lincomycin	250/500 mg
Moxifloxacin	400 mg
Norfloxacin	400mg
Ofloxacin	200/400 mg
Ofloxacin+Ornidazole	200 + 500 mg
Sparfloxacin	100/200/400 mg
Terbinafine	250 mg
Tinidazole	500 mg
Esomeprazole+Clarithromycin +Tinidazole	40+250+500 gm kit
Fluconazole+Azithromycin +Secnidazole	150+1+1gm kit
Omeprazole+Clarithromycin +Tinidazole	30+250+500 gm kit

### CEPHALOSPORINS

Cefixime Dispersible	50/100/200/400 mg
Cefixime Tablets	100/200/400 mg
Cefixime + Clavulanate Potassuim	31.25/62.50/125 mg
Cefixime + Ofloxacin	100/200 mg Tablets
Cefuroxime Axetil	125/250/500 mg
Cefuroxime Axetil + Clavulanate Potassium	250/500+125 mg
Cefpodoxime Proxetil Dispersible	100/200 mg
Cefpodoxime Proxetil Tablets	100/200 mg
Cefpodoxime Proxetil + Clavulanate Potassium	100/200+ 62.50/125 mg
Cefadroxil Dispersible	250 mg
Cefadroxil	500 mg
Cefadroxil and Clavulanate Potassium	500+125 mg
Cephalexin Dispersible	125/250 mg
Cephalexin	250/500 mg
Cefixime	100/200/400 mg
Cefalexin	BP 250/500 mg
Cefdinir	300 mg
Cefixime For Oral Suspension	50/100 mg/5 ml
Cefixime + Clavulanate Potassium For Oral Suspension	100 mg/62.50 mg Per 5ml
Cefuroxime Axetil For Oral Suspension	125/250 mg/5 ml
Cefpodoxime Proxetil For Oral Suspension	50/100 mg/5 ml
Cefpodoxime + Clavulanate Potassium For Oral Suspension	100 mg/62.50mg Per 5ml
Cefadroxil For Oral Suspension	125/250 mg/5ml
Cefadroxil and Clavulanate Potassium For Oral Suspension	250 mg/62.50 mg/5ml
Cefalexin For Oral Suspension	125/250 mg/5ml

### BETALACTUM

Amoxicillin + Clavalunic acid	250+125 mg/500+125/875+125 mg
Amoxicillin + Clavalunic acid Suspension	125+31.5 mg/5 ml,200+28.5 mg/5 ml
Amoxicillin + Potassium Clavulanate Suspension	250+62.5 mg/5 ml
Amoxicillin + Potassium Clavulanate Suspension	400+57 mg/5 ml
Amoxicillin	250/500 mg
Amoxicillin Suspension	125+250 mg/5 ml
Amoxicillin	250 mg
Ampicillin + Cloxacillin	250+250 mg
Ampicillin dispersible	250 mg
Cloxacillin	500 mg

### ONCOLOGY

Abiraterone	250/500 mg
Alectinib Hydrochloride	150 mg
Anastrozole	1 mg
Apalutamide	60 mg
Aprepitant	60/80 mg
Axitinib	1/5 mg
Baricitinib	4 mg
Bicalutamide	50/150 mg
Binimetinib	15 mg
Busulphan	2 mg
Cabozantinib	20/80 mg
Capecitabine	150/500 mg
Cyclophosphamide	25/50 mg
Cyclosporine	25/50/100 mg
Dasatinib	20/50 mg
Enzalutamide	40 mg
Erlotinib	100/150 mg
Estramustine	140 mg
Etoposide	50/100 mg
Eltrombopag	25/50 gm
Exemestane	25 mg
Fingolimod	0.5 mg
Fludarabine Phosphate	10 mg
Flutamide	250 mg
Gefitinib	250 mg
Hydroxyurea	500 mg
Ibrutinib	140 mg
Imatinib Mesylate	100/400 mg
Larotrectinib	25/100 mg
Lenalidomide	5/10/15/25 mg
Lenvatinib	4/10 mg
Letrozole	2.5 mg
Lorlatinib	100 mg
Megestrol Acetate	160 mg
Melphalan	2/5 mg
Mercaptopurine	50 mg
Methotrexate	2.5/5/10 mg
Nintedanib	100/150 mg
Niraparib	100 mg
Olaparib	50/150 mg
Osimertinib	80 mg
Palbociclib	75/100/125 mg
Pazopanib	200/400 mg
Ponatinib	15/45 mg
Procarbazine	50 mg
Rucaparib	200/250/300 mg
Sorafenib Tosylate	250 mg
Sunitinib	25/50 mg
Tamoxifen	10/20 mg
Temozolomide	20/100/250 mg
Thalidomide	50/100 mg
Tucatinib	150 mg
Venetoclax	100 mg



# GLOBAL PRESENCE



## NORTH AMERICA



United State of America



European Union

## AFRICA



Nigeria



Kenya



Ivory Coast



Tanzania



Ghana



Uganda



Angola



Togo



Ethiopia



Malawi



Rwanda



South Sudan



Cameroon



Madagascar



Liberia



Libya



Senegal



Burundi



Namibia



Eritrea

## LATIN AMERICA / CENTRAL AMERICA



Ecuador



Chile



Paraguay



Colombia



Venezuela



Peru



Guyana



Bolivia



Suriname



Honduras



Cuba



Guatemala

## MIDDLE EAST



Yemen



Iraq



Iran



Syria

## ASIA



Sri Lanka



Myanmar



Philippines



India



Cambodia



Vietnam



Afghanistan

## CIS



Tajikistan



Kyrgyzstan



Uzbekistan



Serbia



Ukraine



Russia



Moldova



Turkmenistan

## CERTIFICATIONS & AWARDS



Dun & Bradstreet



EU GMP



GMP



WHO-GMP



INDIAN PHARMA AWARD

Globela® Laboratories

Works :  
Plot No. 5536, Road No. 55, G.I.D.C Estate,  
Sachin, Surat-394230, Gujarat, (India).  
www.globelapharma.com

Globela® Industries

Works :  
Plot No. C-198, G.I.D.C., Saykha,  
Bharuch-392140, Gujarat, India.  
www.globela.in

Globela® Pharma LLC

Office :  
P.O. BOX 8236, JERSEYCITY, NJ-07308  
Cell : +1(917) 724-1489  
Email : priyank@globelapharma.com